

FOR CATTLE OILERS — BACK RUBBERS OR FACE RUBBERS —
To control horn flies on dairy and beef cattle - dilute 1 pint Roberts Ciodrin Emulsion Concentrate with 2½ gallons diesel or fuel oil or other base oil. Recharge oiler, back rubber or face rubber as needed. Do not spray animals with oil solution.

PRECAUTIONS IN USING

Do not contaminate foodstuffs, or feed. Do not apply to calves under 6 months of age. Brahman cattle should not be treated as they may also show hypersensitivity to organic phosphates. During prolonged exposure in spraying and loading operations wear clean synthetic rubber gloves and a mask or respirator of a type found suitable by the U.S. Department of Agriculture for protection against Ciodrin Concentrate Insecticide. Use this material only for recommended purposes and at recommended dosage.

KEEP COVER TIGHTLY CLOSED WHEN NOT IN USE

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not re-use container. Destroy when empty.

ROBERTS LABORATORIES, INC. makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

Manufactured by
ROBERTS LABORATORIES, INC.
Rockford, Illinois 61101

USDA Reg. No. 523-46



**CIODRIN[®]
CONCENTRATE**

A RESIDUAL LIVESTOCK
AND BARN SPRAY
MIX AND SPRINKLE
OR SPRAY

ACTIVE INGREDIENTS:

Dimethyl phosphate of alpha-methylbenzyl
3-hydroxy-cis-crotonate 14.40%
Aromatic petroleum hydrocarbons 75.06%

INERT INGREDIENTS: 10.54%

WARNING: KEEP OUT OF REACH OF CHILDREN

HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN. Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Keep away from heat and open flame. See side panel for additional precautions.

NET CONTENTS ONE QUART

DIRECTIONS

FOR DAIRY CATTLE — To control horn flies, and stable flies, face flies, house flies — dilute 1 pint Roberts Ciodrin Emulsion Concentrate with 2 gallons water. Apply 1 pint of diluted spray per adult animal once each week, or dilute 1 pint Roberts Emulsion Concentrate with 1 gallon of water and apply 1 - 2 oz. per adult animal daily, or as needed.

TO CONTROL LICE AND SOME TICKS (such as the lone star tick or winter ticks) - dilute 1 pint Roberts Ciodrin Emulsion Concentrate with 6 - 12 gallons water. Apply 1 - 2 pints spray per adult animal. Repeat as needed, not oftener than once per week.

FOR BEEF CATTLE — To control horn flies, stable flies, face flies, house flies and ticks - dilute 1 pint Roberts Ciodrin Emulsion Concentrate with 6 - 12 gallons water. Apply 1-2 pints spray per adult animal. Repeat as necessary to maintain control, but not oftener than once weekly.

AIDS IN THE REDUCTION OF FLY POPULATIONS: In dairy barns, beef barns, hog houses, other farm buildings - Mix 1 pint Roberts Ciodrin Emulsion Concentrate with 6 - 12 gallons water. Apply to ceilings, walls, doorways, windows, or other areas where flies congregate. On refuse areas, soak surfaces with spray and repeat as needed. **DO NOT USE IN POULTRY HOUSES.**

SHEEP, GOATS, SWINE, AND HORSES — To control lice and the lone star tick or winter ticks on sheep and goats, lice on swine, lice, ticks, horn flies, and stable flies on horses, dilute 1 pint Roberts Ciodrin Emulsion Concentrate with 6-12 gallons water and spray animals thoroughly. Apply 1-2 pints of spray to larger animals, and use proportionately less for smaller animals. Repeat as needed, but not oftener than once weekly.