

INSTRUCTIONS

RATS Place 1/2 ounce Throw-Pax next to burrows, in runways, or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 15 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

MICE Bait stations at intervals of 8 to 12 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days.

This rodenticide contains **WARFARIN**, an anticoagulant chemical that reduces the clotting ability of the blood and upon repeated feedings, causes fatal hemorrhage in rats and mice. It also contains an antibacterial agent which inhibits bacteria that produce vitamin K, an antidote for **WARFARIN**.

Achieve better results with proper baiting. Place baits along runways, behind boards, against wall. Keep bait dry.



CAUTION

Baits should be placed in areas inaccessible to children, pets, wildlife and domestic animals or in tamper proof bait boxes.

If swallowed by humans, domestic animals, wildlife or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K, combined with blood transfusions are indicated as in the case of hemorrhage caused by overdose of bishydroxycoumarin.

DO NOT REUSE CONTAINER.

DESTROY WHEN EMPTY.

THROW-PAX PELLETTED RAT BAIT

MFD. BY

ROBERTS LABORATORIES, Rockford, Illinois

EPA REG. NO. 523-33AA



R-30

10
copies

ACCEPTED
12-28-71
12-523-48
UNDER THE FEDERAL INSECTICIDE
ACT FOR USE AS AN INSECTICIDE
FOR CONTROL OF PESTS. SUBJECT
TO ATIA REQUESTS.

RODEX™

WATER SOLUBLE RAT AND MOUSE KILLER

CONCENTRATE
(Use Wet or Dry)

Makes One Gallon or 2½ lbs. Dry Bait

ACTIVE INGREDIENTS:	BY WEIGHT
*Sodium Salt of 2-Diphenylacetyl-1,3 indandione	0.106%
INERT INGREDIENTS99.894%
TOTAL	100.000%

*Equivalent to 0.1% by weight of 2-Diphenylacetyl-1,3 indanedione.

CAUTION
KEEP OUT OF REACH OF CHILDREN
See Below for Additional Precautions

DIRECTIONS

TO PREPARE LIQUID BAITS. Dissolve 1 tablespoonful RODEX in one quart of water.

Expose in glass water founts in protected stations. Set fount on a clean level surface where rats and mice are accustomed to feed. Place baits along walls, in corners and in concealed locations where rats and mice can feed undisturbed. Discard at once water in which mold or filth is noted. Keep fount clean and well supplied at all times. Do not place fount where freezing may occur.

TO PREPARE DRY CEREAL BAITS. Thoroughly mix one part by weight RODEX with nineteen parts of dry cereal grain.

Place in shallow trays in protected bait stations in areas where rats and mice feed. Place baits along walls, in corners and in concealed locations where feed-