

CAUTION

Avoid inhaling dust. Avoid contamination of foodstuffs and feed. Avoid contact with the skin and eyes. May be absorbed through skin and dust may irritate nasal membranes. Wash thoroughly after handling. Do not apply spray or dust within 30 days of slaughter. Do not dip within 60 days of slaughter. Do not reuse container. Destroy when empty.

NOTICE OF WARRANTY

Roberts Laboratories, Inc., makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statement on this label.

ROBERTS LABORATORIES, INC.
ROCKFORD, ILLINOIS



**HOG MANGE
POWDER
WET OR DRY**

ACTIVE INGREDIENTS	By Wt.
Gamma Isomer of Benzene	
Hexachloride from Lindane	1.00%
INERT INGREDIENTS	99.00%
TOTAL	100.00%

**CAUTION KEEP OUT OF THE
REACH OF CHILDREN**

Apply only as specified on this label.
See side panel for additional precautions.
Net Contents 1 lb.

DIRECTIONS FOR USE

For the control of sarcoptic mange, and as an aid in the control of certain ticks and lice on hogs, especially in the winter when the use of a dip is not advised. Simply disperse a visible coating of the powder over the animal to be treated by means of a dust gun or shaker can. Treat the animal and its quarters. If it is desired to use this treatment in a wet form, simply suspend the powder in water at the rate of one half pound of powder to each gallon of water. Spray areas of heavy infestation or Dip if preferred. Do not treat sows within 2 weeks of farrowing nor for at least 3 weeks after. Do not spray or dip animals in cold stormy weather. Avoid treating overheated or sick animals. Do not treat animals under 3 weeks of age.

MADE IN U.S.A.
U.S.D.A. REGISTRATION #523-10