

(1 of 2)

**DIRECTIONS FOR USE
HOUSEFLIES**

Residual Wall Sprays: For the control of houseflies including resistant strains, in dairy barns, hog pens, calf barns, poultry houses and other farm buildings, apply a 1% residual spray to the ceilings, walls, stanchions, etc. Prepare the spray by mixing 1 pint Flygon 2-E in 3 gallons of water. Thoroughly wet all fly resting areas to the point of run-off. One gallon of spray will cover 500 to 1,000 square feet of surface. Repeat applications should be made when necessary.

Remove all animals from buildings when applying residual wall sprays.

Flygon 2-E is acceptable for use as a residual-type insecticide in and around federally inspected meat packing plants except where meat food products are processed or handled.

Flygon 2-E controls flies up to 8 weeks or longer.

For Localized Housefly Control: apply a spray containing 1/2 pint of Flygon 2-E in 5 quarts of water with a knapsack or similar type sprayer to areas frequented by flies, such as doorways, around windows, etc.

Repeat applications should be made when necessary. Good sanitation is a necessary part of any effective fly control program.

Maggot Sprays: For the control of housefly maggots, mix 1/2 pint of Flygon 2-E in 5 quarts of water and apply as a coarse spray or with a sprinkling can to fly-breeding areas, such as poultry droppings in caged-layer houses, garbage dumps and manure piles. Repeat application as additional manure or garbage is added.

General Outside Use:

For the control of houseflies around homes and recreation areas, garbage cans, animal quarters, food-processing plants, warehouses, loading docks and refuse areas, thoroughly spray exposed surfaces, such as walls, fences, garbage and refuse containers with 1 pint of Flygon 2-E in 3 gallons of water. Repeat applications should be made when necessary.


DO NOT Contaminate Feed and Foodstuffs, Drinking Fountains, Litter and Feed Troughs.

DO NOT use in Milk-Processing Rooms, Including Milk Houses and Milk Storage Rooms.

DO NOT use in Homes.



FLYGON 2-E



Systemic Insecticide

CONTROLS HOUSEFLIES

**For control of Insects and Mites on Ornamental Plants
Refer to Attached Leaflet**

ACTIVE INGREDIENTS:

Dimethoate (O-O-dimethyl S-[N-methyl-carbamoylmethyl] phosphorodithioate)	23.4%
Xylene	38.4%
INERT INGREDIENTS	38.2%
TOTAL	100.0%

WARNING

KEEP OUT OF REACH OF CHILDREN

See Side Panel for Antidote and Other Cautions

EPA Reg. No. 523-79
EPA Est. No. 523-IL-1

NET CONTENTS GALLONS

WARNING

VAPOR HARMFUL - HARMFUL OR FATAL IF SWALLOWED - CONCENTRATED MATERIAL CAUSES EYE IRRITATION - COMBUSTIBLE

Avoid breathing vapor or spray mist.

Avoid contact with skin or eyes.

Use only with adequate ventilation.

Wash thoroughly after handling.

Do not contaminate feed or feed products.

Do not use, pour, spill or store near heat or open flame.

In case of contact with eyes, flush eyes with plenty of water for at least 15 minutes; get medical attention.

NOTE TO PHYSICIANS: This product may cause cholinesterase inhibition. Atropine is Antidotal.

DO NOT STORE BELOW FREEZING TEMPERATURES.

Rinse spills on outside of bottle after use. When empty, wash thoroughly with water containing a household detergent. Recap and discard bottle. Never re-use.

This product is toxic to birds and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTICE OF WARRANTY

ROBERTS LABORATORIES, makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

Manufactured by
ROBERTS LABORATORIES
ROCKFORD, IL 61101

ITU

(2 of 2)

The uses listed below are foliar sprays unless otherwise indicated.


Ornamental Plants Outdoor Plants only	Pest Controlled	Teaspoonful per 1/2 Gal. Water	Tablespoonful per 3 Gals. Water
Azaleas	Lace Bugs, Leaf Miners Mites Tea Scale White Flies	1	2
Camellias	Aphids Camellia Scale Mites Tea Scale	1 Foliar Spray: Apply 2 sprays 6 weeks apart the first year followed by annual applica- tions soon after first growth begins in the spring. Soil Drench: Apply Cygon 2-E as a soil drench around the base of plants in early spring at the rate of 4 tablespoonsful per gallon of water per plant up to 6 feet tall. Increase this rate proportionately for larger plants.	2
Roses	Aphids Leafhoppers Mites Thrips	1	2
Birch	Aphids Leaf Miners	1/2 For leaf miners apply when leaves are ex- panded (about mid-May) and repeat in early July.	2
Boxwood	Leaf Miners Mealy Bugs Mites	1 For leaf miners, apply in spring when leaf miner flies first appear or in early sum- mer for control of larvae in the infested leaves.	2
Euonymus	Aphids Scale	2	4
Gladiolus	Aphids Thrips	1	2
Hemlock	Mites Scale	1	2
Juniper	Aphids Bagworms Midges Mites	2	4
Pine	Aphids Bagworms European Pine Shoot Moth Nantucket Pine Tip Moth Zimmerman Pine Moth	2	4



FLYGON 2-E

Systemic Insecticide

EPA Reg. No. 523-79
EPA Est. No. 523-IL



Manufactured by
ROBERTS LABORATORIES
ROCKFORD, IL 61101

OUTDOOR ORNAMENTAL PLANTS

This product is toxic to birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

FLYGON 2-E is generally effective in controlling aphids, thrips, leaf miners, scales, leafhoppers and mites. For proper timing of treatments for the control of specific pests on ornamental plants, consult your state agricultural experiment station or state agricultural extension service.

Make adequate sprays when pests appear or when damage is first observed. Do not overdose or overspray. Do not use on ornamental plants not listed. Do not use on any ornamental stock plants grown as a source of propagation material, such as cuttings, layers, root stocks or scions for grafting or budding. Do not use in spray mixtures containing oil. Do not use on plants growing in greenhouses.