



11-200
00110

2,4-D

LOW VOLATILE ESTER WEED KILLER

Emulsifiable concentrate for controlling certain weeds
in corn, small grains, pastures, and other non-crop
areas.

READ "WARNING NOTICE" BEFORE BUYING OR USING.
IF NOTICE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.
Keep out of reach of children.

CAUTION: MAY CAUSE SKIN IRRITATION.

Avoid contact with eyes, skin, or clothing.
Do not store near food, fertilizer, seeds, insecticides, or fungicides.
Do not contaminate water used for irrigation, domestic or spray purposes.
In case of fire, use water spray, foam, dry chemical or CO₂.
Harmful if swallowed.


ACTIVE INGREDIENTS:

Isocetyl ester of
2,4-Dichlorophenoxyacetic acid 69.7%


INERT INGREDIENTS:

36.3%
100.0%
Equivalent to 46.2% 2,4-D acid
or 4.0 pounds per gallon.

NET CONTENTS 1 U.S. GALLON



40



2,4-D

LOW VOLATILE ESTER

WEED KILLER

Emulsifiable concentrate for controlling certain weeds in corn, small grains, pastures, and other non-crop areas.

READ WARNING NOTICE BEFORE BUYING OR USING.
IF NOTICE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.
Keep out of reach of children.

CAUTION: MAY CAUSE SKIN IRRITATION.

Avoid contact with eyes, skin, or clothing.
Do not get on skin or clothes. Wash thoroughly with soap and water.
Do not get on skin or clothes. Wash thoroughly with soap and water.
Do not get on skin or clothes. Wash thoroughly with soap and water.

ACTIVE INGREDIENTS	
2,4-D (2,4-Dichlorophenoxyacetic acid)	69.7%
Inert ingredients	30.3%
100.0%	

NET CONTENTS 1 U.S. GALLON



WARNING—Extreme care must be used when applying 2,4-D to prevent injuring desirable plants and crops. 2,4-D should not be used in susceptible crops such as cotton, tomatoes, vegetables, grapes, fruit trees, soybeans, tobacco, or beans. Do not allow spray mist to drift on to such plants. Since even minute quantities of spray may cause severe injury. Use low pressures when spraying to produce coarse sprays which are less likely to drift. Do not spray on windy days. Esters of 2,4-D produce vapors after application which may injure susceptible plants in the vicinity.

Do not use on newly seeded lawns. Creeping grasses, such as bent, are injured by 2,4-D and should not be sprayed unless spot applications are made on weeds and incured turf injury can be tolerated. White clover, or other legumes, are damaged by 2,4-D.

Do not use on newly seeded lawns.

Do not use on newly seeded lawns.

Do not use on newly seeded lawns.

Do not use on newly seeded lawns.

Do not use on newly seeded lawns.

DIRECTIONS FOR USE

SMALL GRAINS (Wheat, Barley, Rye, Grasses)—Apply 1 pint per acre when weeds are in the first stage of growth. Do not apply after the milk stage. For perennial weeds such as Canada Thistle and Wild Mering, apply 1 pint per acre when weeds are in the first stage of growth. Do not apply after the milk stage. Do not apply to grain fields with corn 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.

CORN—For controlling susceptible annual or perennial weeds, apply not more than 1 pint per acre. If crop is 8 inches or taller, avoid application of 2,4-D to the growing point. Use drop nozzles and direct spray near base of the crop. Do not use after tassels appear. Different varieties may vary in susceptibility to 2,4-D. Do not use 2,4-D in corn grown for the production of hybrid seed. Consult your state agricultural experiment station or extension service weed specialist for specific recommendations. Avoid cultivating soon after treatment when corn may be brittle.

For pre-emergence treatment of weeds in corn, use 1 pint per acre applied as a broadcast spray 2 to 3 days before crop emergence. Do not use pre-emergence sprays in light sandy soils.

SORGHUM—Apply 1 pint per acre. Crop is most resistant to 2,4-D when it is in the first stage of growth. Use drop nozzles when crop is over 12 inches high. Do not use when plants are less than 12 inches high. Do not apply during flowering. Heavy rain may cause 2,4-D to settle in the soil. Consult your state agricultural experiment station or extension service weed specialist for specific recommendations.

PASTURES, LAWNS, GOLF COURSES AND SIMILAR TURF—Apply 1 pint per acre. Use drop nozzles. Do not use on newly seeded lawns. Do not use on lawns where creeping grasses are present. Do not use on lawns where white clover is present. Do not use on lawns where white clover is present. Do not use on lawns where white clover is present.

NON-CROP AREAS—Do not use on areas where crops are growing. Do not use on areas where crops are growing. Do not use on areas where crops are growing. Do not use on areas where crops are growing.

NOTE—For specific information on use, consult the label on the container of 2,4-D.

WARNING NOTICE

This product is a weed killer. It is not a herbicide. It is not a fertilizer. It is not a fungicide. It is not a bactericide. It is not a virus. It is not a nematode. It is not a mite. It is not a tick. It is not a flea. It is not a mosquito. It is not a fly. It is not a beetle. It is not a caterpillar. It is not a worm. It is not a slug. It is not a snail. It is not a crab. It is not a spider. It is not a scorpion. It is not a centipede. It is not a millipede. It is not a tick. It is not a flea. It is not a mosquito. It is not a fly. It is not a beetle. It is not a caterpillar. It is not a worm. It is not a slug. It is not a snail. It is not a crab. It is not a spider. It is not a scorpion. It is not a centipede. It is not a millipede.

MONSANTO COMPANY AGRICULTURAL DIVISION ST. LOUIS, MO., U.S.A.

