

GENERAL INFORMATION

Randox Granular controls these weeds:

Annual Bluegrass <i>Poa annua</i>	Great Foxtail (Wild Millet) <i>Setaria faberii</i>
Barnyardgrass (Watergrass) <i>Echinochloa crusgali</i>	Green Foxtail <i>Setaria viridis</i>
Cardinalweeds <i>Amaranthus palmeri</i>	Yellow Foxtail <i>Setaria glauca</i>
Carpetweed <i>Mollugo verticillata</i>	Pigweed (Redroot) <i>Amaranthus retroflexus</i>
Chenopod <i>Bromus setaceus</i>	Pigweed (Spiny) <i>Amaranthus spinosus</i>
Crabgrass <i>Digitaria sp.</i>	Purslane <i>Portulaca oleracea</i>
	Stinkgrass <i>Eragrostis canadensis</i>
Smooth top cyrtophol <i>Cyrtopholus sp.</i>	
Crab Purslane <i>Portulaca oleracea</i>	Quackgrass <i>Agropyron repens</i>
Barnyardgrass <i>Echinochloa crusgali</i>	Broadleaf Purslane <i>Portulaca oleracea</i>
Johnsongrass <i>Sorghum halepense</i>	Cocklebur <i>Xanthium sp.</i>
Notgrass <i>Cyperus sp.</i>	Field Bindweed <i>Convolvulus arvensis</i>

Calibrating Granular Equipment

For successful results, Randox Granular must be applied evenly at the proper rate per acre and frequent calibration of applying equipment in the field is recommended. Equipment which discharges granules through spouts and tubes may be calibrated for Randox Granular by using a wide

SPECIFIC CROP USE

CORN Hybrid Seed Corn, Field Corn, Sweet Corn, Popcorn


SORGHUM Grain, Forage. Do not use on seed production fields. Do not use on later plantings or similar fields where Sorghum is planted at the bottom of furrows.

BEANS East of Rocky Mountains only. Soybeans, Lima and Snap Beans

For **CORN**, **SORGHUM** and **BEANS** as shown above, apply Randox Granular when planting, or before either the crop or weeds appear above the soil surface. For over all treatment, use 20 pounds per acre. For band treatment, reduce the amount per acre in proportion to the area treated. Calculate according to the formula given on this label. In corn and soybeans use 25 lbs. per acre broadcast where recommended by competent local agricultural authorities.

ONIONS—Apply 30 pounds of Randox Granular per acre as an over all treatment just before onions emerge. If a windbreak crop is used, use barley instead of oats. Do not use windbreak barley for food or feed purposes. Heavy rains immediately after application may result in some stand reduction of onions.

Where seeded onions require treatment with certain insecticides or fungicides, or if fertilizer has been placed too close to the seed, onions may already be in danger of chemical injury. In such cases, it may be advisable to reduce the rate of Randox Granular to 20 pounds per acre over all for the first application, though this may reduce weed control somewhat.



MINIATURE FACSIMILE
Smallest Type size on package 12 pt

WARNING NOTICE

Monsanto Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. This product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as for example incompatibility with other products, the manner of its use or application, or the presence of other products or material in or on the soil or crop. MONSANTO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY. The exclusive remedy of the user or buyer and the limit of the liability of Monsanto Company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this notice which may not be varied by any verbal or written agreement.

Keep container tightly closed. Keep out of reach of children. Do not breathe dust or fumes. Wash hands thoroughly after use.


Use only as directed. Do not use on crops or plants not listed on the label. Do not use on crops or plants listed on the label if the crop or plant is not listed on the label.

Do not contaminate water, soil, or equipment. Do not use in water.

Do not use in water. Do not use in water.

Do not use in water. Do not use in water.

NET 50 LBS.



MS

RANDOX

Granular

18



Granular RANDOX

SELECTIVE HERBICIDE

For weed control at planting time in corn, sorghum, beans, onions, cabbage, sweet potatoes, tomatoes, sugar cane, celery, potatoes and castor beans. Do not use on other crops.

READ "WARNING NOTICE" BEFORE BUYING OR USING. IF NOTICE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.

KEEP OUT OF REACH OF CHILDREN

WARNING!
MAY CAUSE SKIN IRRITATION
OR EYE INJURY

Do not get in eyes or on skin
Avoid breathing dust
Do not take internally
In case the dust from the granules comes in contact with the eyes, flush with plenty of water for at least 15 minutes and get medical attention; for skin, wash with soap and water
Avoid contamination of seed, feed and foodstuffs.

50 LBS.

Use only according to label instructions.

ACTIVE INGREDIENT	
2-chloro-N,N-diallylacetamide	20%
INERT INGREDIENTS	
	80%
	100%

MONSANTO COMPANY • AGRICULTURAL DIVISION • ST. LOUIS, MISSOURI • U.S.A. COPYRIGHT 1960 MONSANTO COMPANY

MINIATURE FACSIMILE
Smallest Type size on package 12 point.


Granular RANDOX

NET 50 LBS.

RANDOX

Granular

NET 50 LBS.



Granular RANDOX

SELECTIVE HERBICIDE

For weed control at planting time in corn, sorghum, onions, cabbage, sweet potatoes, tomatoes, sugarcane, potatoes and castor beans. Do not use on other crops.

Sites

READ "WARNING NOTICE" BEFORE BUYING OR USING. IF NOTICE TERMS ARE NOT ACCEPTABLE, RETURN TO SELLER.

KEEP OUT OF REACH OF CHILDREN.

WARNING!
MAY CAUSE SKIN IRRITATION
OR EYE INJURY


Do not get in eyes or on skin.
Avoid breathing dust.
Do not take internally.
In case the dust from the granules comes in contact with the eyes, flush with plenty of water for at least 15 minutes and get medical attention; for skin, wash with soap and water.
Avoid contamination of seed, feed and foodstuffs.

Use only according to label instructions.

ACTIVE INGREDIENT
2,4-Diaminodinitrophenol
INERT INGREDIENTS

5

MONSANTO COMPANY • AGRICULTURAL DIVISION • ST. LOUIS, MISSOURI



Granular RANDOX