


NET 50 LBS. **T** **RAN DOX**  
Granular



Granular **T**  
**RAN DOX**

SELECTIVE HERBICIDE FOR CORN


For pre-emergence control of certain annual weeds in field corn, silage corn, sweet corn and popcorn.

REG. U.S. PAT. & TM. OFF. © 1977 MONSANTO CO. ST. LOUIS, MO. 63166

**WARNING!**


MAINTAIN A SAFE DISTANCE FROM THE TREATMENT AREA. DO NOT ENTER THE TREATMENT AREA UNTIL THE TREATMENT HAS DRIED. WEEDS WILL BE CONTROLLED BY THE TREATMENT. DO NOT APPLY TO CROPS OR SOILS NOT LISTED ON THE LABEL. ALWAYS WEAR PROTECTIVE CLOTHING AND GEAR. ALWAYS USE PROTECTIVE EQUIPMENT. ALWAYS FOLLOW THE DIRECTIONS ON THE LABEL. ALWAYS WEAR PROTECTIVE CLOTHING AND GEAR. ALWAYS USE PROTECTIVE EQUIPMENT. ALWAYS FOLLOW THE DIRECTIONS ON THE LABEL.

NET WT. 50



# Granular **RANDOX T**

NET 50 LBS.



### CALIBRATING GRANULAR EQUIPMENT

Granular equipment should be calibrated at least once a year. To insure accurate and frequent calibration, use a calibrated container. The following instructions apply to all types of granular equipment and tubes may be calibrated for use in this way by using a calibrated measuring bottle or similar measuring device that is graduated in full inches.

Detach the delivery tube from the bottom of the hopper and insert the granular material through this opening while holding the material in a pan. Measure the amount of granular material that falls from the hopper into the pan. Repeat this procedure until the amount of granular material is approximately 100 grams.



Compare the amount of granular material that falls from the hopper into the pan with the amount of granular material that falls from the calibrated measuring device. If the amount of granular material that falls from the hopper is less than the amount of granular material that falls from the calibrated measuring device, the hopper should be calibrated.

### WARNING

Granular material is highly toxic. It should be handled with care. Avoid contact with skin, eyes, nose, mouth, and clothing. If contact occurs, wash thoroughly with water. Do not eat, drink, or smoke while handling this material.

Keep this material away from children and animals. Do not use in areas where children and animals are likely to be present. Do not use in areas where food is prepared or stored.

Do not breathe dust or fumes. Use a respirator if necessary. Do not get material on clothing. If it does, remove clothing and wash thoroughly with water.

Do not dispose of this material in the environment. It is highly toxic to fish and other aquatic life. Do not dispose of this material in the soil.

### WARNING NOTICE

Use this material only in accordance with the directions on the label. Do not use in areas where children and animals are likely to be present. Do not use in areas where food is prepared or stored.

Do not breathe dust or fumes. Use a respirator if necessary. Do not get material on clothing. If it does, remove clothing and wash thoroughly with water.

### WEEDS CONTROLLED

Randex T Granular controls these weeds:

- |                 |        |             |          |         |           |                              |               |                |           |              |                    |              |         |               |                  |                 |           |         |                   |         |           |           |                 |         |
|-----------------|--------|-------------|----------|---------|-----------|------------------------------|---------------|----------------|-----------|--------------|--------------------|--------------|---------|---------------|------------------|-----------------|-----------|---------|-------------------|---------|-----------|-----------|-----------------|---------|
| Annual Ryegrass | Barley | Chenopodium | Cockspur | Cyperus | Crabgrass | Curtain Foxtail (White Star) | Green Foxtail | Yellow Foxtail | Groundnut | Lambquarters | Medicinal Plantain | Mile Mowbray | Mustard | Parrot's Foot | Pigweed (Redtop) | Pigweed (Spiny) | Portulaca | Ragwort | Russell's Thistle | Sandbar | Smartweed | Spiny Top | Stinging Nettle | Thistle |
|-----------------|--------|-------------|----------|---------|-----------|------------------------------|---------------|----------------|-----------|--------------|--------------------|--------------|---------|---------------|------------------|-----------------|-----------|---------|-------------------|---------|-----------|-----------|-----------------|---------|

- |          |           |              |
|----------|-----------|--------------|
| Illinois | Michigan  | Ohio         |
| Indiana  | Minnesota | South Dakota |
| Iowa     | Missouri  | Wisconsin    |
| Kansas   | Nebraska  |              |

### DIRECTIONS FOR USE

For use in corn, soybeans, and sorghum. Apply 1/2 lb. per acre. For use in wheat, apply 1/2 lb. per acre. For use in oats, apply 1/2 lb. per acre. For use in rice, apply 1/2 lb. per acre. For use in sorghum, apply 1/2 lb. per acre.

For best results, apply this material when weeds are young. Do not apply after the weeds have reached the three-leaf stage. Do not apply after the weeds have reached the four-leaf stage.


Do not apply to crops that are sensitive to this material. Do not apply to crops that are in the seedling stage. Do not apply to crops that are in the flowering stage.

Do not apply to crops that are in the seedling stage. Do not apply to crops that are in the flowering stage. Do not apply to crops that are in the seedling stage.

Do not apply to crops that are in the seedling stage. Do not apply to crops that are in the flowering stage. Do not apply to crops that are in the seedling stage.

NET 50 LBS.

1 X0



MINIATURE FACSIMILE  
Smallest Type size on package 12 point.



Granu

RA

SELECTIVE

For pre-emergent  
silage corn, s


READ WARNING LABEL BEFORE

KEEP OUT OF

**WARNING!**  
MAY CAUSE SKIN IRRITATION  
OR EYE INJURY

Do not get on eye or skin  
Avoid breathing dust  
If you take internally  
... ..  
... ..  
... ..

**APPLYING GRANULAR EQUIPMENT**



**WARNING**

**WEEDS CONTROLLED**

Randox Granular controls these weeds:

- Annual Bluegrass
- Barnyardgrass
- Bromegrass (Watergrass)
- Clawweed
- Crab
- Crabgrass
- Crabapple
- Cyprus
- Field Bindweed
- Flourgrass
- Green Foxtail
- Johnsongrass
- Lambquarters
- Lind
- Marestail
- Millett
- Paragrass
- Pigweed (Spiny)
- Ragwort
- Riverbank Swallowtail
- Riverbank Swallowtail
- Riverbank Swallowtail
- Sagebrush
- Sheep Sorrel
- Smartweed
- Soybean
- Spiny Top
- Stinkweed
- Sweet Clover
- Sycamore
- Tanacetum
- Tanacetum
- Timothy
- Trifolium
- Wild Mustard
- Wild Radish
- Wild Yarrow
- Yellow Foxtail
- Zinnia

- Mustard
- Pigweed (Redtop)
- Pigweed (Smooth)
- Purslane
- Ragwort
- Riverbank Swallowtail
- Sheep Sorrel
- Smartweed
- Spiny Top
- Stinkweed
- Sweet Clover
- Tanacetum
- Timothy
- Trifolium
- Wild Mustard
- Wild Radish
- Wild Yarrow
- Yellow Foxtail
- Zinnia

- Illinois
- Indiana
- Michigan
- Minnesota
- Missouri
- Nebraska
- Ohio
- South Dakota
- Wisconsin

**DIRECTIONS FOR USE**

Apply granular material to all open tiller and popcorn

S/1

**WARNING**

This product is a restricted use pesticide. It is to be used only by or under the direct supervision of a certified applicator. For more information on the safe use of this product, consult the label and the pesticide laws of your state.

EYE INJURY

Use only according to label instructions

Do not get in eyes or on skin

Avoid breathing dust

Do not take internally

When the dust from the granules comes in contact

with the eyes, flush with plenty of water

for at least 15 minutes and get medical attention

for skin, wash with soap and water.

Avoid contamination of seed, feed, and foodstuffs

ACTIVE INGREDIENT

INERT INGREDIENTS

100.0  
55.0  
100.0

NET WT

SANTO COMPANY • AGRICULTURAL DIVISION • ST. LOUIS, MISSOURI • U.S.A.

Granular

RANDOX

T

NET 50 LBS.

Groundsel  
Velvetleaf  
Lambquarters

Illinois Michigan Ohio  
Indiana Minnesota South Dakota  
Iowa Missouri Wisconsin  
Kansas Nebraska

DIRECTIONS FOR USE

FIELD CORN, SILAGE CORN, SWEET CORN AND POPCORN

For best results

Apply to soil in rows of corn at planting time. Apply to silage corn at planting time. Apply to sweet corn at planting time. Apply to popcorn at planting time.

WARNING

Do not get in eyes or on skin. Avoid breathing dust. Do not take internally. When the dust from the granules comes in contact with the eyes, flush with plenty of water for at least 15 minutes and get medical attention for skin, wash with soap and water. Avoid contamination of seed, feed, and foodstuffs.

WARNING NOTICE

Do not get in eyes or on skin. Avoid breathing dust. Do not take internally. When the dust from the granules comes in contact with the eyes, flush with plenty of water for at least 15 minutes and get medical attention for skin, wash with soap and water. Avoid contamination of seed, feed, and foodstuffs.