

PM 25

524-124

4/9/69

MONSANTO

131
1/2



BW

AVADEX®

SELECTIVE HERBICIDE

Emulsifiable concentrate for control of wild oats in wheat, barley and field dried peas. Do not use on other crops.

Use only according to label instructions. READ "WARNING NOTICE" BEFORE BUYING OR USING. IF NOTICE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED. Keep out of reach of children.

CAUTION! MAY CAUSE SKIN AND EYE IRRITATION. MAY BE HARMFUL IF SWALLOWED.

Do not get in eyes or on skin. Do not take internally. Do not store near feed and foodstuffs. To avoid contact with skin or eyes, wear rubber gloves and goggles when handling or spraying.

In case of contact, immediately wash skin with soap and plenty of water; for eyes, flush with plenty of water for at least 15 minutes and get medical attention. Remove contaminated clothing. Wash clothing before reuse.

In case of fire, use water spray, foam, dry chemical or CO₂.

ACTIVE INGREDIENT:
S-(2,3,5-Trichloroallyl) diisopropylthiocarbamate


46.3%
53.7%
100.0%

INERT INGREDIENTS:

*Contains 4 lbs. S-(2,3,5-Trichloroallyl) diisopropylthiocarbamate per gallon.

NET CONTENTS: 5 U.S. GALLONS

STORE ABOVE 32° F. TO KEEP FROM FREEZING. Freezing will result in crystals which settle to the bottom of the can. If allowed to freeze, place in a warm room (72° F.) and roll and shake the can frequently for several days to redissove.



Recommended rates should be followed exactly in order to

1. Avoid crop injury

2. Control wild oats

When using Avadex BW, a strip should be left untreated for proof of results. Weed control may be evaluated also by removing a surface inch or two of soil at the time of germination to inspect the number of wild oats that were killed before emergence.

Wild oats are usually killed before emergence, but occasionally, and particularly under dry conditions, plants may reach the 3 or 4 leaf stage before they die.

Under conditions of prolonged high temperature at the time of wild oat germination, or drought in spring, Avadex BW may not provide the usual high standard of wild oat control.

Seeding may be done either before or after spraying and incorporation, depending upon the crop that is to be sown. If seeding is delayed, shallow reworking of the treated area before seeding will not destroy the effects of the chemical.

Do not graze livestock on treated crops.

NOTE: Domestic oats should not be sown where Avadex BW was used the previous year. There is a possibility that Avadex BW may carry over sufficiently to injure this crop.

GENERAL INFORMATION

Keep Avadex BW container closed to prevent spills, evaporation and contamination. Avadex BW is recommended for use in spring wheat, durum wheat, barley and field dried peas only. Other crops should not be treated with Avadex BW because injury may occur.

Spraying equipment must be properly calibrated—application of too much Avadex BW may injure the crop—application of too little Avadex BW may result in poor wild oat control.


Avadex BW should be sprayed on the soil before wild oats germinate. If wild oats have emerged, tillage prior to spraying must be relied upon for their control.

Before applying Avadex BW be sure the soil is in good working condition. This applies particularly to clay soils. Avoid application to a field left in a lumpy condition. Application in a field which is wet, lumpy, rough or ridged will result in reduced wild oat control and increased crop thinning.

When the incorporation of the chemical into the soil must be accomplished as soon as possible on the day of spraying. The incorporation should not be deeper than 2 inches, because deeper incorporation tends to dilute the chemical, decreasing wild oat control and increasing the risk of crop thinning.

When summer fallow or plowed ground is being treated, a heavy trash cover is required. If a heavy trash cover is not available, the soil should be wet or lumpy, (making proper incorporation of the chemical into the soil) a second disking either before or after spraying is recommended. If the soil is loose and lumpy, incorporation with 7 harrows at right angles is adequate.

ACCEPTED FOR REGISTRATION
APR - 9 1969
MONSANTO COMPANY
ST. LOUIS, MISSOURI 63166



MINIATURE FACSIMILE
Smallest Type size on package 12 point.



BW

AVADEX

SELECTIVE HERBICIDE

Emulsifiable concentrate for control of wild oats in wheat, barley and field dried peas. Do not use on other crops.

Use only according to label instructions.
READ "WARNING NOTICE" BEFORE BUYING OR USING.
IF NOTICE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.

Keep out of reach of children.

CAUTION! MAY CAUSE SKIN AND EYE IRRITATION.
MAY BE HARMFUL IF SWALLOWED.

Do not get in eyes or on skin.
Do not take internally.
Do not store near feed and foodstuffs.
To avoid contact with skin or eyes, wear rubber gloves and goggles when handling or spraying.
In case of contact, immediately wash skin with soap and plenty of water; for eyes, flush with plenty of water for at least 15 minutes and get medical attention.
Remove contaminated clothing. Wash clothing before reuse.
In case of fire, use water spray, foam, dry chemical or CO₂.

ACTIVE INGREDIENT:
S-(2,3,3-Trichloroallyl) disopropylthiocarbamate 46.3%

INERT INGREDIENTS: 53.7%
100.0%

*Contains 4 lbs. S-(2,3,3-Trichloroallyl) disopropylthiocarbamate per gallon.
NET CONTENTS: 5 U.S. GALLONS
STORE ABOVE 32° F. TO KEEP FROM FREEZING. Freezing will result in crystals which settle to the bottom of the can. If allowed to freeze, place in a warm room (72° F.) and roll and shake the can frequently for several days to redissolve.



SPECIFIC CROP USE

Before using, be sure to read and understand the "Caution," "General Information" and "How to Mix and Apply" sections of this label.

Avadex BW is not recommended for use in Arizona.

BARLEY

Avadex BW is recommended after seeding as follows:

- After Seeding Method**
1. Seed to a depth of 2 to 3 inches.
 2. Apply 1½ quarts per acre BEFORE crop emergence, and before kernel sprouts exceed ¼ inch in length. (See "How to Mix and Apply" section.)
 3. Incorporate with 2 harrowings at right angles as soon as possible on the day of spraying. This method of incorporation is not effective if compact soil prevents penetration of harrow teeth, or if trash accumulates in the harrow sections.

Before Seeding Method—While application after seeding barley is recommended, Avadex BW can also be used before seeding. Heavy stubble should be thoroughly tilled either before or after application as follows:

1. Apply 1½ quarts Avadex BW per acre. (See "How to Mix and Apply" section.)
2. Incorporate with a disk-type implement or cultivator to a depth of 2 inches or less, followed by a harrow as soon as possible on the day of spraying. Two harrowings at right angles after spraying is adequate incorporation if the soil is loose and free of trash.
3. Seed barley BELOW the chemical layer in the soil to a depth of 3 inches, any time from immediately after, up to 3 weeks after spraying.

NOTE—Crop thinning has been known to occur. This is usually avoided by applications AFTER seeding and minimized in applications before seeding when the seed is placed ½ to 1 inch below the Avadex BW treated soil layer. Thinning is usually more than offset by tillering and increased yields.

SPRING AND DURUM WHEAT

Properly incorporated, 1 quart of Avadex BW per acre should provide reasonable wild oat control.

AVADEX BW IS RECOMMENDED AFTER SEEDING ONLY AS FOLLOWS:

1. On summerfallow or plowed ground, seed to a depth of 2 to 3 inches. If drilling leaves ridges in loose soil, the field should be harrowed prior to spraying.

SPECIFIC CROP USE (continued)

2. Spray Avadex BW at 1 quart per acre BEFORE crop emergence and before kernel sprouts exceed ¼ inch in length. (See "How to Mix and Apply" section.)

3. Incorporate with 2 harrowings at right angles as soon as possible on the day of spraying. Two harrowings is adequate incorporation if the soil is loose and free of trash.

NOTE—Occasionally some crop thinning may occur, particularly when there is abundant rainfall after spraying and before crop emergence. Thinning is usually avoided when the seed is placed ½ to 1 inch below the treated Avadex BW soil layer. Thinning is usually more than offset by tillering and increased yields.

FIELD DRIED PEAS

Avadex BW is recommended either before or after seeding.

Before Seeding Method

1. Apply 1½ quarts Avadex BW per acre (See "How to Mix and Apply" section.)
2. Incorporate with a disk-type implement or cultivator to a depth of 2 inches or less, followed by a harrow as soon as possible on the day of spraying. Two harrowings at right angles after spraying is adequate incorporation if the soil is loose and free of trash.
3. Seed peas any time from immediately after, up to 3 weeks after spraying.

NOTE—Leaf crinkling and delayed maturity of dry peas may occur, particularly on clay points in the Northwest, but they are usually more than offset by a reduction of wild oats.

After Seeding Method

1. On summerfallow or plowed ground, seed to a depth of 2 to 3 inches.
2. Spray Avadex BW at 1½ quarts per acre soon after seeding. (See "How to Mix and Apply" section.)
3. Incorporate as for barley "AFTER seeding"

895 15-000.11 (USDA Reg. No. 524-124)

WARNING NOTICE

Monsanto Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. This product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as for example incompatibility with other products, the manner of its use or application, or the presence of other products or materials in or on the soil or crop. Monsanto makes no other express or implied warranty of fitness or merchantability. The exclusive remedy of the user or buyer, and the limit of the liability of Monsanto Company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this notice which may not be varied by any verbal or written agreement.

HOW TO MIX AND APPLY AVADEX BW

Dilute Avadex BW with 5 or more gallons of water per acre for an over-all treatment. Higher volumes approaching 10 gallons per acre will usually result in more uniform spraying and less evaporation, and slightly better wild oat control.

Spraying during winds above 10 miles per hour will result in a loss of Avadex BW below the amount required for good wild oat control.

For best mixing of Avadex BW with water, proceed as follows:

1. Fill the tank approximately three-quarters full with water.

HOW TO MIX AND APPLY AVADEX BW (continued)

2. Carefully add the proper amount of Avadex BW while mixing.
3. Fill the remaining portion of the tank with water.
4. Mix thoroughly again if the solution has been left in the tank for any length of time.
5. The sprayer by-pass valve should be in operation before and during application.

Before spraying, check all hose connections to see that they are properly secured. Adjust the height of the spray boom in the field to ensure an even distribution of the chemical on the soil. Avoid overlapping of the spray. Pressure should not be above 30 p.s.i. and just high enough to provide a uniform spray pattern. Boomless sprayers should not be used because the distribution of Avadex BW will be too uneven.

A field marker or chain should be attached to both ends of the sprayer boom to serve as a marker, thus avoiding overlapping or untreated strips. Overlapping will double the recommended rate and crop injury may result.

Promptly change clothing which becomes wet by accidental spillage or drift from spray. Wash clothing before re-use.

For methods of incorporation, see "Specific Crop Use" section.

