

# METHYL PARATHION 80

Stabilized

FOR USE

## ACTIVE INGREDIENTS:

O,O-dimethyl O-p-allyloxyphenyl phosphorothioate ..... 80.0%

Xylene ..... 12.5%

INERT INGREDIENTS: ..... 7.5%

Keep out of reach of children.

## DANGER!

POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING.

RAPIDLY ABSORBED THROUGH SKIN.

REPEATED EXPOSURE MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams, ponds and public water. Do not contaminate water by cleaning of equipment or discharge of wastes.

NOTE: DO NOT HEAT THIS MATERIAL ABOVE 131°F (55°C). STORE AT TEMPERATURES ABOVE 65°F (18.3°C) TO AVOID CRYSTALLIZATION.

## DO NOT GET ON SKIN, IN EYES, ON CLOTHING:

Wear clean heavy natural rubber gloves, goggles, and clean waterproof or freshly laundered protective clothing (coveralls, rubber boots, cap, etc.). Destroy and replace gloves frequently.

Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact immediately remove contaminated clothing and wash skin thoroughly with soap and water; for eyes flush with water for 15 minutes.

## DO NOT BREATHE VAPOR OR DUST:

Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for METHYL PARATHION protection.

IN CASE OF SPILL OR LEAK, keep all unprotected persons away. Cover with absorbent such as soda ash, lime, clay or saw dust. Sweep up and bury. Wash area thoroughly with strong lye solution.

EMPTY DRUMS should be decontaminated by washing with hot, 5% caustic soda solution. Perforate to prevent use other than for this insecticide. Bury or discard in a safe place away from water supplies.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. In case of fire, use water spray, foam, dry chemical or CO<sub>2</sub>.

## POISON

FIRST AID: ATROPINE IS AN ANTIDOTE.

CONSULT PHYSICIAN FOR EMERGENCY

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED METHYL PARATHION POISON

If symptoms or signs of poisoning include vision, abdominal cramps, and tightness in the chest, do not wait for a doctor but give two atropine tablets (each 1/100 grain or 0.65 milligrams) at once to children under five years of age.)

Move patient immediately from the area where methyl parathion is present.

Remove contaminated clothing and wash thoroughly clean with plenty of soap and water to remove traces of methyl parathion. If swallowed, vomit by giving salty or soapy warm water. Victim lie down and keep quiet.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

MONSANTO COMPANY, AGRICULTURAL DIVISION, ST. LOUIS, MO.

**ACCEPTED**

**DEC 14 1972**

**UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON PREVENTION  
EP ORDER NO. 524-136**

**FOR USE BY MANUFACTURERS IN PREPARATION OF PESTICIDES**

**POISON** 

**ATROPINE IS AN ANTIDOTE—**

**PHYSICIAN FOR EMERGENCY SUPPLY.**

**PHYSICIAN AT ONCE IN ALL CASES OF  
METHYL PARATHION POISONING.**

**is or signs of poisoning include blurred  
ominal cramps, and tightness in the chest,  
for a doctor but give two atropine tablets  
0 grain or 0.65 milligrams) at once. (One  
children under five years of age.)**

**ent immediately from the area where  
athion is present.**

**contaminated clothing and wash the skin  
plenty of soap and water to remove all  
methyl parathion. If swallowed, induce  
y giving salty or soapy warm water. Have  
down and keep quiet.**

**IVE ANYTHING BY MOUTH TO AN  
IOUS PERSON.**

**PHYSICIANS NOTE:** This product is a cholinesterase inhibitor. Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

**TREATMENT:** Large doses of atropine are required. For adults, give 2.0 mg. to 4.0 mg. of atropine, preferably by intravenous injection, at once and repeat every 10 to 15 minutes until pupils dilate. In addition to atropine, if 2-PAM Chloride (pralidoximechloride) is available, administer according to manufacturer's directions. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

LOT NO. \_\_\_\_\_

PACKER \_\_\_\_\_

**NET LEGAL**

**LBS.**

**CULTURAL DIVISION, ST. LOUIS, MO., U.S.A. 896.07-000.08/53 (EPA Reg. No. 524-136-A)**



**W A R N I N G**  
**KEEP AWAY from FEED**  
**P O I S O N**  
**CAUTION—DO NOT**  
**IF LEAKING**  
**DO NOT**

This is to certify that the  
are properly described by  
and marked and are in  
transportation according  
prescribed by the  
Transportation

**MONSANTO**

Shipper's name  
hereon for  
ments  
EXPR

**METHYL PARATHION**

In case of any evidence of damage  
telephone collect, (205) 236-6388