




PRECAUTIONARY STATEMENT
Hazard to Humans and Domestic Animals

DANGER—POISON—PRECAUTIONS



CAN KILL YOU IF SWALLOWED

This product is highly toxic. Small amounts can be fatal. Do not swallow.



CAN KILL YOU BY SKIN CONTACT

This product is highly toxic. Do not allow it to come in contact with your skin.



CAN KILL YOU IF BREATHED

This product is highly toxic. Do not breathe the vapors.

MINIATURE FACSIMILE
Smallest Type Size on Package is Legal

POISON
See side panels for antidote and precautions

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

STOP! READ THE LABEL



CAN KILL YOU

DANGER
Keep out of reach of children

POISON SIGNS (Symptoms)

Parathion is a very dangerous substance which enters the body on contact with skin, clothes, eyes and eyes. Clothing wet with the material must be removed immediately. Exposed persons must receive prompt medical treatment if they have it. Some of the signs and symptoms of poisoning are: headache, nausea, vomiting, diarrhea, weakness, blurred vision, pinpoint pupils, tightness in the chest, labored breathing, muscular twitching, watering of eyes, drooping eyelids, difficulty in breathing, muscle spasms and coma.

If inhaled and victim is awake, conscious, make him vomit quickly. Have victim wear goggles, gloves, and induce vomiting by forcing back of throat with finger. Do not induce vomiting by pouring anything into mouth to an unconscious person. Have a doctor called and keep quiet. Get medical attention. In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Seek medical attention.

NOTE TO PHYSICIAN

Parathion is a very dangerous substance which enters the body on contact with skin, clothes, eyes and eyes. Clothing wet with the material must be removed immediately. Exposed persons must receive prompt medical treatment if they have it. Some of the signs and symptoms of poisoning are: headache, nausea, vomiting, diarrhea, weakness, blurred vision, pinpoint pupils, tightness in the chest, labored breathing, muscular twitching, watering of eyes, drooping eyelids, difficulty in breathing, muscle spasms and coma.

POISON
ATROPINE IS AN ANTIDOTE
CONSULT PHYSICIAN FOR
EMERGENCY SUPPLY

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED PARATHION POISONING. Symptoms of signs of poisoning include: headache, nausea, vomiting, diarrhea, weakness, blurred vision, pinpoint pupils, tightness in the chest, labored breathing, muscular twitching, watering of eyes, drooping eyelids, difficulty in breathing, muscle spasms and coma.


FIRST AID (PRACTICAL TREATMENT)

Call a doctor, physician or hospital. Follow their advice. Explain that the victim has been exposed to parathion and describe this label. If breathing has stopped, start artificial respiration immediately and transfer to hospital as quickly as possible.

PARATHION INSECTICIDE
Niran 6-3

Monsanto

NOT FOR HOME USE



NET U.S. GAL.

BEST DOCUMENT AVAILABLE

DEC 21 1982

Ms. Glynadee Edwards
Monsanto Company
1101 17th Street, NW
Suite 604
Washington, D.C. 20036

Dear Ms. Edwards:

Subject: Miran S-3
Revised Ingredient Statement
EPA Registration No. 524-144
Your Application Dated November 30, 1982

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

TS-767:Ellenberger:DCR-28989:WANG-1657C:hk:Raven:479-2013:12/16/82

