

READ THE ENTIRE LABEL FOR FAR-GO BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the Label Booklet for Far-Go and this supplement.

Far-Go®

HERBICIDE BY **Monsanto**

(EPA Reg. No. 524-145)

FOR DISTRIBUTION AND USE ONLY WITHIN KANSAS,
OKLAHOMA, AND TEXAS.

For FALL treatment to control **BROMUS SPECIES (B. TECTORUM, B. JAPONICUS, B. SECALINUS)** in WINTER wheat.

Keep out of reach of children.

WARNING!

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000

Far-Go™ is a registered trademark of Monsanto Company



MONSANTO COMPANY • AGRICULTURAL PRODUCTS • ST. LOUIS, MISSOURI, 63167 U.S.A.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with the labeling.

This label must be in the possession of the user at the time of herbicide application.

WINTER WHEAT—PLANTED WITH DOUBLE DISK PRESS DRILL

GENERAL INFORMATION

WINTER WHEAT THINNING AND STUNTING MAY BE OBSERVED ESPECIALLY IF WINTER wheat is under stress from soil crusting, excessive rainfall or irrigation, snow cover, cool-wet conditions, drought, soil compaction, deep seeding, diseases, insects, or any factor which inhibits emergence or development. THINNING IS USUALLY MORE THAN OFFSET BY TILLERING.

Under certain conditions, Far-Go may cause unacceptable injury to certain wheat varieties. The grower should consult the state university extension specialist, seed producer or supplier, or the local Monsanto representative for those wheat varieties which can be used safely with Far-Go.

DO NOT GRAZE LIVESTOCK ON TREATED CROP.

FIELD PREPARATION

Before applying this herbicide, all deep tillage must be completed and the soil should be in good working condition. A field with heavy trash cover should be worked to remove as much surface trash as possible, prior to application. Do not apply if soil is wet or lumpy, making proper incorporation difficult.

LUMPY SOIL ALSO MAY RESULT IN BROMUS GERMINATION AND EMERGENCE ABOVE THE TREATED SOIL LAYER AS LUMPS ARE MELTED DOWN BY SUBSEQUENT RAINFALL. BROMUS GERMINATED AT TIME OF HERBICIDE APPLICATION MUST BE DESTROYED BY TILLAGE OR OTHER METHODS.

RATE OF APPLICATION

Apply this product at an overall rate of 1.5 quarts per acre. Do not use other soil-applied herbicides in conjunction with this herbicide treatment.

INCORPORATION PRIOR TO SEEDING

Apply this product in the FALL before seeding and shallowly incorporate this herbicide immediately to a depth of no more than 2 inches, using a field cultivator with harrow attachment. This is accomplished by setting the incorporation implement to work the soil no deeper than 3 to 4 inches. Tillage implements usually incorporate to one-half of the tillage depth. A second incorporation (similar to the first incorporation) at an angle should provide best results. To avoid deep incorporation, do not use disk implements.

SEEDING INFORMATION


WINTER WHEAT SHOULD BE PLANTED $\frac{3}{4}$ TO $1\frac{1}{4}$ INCHES DEEP TO FACILITATE RAPID EMERGENCE. EARLY SEEDING WILL MINIMIZE THE PROBABILITY OF WHEAT INJURY DUE TO SLOW EMERGENCE AND GROWTH DUE TO COLD LATE FALL WEATHER CONDITIONS.

Read "LIMIT OF WARRANTY AND LIABILITY" in the Far-Go pamphlet before buying or using this product. If the terms are not acceptable, return the product unopened at once.

A-MAP-2707-2-12-85

©MONSANTO COMPANY 1985

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167



Emulsifiable herbicide for control of wild oats in winter wheat, spring
and durum wheat, barley, green peas, field dried peas and lentils.

Carefully follow detailed instructions in attached booklet

MINIATURE FACSIMILE:
Type Size Meets Requirements

PRINT: PMS 347 Green

LIMIT OF WARRANTY AND LIABILITY

This company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this company, including but not limited to incompatibility with products other than those set forth in the Directions, unusual weather or weather conditions which are

outside the range considered normal at the application site and for the time period when the product is applied with the normal range being determined on the basis of the average range for the prior 40 years computed from the best available information, and all weather perils including but not limited to hurricanes, tornadoes and floods, as well as weather considerations set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil or crop.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY

THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES.

The buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

© MONSANTO COMPANY 1984

*Trademark Monsanto Company

EPA Est. 17507 MT 1

815 15 000 467

Read the
Use only
Read the
buying
once un
PP
Hazar
Keep o
WA
CAUSES
HARMF
Do not
Avoid br
Wash the
To avoid
and go

FIRST AID IN
with soap
plenty of
medical at
which beco
from spray

In case
Call Collec

Far-Go

HERBICIDE BY

Monsanto

Emulsifiable herbicide for control of wild oats in winter wheat, spring and durum wheat, barley, green peas, field dried peas and lentils.

Complete Directions for Use

Far-Go® is a registered trademark of Monsanto Company

MONSANTO COMPANY 1984

EPA Reg. No. 524-145-AA 815.15.000.47/53

Read the entire label before using this product.

Use only according to label instructions.

NEVER REFORMULATION OR REPACKAGING

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

LIMIT OF WARRANTY AND LIABILITY

This company warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this company, including but not limited to: incompatibility with products other than those set forth in the Directions; unusual weather or weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied with the normal range being determined on the basis of the average range for the prior 40 years computed from the best available information; and weather perils, including but not limited to hurricanes, tornadoes and floods, as

well as weather considerations set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil or crop.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED. OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES.

The buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

PRECALCULATORY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

WARNING!

CAUSES SKIN AND EYE IRRITATION.

HARMFUL IF SWALLOWED.

Do not get in eyes, on skin or on clothing.

Avoid breathing vapors or spray mist.

Wash thoroughly after handling.

To avoid contact with skin or eyes, wear rubber gloves and goggles when handling or spraying.

FIRST AID: IN CASE OF CONTACT, immediately wash skin with soap and plenty of water. For eyes, flush with plenty of water for least 15 minutes and get medical attention. Remove contaminated clothing which becomes wet by accidental spillage or drift from spray. Wash clothing before reuse.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Avoid direct application to any body of water. Do not contaminate water by disposal of waste or cleaning of equipment.

Physical or Chemical Hazards

COMBUSTIBLE.

Do not use, pour, spill or store near heat or flame.

Use only with adequate ventilation.

In case of:

FIRE, use water spray, foam, dry chemical or CO₂.
SPILL or LEAK, flush area with water spray.

Storage and Disposal

Do not contaminate water, foodstuffs, seed or feed by storage and disposal.

STORAGE:

STORE ABOVE 32°F TO KEEP FROM FREEZING.

Freezing will result in crystals which settle to the bottom of the can. If allowed to freeze, place in a warm room (72°F.) and roll and shake the container frequently for several days to redissolve before using.

Keep this container closed to prevent spills, evaporation and contamination.

DISPOSAL:

Wastes of this pesticide may cause skin and eye irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

(See individual container label for disposal information.)

ACTIVE INGREDIENT

S-(2,3,3-Trichloroallyl)-disopropylthiocarbamate	46.3%
INERT INGREDIENTS	53.7%
	100.0%

Contains 4 pounds S-(2,3,3-Trichloroallyl)-disopropylthiocarbamate per gallon.

GENERAL INFORMATION

This herbicide or any labeled tank mixture of this herbicide should be sprayed on the soil and incorporated before weeds germinate. If weeds have emerged prior to application, they must be controlled by tillage.

Before applying this herbicide or any labeled tank mixture of this herbicide, the soil should be in good working condition. This applies particularly to clay soils. Avoid application to a field left in a ridged condition. Application to a field which is wet, lumpy, rough or ridged will result in reduced wild oat control and increased crop thinning.

This product or any labeled tank mixture of this herbicide must be incorporated into the soil immediately after spraying. Incorporation should not be deeper than 2 inches and should be done at 6 to 7 miles per hour to ensure good mixing into the soil.

Seeding may be done either before or after spraying, and incorporation depending upon the crop that is to be sown. See "Directions for Use" section of this label. If seeding is delayed, shallow reworking of the treated area before seeding will not destroy the effects of the chemical.

Weed control may be evaluated by removing a surface inch of soil at the time of germination to inspect the number of wild oats that were killed before emergence.

This herbicide is primarily absorbed by wild oat shoots from the treated layer of the soil. Wild oats are usually controlled before they emerge through the soil. Occasionally and particularly under dry conditions and cold soil temperatures, some wild oats may emerge and reach the 3 to 4 leaf stage before dying. Under conditions of prolonged high temperature at the time of wild

30F6

out germination or drought in the spring, this product may not provide the usual high standard of wild oat control.

ATTENTION

Occasionally some crop thinning may occur in wheat and barley from fall or spring treatments, particularly where water accumulates in low places after spraying and before crop emergence. Crop thinning is minimized in applications before seeding when the seed is placed $\frac{1}{2}$ to 1 inch below the herbicide-treated soil layer. Thinning is usually more than offset by increased tillering.

Do not graze livestock on treated crops.

Keep this container closed to prevent spills, evaporation and contamination.

Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST

Herbicides may not always mix evenly throughout a sprayable fluid fertilizer or the compounds may separate too quickly to make their combined use of practical value. This may be due to certain characteristics of the different fluid fertilizers. A simple test using small quantities of the components is suggested to provide compatibility potential.

The test follows:

A. Materials Required For A Compatibility Test

1. Two one-quart jars with lid or stopper (marked "with" and "without").
2. Teaspoons (for a more exacting test, a five to ten milliliter (ml.) pipette or graduated cylinder is desirable).
3. A minimum of one quart of the sprayable fluid fertilizer to be tested.
4. The herbicide to be mixed.
5. A compatibility agent (the purpose of the adjuvant is to help keep the fertilizer and crop protection chemical in suspension if this assistance is needed).

B. Procedure

1. Add one pint of the sprayable fluid fertilizer that will be used or other herbicide carrier to each jar marked "with" and "without".



Add One Pint Liquid Fertilizer To Two Quart Jars



2. To the jar marked "with" add $\frac{1}{4}$ TEAspoon or 1.2 milliliters of a suitable compatibility agent. Shake gently for five to ten seconds to mix. ($\frac{1}{4}$ TEAspoon is the equivalent of two pints per 100 gallons of liquid fertilizer.)



To Jar Marked "With" Add Compatibility Agent And Shake To Mix



3. To each jar add the appropriate amount of herbicide(s). Shake gently for five to ten seconds after each addition.



Add Herbicide(s) To Both Jars And Shake To Mix



HERBICIDE	RATE	Amount To Be Added Per Pint Of Sprayable Fluid Fertilizer (Assuming Volume is 25 gal. A)	
		Level	TEAspoons, Milliliters
Far-Go*	1 qt.	=	0.9 or 4.7
	1.25 qt.	=	1.2 or 5.9

This compatibility test is designed for 25 gallons of spray per acre with the maximum labeled rate of herbicide. For changes in spray volume, make appropriate changes in the ingredients of the test. Regardless of spray volume, the amount of compatibility agent should be equal to two or three pints (two pints = $\frac{1}{4}$ TEAspoon or 1.2 milliliters, three pints = $\frac{3}{4}$ TEAspoon or 1.8 milliliters per pint of sprayable fluid fertilizer) per 100 gallons of liquid fertilizer.

C. Observations And Decisions

1. If the herbicide(s) and the sprayable fluid fertilizer are compatible.

compatibility agent is necessary.

Five minutes after the final addition and mixing, observe both jars for the formation of large flakes, sludge, gels or other precipitates. Observe if the herbicide(s) cannot be physically mixed with the liquid fertilizer but remains as small oily particles in the solution.

If compatibility in any form described above occurs in the jar "with" the compatibility agent added, the liquid fertilizer and the herbicide(s) should not be used together in the same spray tank.

If incompatibility as described above occurs in the jar "without" the adjuvant but not in the jar "with" adjuvant, the use of a compatibility adjuvant is recommended.

Both jars should be allowed to stand and to be observed periodically for one-half hour. If the separate layers of liquid fertilizer and additives can be resuspended by shaking, commercial application is possible. This product normally will go to the top after standing.

If the herbicide(s) is compatible with fluid fertilizer in the foregoing test without having to use a compatibility agent, fluid fertilizer may be used for the premixing. If it is not compatible, the herbicide(s) should be premixed with water before adding to the spray tank.

MIXING INSTRUCTIONS

Always predetermine the compatibility of this herbicide or labeled tank mix of this herbicide with water carrier or sprayable fluid fertilizer carrier by mixing small proportional quantities in advance. See the compatibility test section of the label for instructions as to how to determine the compatibility of this herbicide with sprayable fluid fertilizer.

NOTE: This herbicide can be applied in sprayable fluid fertilizer in the spring to barley, spring and durum wheat only.

Mix Far-Go or labeled tank mix of Far-Go with the appropriate carrier as follows:

1. Fill the tank approximately three-quarters full with water. Check for adequate agitation.
2. Carefully add the proper amount of herbicide(s) into the tank while mixing.
3. Fill the remaining portion of the tank with water.
4. Maintain good agitation at all times until the contents of the tank are sprayed.
5. Mix thoroughly again if the solution has been left in the tank for any length of time.
6. The sprayer by-pass valve should be in operation before and during application.

APPLICATION INFORMATION

Before spraying, check all hose connections to see that they are properly secured. Spraying equipment must be properly calibrated. Adjust the height of the spray boom in the field to ensure an even distribution of the chemical on the soil. Pressure should not be above 30 p.s.i. and just high enough to provide a uniform spray pattern. Boomless sprayers should not be used because the distribution of this product will be too uneven.

A field marker or chain should be attached to both ends of the sprayer boom to serve as a mark, avoiding overlapping or untreated strips. Overlapping will double the recommended rate and crop injury may result.

Apply in 10 or more gallons of water or sprayable fluid fertilizer per acre. Higher volumes will usually result in more uniform spraying and less evaporation.

Spraying during winds above 10 miles per hour will result in a loss of herbicide below the amount recommended for good wild oat control.

WEEDS CONTROLLED WITH FAR-GO*

When applied as directed under conditions described on this label, Far-Go CONTROLS the following weeds:

Oats (wild)

Avena fatua

Ryegrass (annual)

Lolium multiflorum
(Oregon only)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in any manner inconsistent with its labeling.

FIELD PREPARATION

Before applying this herbicide or any labeled tank mixture, the soil should be in good working condition. All deep tillage must be completed prior to application of this herbicide. If plowed ground is being treated, a single cultivation is recommended prior to spraying. If stubble ground with a heavy trash cover is being treated, or if the soil is wet or lumpy (making proper mixing difficult), two cultivations before spraying are recommended.

FERTILIZER IMPREGNATION WITH FAR-GO®

FOR USE IN ALL STATES EXCEPT CALIFORNIA

The application of Far-Go impregnated on fertilizer may be made as follows

Crops: Winter Wheat, Spring Wheat and Barley

Time of Application, Rate and Fertilizer Carrier

• Fall Applied and Fall Incorporated

Spring wheat and barley
1.25 lbs. ai. A of Far-Go (the 1.5 lb. rate may be used only on fine textured soil)

Urea or dry blended fertilizer carrier

Winter wheat

1.25 lbs. ai. A of Far-Go

Urea fertilizer carrier

Preplant incorporated

• Spring Applied and Preplant Incorporated

Spring barley only

1.25 lbs. ai. A of Far-Go

Dry blended fertilizer carrier

APPLICATION WITH DRY BULK FERTILIZER

FOR USE IN ALL STATES EXCEPT CALIFORNIA

The herbicide-fertilizer impregnation process must be done at commercial fertilizer or chemical dealerships that are properly equipped for this procedure.

Dry bulk fertilizer may be impregnated with Far-Go herbicide. Far-Go must be impregnated on a minimum of 150 pounds per acre of dry fertilizer. Applications of dry bulk fertilizer impregnated with Far-Go herbicide may be made in the fall or spring, but only preplant incorporated treatments to barley may be made in the spring. Spring planted wheat or barley may be seeded into fall Far-Go herbicide applications in accordance with label directions. Applications should be made as soon as possible after blending.

The following table provides a reference to determine the amount of liquid herbicide to be mixed per ton of dry bulk fertilizer.

DRY FERTILIZER (Lbs. Acre)	ACRES COVERED (Per Ton)	FAR-GO® HERBICIDE RATE (Quarts Per Acre)	
		1.25	1.50
150	13.3	16.5	20.0
160	12.5	15.8	18.8
170	11.7	14.8	17.5
180	11.1	14.0	16.8
190	10.5	13.0	15.8
200	10.0	12.5	15.0
210	9.5	12.0	14.3
220	9.1	11.5	13.5
230	8.7	11.0	13.0
240	8.3	10.5	12.5
250	8.0	10.0	12.0
260	7.7	9.6	11.6
270	7.4	9.3	11.1

Mix and blend the dry fertilizer and herbicide in a closed rotary drum-type mixer allowing sufficient time to ensure uniform coverage. The nozzle used to spray the herbicide treatment must be placed inside the mixer to provide uniform spray coverage of the tumbling fertilizer.

Spread the fertilizer/herbicide mixture with a properly calibrated applicator. Be certain the material is applied uniformly to the soil surface. When using fan spreaders, a 100% overlap is recommended. Fan spreaders should be calibrated to put out one-half the desired rate per acre and application overlapped by doubling back to cover one-half of the previous swath. Dribble spreaders must be calibrated with the fertilizer/herbicide mixture since the flow pattern will generally be slower than straight dry fertilizer. Non-uniform spreading of the fertilizer/herbicide mixture may result in unsatisfactory weed control or crop injury.

Follow normal Far-Go herbicide incorporation procedures to incorporate the fertilizer/herbicide mixture.


A partial list of fertilizers which may be impregnated with Far-Go are:

Diammonium phosphate	18.46.0
Treble super phosphate	0.46.0
Urea	45.0.0
Ammonium sulfate	21.0.0
Potassium chloride	0.0.60
Potassium sulfate	0.0.52

NOTE: Far-Go can be impregnated on any commonly used dry fertilizer except ammonium nitrate and those containing ammonium nitrate, potassium nitrate or sodium nitrate. All individual state regulations relating to dry bulk fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the fertilizer/herbicide mixture.

EPA Reg. No. 524-145-AA

815.15.000.47/53



MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

HOW TO INCORPORATE

DO NOT INCORPORATE FAR-GO MORE THAN 2 INCHES. This is accomplished by setting the incorporation implement to work the soil no deeper than 3 to 4 inches. Tillage implements usually incorporate to one-half of the tillage depth. Shallow incorporation is necessary to prevent diluting the herbicide, thus decreasing wild oat control and increasing the risk of crop injury. Incorporation should be performed immediately after herbicide application. Do not use disc implements for incorporation.

CROPPING SYSTEMS

FAR-GO®

FALL TREATMENT: Winter Wheat

CROP	FAR-GO® RATE (Qts / A)	SEEDING DEPTH (Inches)
Winter Wheat— Apply just before seeding to soon after seeding before wild oats germinate	1.25	2.5 to 3

*DO NOT use on clay knobs. DO NOT use on fields where seeding will be shallow (where seed must be "dusted in" or "floated on"). Winter wheat may be drilled immediately or up to 3 weeks after spraying and incorporating.

For applications to Winter Wheat either before or after seeding, apply and shallowly incorporate with a spike tooth or spring tooth harrow followed with a second incorporation at right angles. When incorporating after seeding Winter Wheat, adjust cultivation depth so as not to disturb wheat seed.

FALL TREATMENT: Barley, Durum Wheat, and Spring Wheat

CROP	FAR-GO® RATE (Qts / A)	SEEDING DEPTH (Inches)
Spring or Durum Wheat	1	2 to 3
Barley — Apply within 3 weeks of soil freezeup	1.25	2 to 3

Breakdown of this product is minimal at 40°F and below, therefore very limited herbicide will be lost after soil temperatures drop below 40°F.

For applications to provide wild oat control in spring planted Spring Wheat, Durum Wheat and Barley, apply on summer fallow ground which is free of lumps and trash, and single pass incorporate with equipment similar to a culti-harrow or a duckfoot cultivator, multiweeder or spring tooth harrow.

For application to stubble ground with heavy trash cover, fields should be worked twice with a field cultivator or chisel plow to remove as much surface trash as possible. When a chisel plow is used it should be followed by a field cultivator before application. Apply this herbicide and single pass incorporate with a field cultivator or culti-harrow.

If soil must be ridged after incorporating to prevent soil erosion by high winds, the depth of ridging should be kept to a minimum. Deep ridging may decrease the effectiveness of the herbicide or result in crop injury in the spring.

In preparing the herbicide-treated area for seeding in the spring, care must be taken to avoid working the soil any deeper than 3 to 4 inches. Deep tillage, greater than 4 inches in depth, tends to reduce the wild oat control and increases the risk of crop injury.

SPRING TREATMENT: Spring Wheat (Montana and North Dakota only), Barley, Green Peas, Field Dried Peas, and Lentils

BEFORE SEEDING APPLICATION

CROP	FAR-GO® RATE (Qts / A)	SEEDING DEPTH (Inches)
Spring Wheat* — Apply just before seeding with HOEDRILLS	1.0	½ inch Below The Incorporated Herbicide
Barley — Apply just before seeding	1.25	2 to 3
Green Peas**	1.25	2 to 3
Field Dried Peas**	1.25	2 to 3
Lentils — Apply up to 3 weeks before seeding	1.25	2 to 3

*Montana and North Dakota only

Leaf crinkling and delayed maturity of peas may occur, particularly on clay points in the Northwest, but this is usually more than offset by a reduction of wild oat competition and increased yields.

Before Seeding Incorporation: For application to stubble ground, work fields twice with a disc, field cultivator or chisel plow to provide soil in a good working condition and to remove surface trash. Apply and incorporate with equipment such as a culti-harrow, springtooth cultivator or duckfoot cultivator followed with a second incorporation at right angles.

FOR SPRING WHEAT IN MONTANA AND NORTH DAKOTA ONLY: Shallowly incorporate immediately using a field cultivator with rod weeder or harrow attachment to a depth of no more than 2 inches. PLACE SEED ½ INCH BELOW THE INCORPORATED HERBICIDE BY USING ONLY HOEDRILLS.

SPRING TREATMENT: Barley, Green Peas, Field Dried Peas, Lentils, Durum Wheat and Spring Wheat

AFTER SEEDING INCORPORATION

CROP	FAR-GO® RATE (Qts / A)	SEEDING DEPTH (Inches)
Spring and Durum Wheat	1	2 to 3
Barley	1.25	2 to 3
Green Peas	1.25	2 to 3
Field Dried Peas	1.25	2 to 3
Lentils	1.25	2 to 3

Leaf crinkling and delayed maturity of peas may occur, particularly on clay points in the Northwest, but this is usually more than offset by a reduction of wild oat competition and increased yields.

Apply after seeding before crop emergence and before wheat and barley kernel sprouts exceed ½ inch in length and before green peas, field dried peas and lentils shoot sprouts exceed ½ inch in length.

After Seeding Incorporation: Apply immediately after seeding and shallowly incorporate with equipment such as a flex multiweeder or harrow, and follow with a

second incorporation at a right angle. Adjust incorporation equipment to a depth so as not to disturb the seed.

FAR-GO® PLUS TREFLAN™ (TANK MIXTURE)

Barley and Wheat

When applied as directed under conditions described on this label, this tank mixture **CONTROLS** these additional weeds:

Foxtail (green) Setaria viridis	Foxtail (yellow) Setaria lutescens
------------------------------------	---------------------------------------

Green and yellow foxtail are commonly referred to as pigeon grass.

*Treflan is a trademark of E.I. du Pont de Nemours & Co.

SPRING TREATMENT: Barley, Spring Wheat, and Durum Wheat

AFTER SEEDING INCORPORATION

CROP	FAR-GO® RATE (Qts / A)	TREFLAN Qts / A
Spring Wheat, Durum Wheat and Barley	1.0	0.5

Seed to a depth of 2 to 3 inches.

Apply immediately after seeding.

After Seeding Incorporation: Apply immediately after seeding and shallowly incorporate with equipment such as a flex multiweeder or harrow, and follow with a second incorporation at a right angle. Adjust incorporation equipment to a depth so as not to disturb the seed.

NOTE: Do not apply the Far-Go herbicide plus Treflan tank mix to soils which are subject to prolonged periods of flooding.

If less than 25 inches of total water is used to produce the crop, do not plant sorghum or oats for 18 months after application.

Applied according to label directions and under normal growing conditions, Far-Go herbicide plus Treflan will not adversely affect spring wheat and barley. Spring wheat and barley may be injured if seeded through a treated layer of Far-Go herbicide plus Treflan. Seeding disease, cold weather, improper seeding or incorporation depth, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of crop damage from Far-Go herbicide plus Treflan. Under these conditions, delayed crop development or reduced yields may result.

Refer to Treflan label for crop rotation restrictions and cautionary statements.

ANNUAL CANARYGRASS SEED PRODUCTION

FALL TREATMENT: Apply the herbicide at a rate of 1 quart per acre within 3 weeks of soil freezeup. The seeding depth of the crop should be 1 to 2 inches.

SPRING TREATMENT: Incorporate this product according to the **BEFORE SEEDING INCORPORATION** instructions in the **SPRING TREATMENT** section of this label. Apply the herbicide at a rate of 1 quart per acre just before seeding. The seeding depth of the crop should be 1 to 2 inches.