

LETTUCE—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, mites and mites use 1 quart per acre.

SWEET POTATOES—For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites use 1 quart per acre.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 7 DAYS BEFORE HARVEST.

ARTICHOKE—For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers, mites and mites use 1 quart per acre.

BROCCOLI—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 to 3 pints per acre. For cabbage looper, stink bug use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 10 days before harvest.

BRISSELS SPROUTS—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 to 3 pints per acre. For cabbage looper, stink bug use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 10 days before harvest.

CABBAGE—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 to 3 pints per acre. For cabbage looper, stink bug use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 10 days before harvest.

CAULIFLOWER—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 to 3 pints per acre. For cabbage looper, stink bug use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 10 days before harvest.

KOHLRABI—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 to 3 pints per acre. For cabbage looper, stink bug use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 10 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 14 DAYS BEFORE HARVEST.

CUCUMBERS—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 quart per acre. Do not apply before vines.

ONIONS—For control of thrips, use 1 pint per acre.

TOMATOES—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, mites and mites use 1 to 3 pints per acre. For cabbage looper, stink bug use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 15 days before harvest.

TURNIPS—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, leaf miners and mites use 1 to 3 pints per acre. For cabbage looper, stink bug use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 15 days before harvest if tops are to be used for food or feed do not apply closer than 21 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 15 DAYS BEFORE HARVEST IF TOPS OF BEETS ARE TO BE USED AS FOOD OR FEED THE INTERVAL BETWEEN LAST APPLICATION AND HARVEST SHOULD BE INCREASED TO 21 DAYS.

BEANS (Dry and Green)—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, lygus bugs and mites use 1 to 3 pints per acre. For pea curculionid, surface feeding and climbing cutworms and stink bugs use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 10 days before harvest.

BEETS (Red)—For control of aphids, armyworms up to third instar, cabbage loopers, flea beetles, leafhoppers, lygus bugs, mites and stink bugs use 1 quart per acre.

CARROTS—For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites, use 1 quart per acre. Do not feed tops.

CELERY—For control of aphids, armyworms up to third instar, cabbage loopers, flea beetles, leafhoppers, lygus bugs, mites and stink bugs use 1 quart per acre.

CUCUMBERS—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 quart per acre. Do not apply before vines.


ONIONS—For control of thrips, use 1 pint per acre.

PEPPERS—For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites, use 1 quart per acre.

SPINACH—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, mites, pea weevils, cutworms and stink bugs use 1 to 3 pints per acre. For cabbage looper, stink bug use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 10 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 21 DAYS BEFORE HARVEST.

LETTUCE—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, mites and mites use 1 to 3 pints per acre.



CEREAL

BARLEY, OATS, RYE AND WHEAT—For control of aphids, armyworms up to third instar, leafhoppers and winter grain flies use 1 to 3 pints per acre. For cutworms use 1 to 2 pints per acre. For chinch bugs, fall chinch bugs and Say's plant bugs use 1/2 pint per acre. For western or brown wheat mites use 1 pint per acre. Do not use more than 1 pint per acre within 15 days of harvest.

CORN—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 to 3 pints per acre. For corn leaf aphid and mites use 1 pint per acre. For sorghum midges use 1/2 pint per acre. For cutworms use 1 to 2 pints per acre. For chinch bugs use 1/2 pint per acre. For Say's plant bugs use 1/2 pint per acre. For western or brown wheat mites use 1 pint per acre. Do not use more than 1 pint per acre within 15 days of harvest.

RICE—For rice stink bugs use 1/2 pint per acre. Do not apply within 15 days of harvest. Other rice and Say's plant bugs are important pests. These are important pests.

SORGHUM—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 to 3 pints per acre. For sorghum midges use 1/2 pint per acre. For cutworms use 1 to 2 pints per acre. For chinch bugs use 1/2 pint per acre. For Say's plant bugs use 1/2 pint per acre. For western or brown wheat mites use 1 pint per acre. Do not use more than 1 pint per acre within 15 days of harvest.

OIL SEED CROPS

SAFFLOWER—For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and mites use 1 to 3 pints per acre.

FRUIT

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATION CLOSER THAN 14 DAYS BEFORE HARVEST. APPLY AS FREQUENTLY AS NEEDED TO CONTROL INSECTS.

APPLES—For control of aphids, including mites, plum curculionid, scales and red banded leaf rollers use 1/2 pint to 1 quart per 100 gallons of water. For codling moth, plum curculionid and red banded leaf rollers apply at petal fall and 2 to 4 applications, 8 to 14 days thereafter, to maintain control. Do not use more than 6 quarts of this product per acre.

GRAPES—For control of aphids, grape leaf folders and grape leafhopper use 1/2 pint to 1 quart per 100 gallons of water. Do not use more than 1/2 pint of this product per acre.

PEACHES, PLUMS AND PRUNES—For control of aphids, peach tree borers, plum curculionid and Oriental fruit moths use 1/2 pint to 1 quart per 100 gallons of water. Spray tree trunks at time of moth flight to control peach tree borer. For plum curculionid and Oriental fruit moth apply at petal fall and 2 to 4 applications, 8 to 14 days thereafter, to maintain control. Do not use more than 1 gallon of this product per acre.

STRAWBERRIES—For control of aphids use 1/2 pint per acre.

RAPIDLY ABSORBED THROUGH SKIN. REPEATED EXPOSURE MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.

DO NOT GET ON SKIN, IN EYES, ON CLOTHING: Wash hands after touching and if clothes wet with water, wash thoroughly. Protective clothing, water, water, water. Do not eat, drink, or smoke while using this product.

Apply to plants with spray and warm water. Do not bathe or shower.

Bathe immediately after work. Wash all clothing with plenty of soap and water. Remove all traces of methyl parathion. Wash with sudsoy sudsing by giving sudsing sudsoy wash water. Have water run down and keep quiet.

Do not eat, drink, or smoke while using this product. Wash hands after work. Wash all clothing with plenty of soap and water. Remove all traces of methyl parathion. Wash with sudsoy sudsing by giving sudsing sudsoy wash water. Have water run down and keep quiet.

DO NOT BREATHE DUST, VAPOR, OR SPRAY MIST: What a major health hazard to the face caused by the U.S. Department of Agriculture, Methyl Parathion, is a highly toxic insecticide. It is a highly toxic insecticide. It is a highly toxic insecticide. It is a highly toxic insecticide.

Do not breathe dust, vapor, or spray mist. What a major health hazard to the face caused by the U.S. Department of Agriculture, Methyl Parathion, is a highly toxic insecticide. It is a highly toxic insecticide. It is a highly toxic insecticide. It is a highly toxic insecticide.

DO NOT STORE NEAR FOOD OR FEED PRODUCTS: Do not store near food or feed products. Do not store near food or feed products. Do not store near food or feed products. Do not store near food or feed products.

Do not store near food or feed products. Do not store near food or feed products. Do not store near food or feed products. Do not store near food or feed products.

DISPOSAL OF EMPTY CONTAINER: Do not reuse empty container. Do not reuse empty container. Do not reuse empty container. Do not reuse empty container.

Do not reuse empty container. Do not reuse empty container. Do not reuse empty container. Do not reuse empty container. Do not reuse empty container. Do not reuse empty container.

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED METHYL PARATHION POISONING: Call a physician at once in all cases of suspected methyl parathion poisoning. Call a physician at once in all cases of suspected methyl parathion poisoning. Call a physician at once in all cases of suspected methyl parathion poisoning.

Call a physician at once in all cases of suspected methyl parathion poisoning. Call a physician at once in all cases of suspected methyl parathion poisoning. Call a physician at once in all cases of suspected methyl parathion poisoning.

Call a physician at once in all cases of suspected methyl parathion poisoning. Call a physician at once in all cases of suspected methyl parathion poisoning. Call a physician at once in all cases of suspected methyl parathion poisoning.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON: Never give anything by mouth to an unconscious person. Never give anything by mouth to an unconscious person. Never give anything by mouth to an unconscious person.

Never give anything by mouth to an unconscious person. Never give anything by mouth to an unconscious person. Never give anything by mouth to an unconscious person. Never give anything by mouth to an unconscious person.

PHYSICIANS NOTE: This product is a cholinesterase inhibitor. Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

TREATMENT: Large doses of atropine are required for adults (1 mg. to 4 mg. intravenously) and possibly by intravenous injection at 15 minute intervals every 15 to 30 minutes until just before the onset of atropine toxicity. Do not use more than 10 mg. of atropine per 100 lbs. of body weight. Do not use more than 10 mg. of atropine per 100 lbs. of body weight.

WARNING: Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist.

Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist.

Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist. Do not breathe dust, vapor, or spray mist.

Combustible. Do not store or use near heat or open flame. Do not store or use near heat or open flame. Do not store or use near heat or open flame. Do not store or use near heat or open flame.

AID
AN
OF

ominal
or but
hion is

plenty
ion. If
water.

IOUS

ibitor.
in the
vation,
mps.

1/100
every
red in
in age.
essary.
ire to
onged
ibitor
enera-

VEGETABLES

ACTION


BARLEY OATS RYE AND WHEAT

CORN

ALFALFA CLOVER AND VEGETABLES

HERBS

HYPERICUM



WARNING
KEEP AWAY from FEED or FOOD PRODUCT

POISON

CAUTION-DO NOT DRINK
IF LEAKING **DON'T** BREATHE FUMES
DO NOT TOUCH CONTAINER
DO NOT SWALLOW

This is to certify that the contents of this package are properly described by name and are packed and marked and are in proper condition for transportation according to the Regulations prescribed by the Interstate Commerce Commission.

MANUFACTURED BY CHEMICAL COMPANY

Shipper's name required
reason for shipment by
EXPRESS

ACCEPTED
FEB 13 1964