

# METHYL PARATHION 4 E

INSECTICIDE

Emulsifiable concentrate for controlling certain insects on the listed field, forage, fruit and vegetable crops.

Use only according to label instructions.

READ "WARNING NOTICE" BEFORE BUYING OR USING.

IF NOTICE TERMS ARE NOT ACCEPTABLE RETURN AT ONCE UNOPENED.

Keep out of reach of children.

**DANGER!** POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING.

RAPIDLY ABSORBED THROUGH SKIN.

REPEATED EXPOSURE MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.

Do not get on skin, in eyes, on clothing.

Do not breathe dust, vapor, or spray mist.

Do not store near food or feed products.

Do not use or store near heat or open flame.

In case of fire, use water spray, foam, dry chemical or CO<sub>2</sub>.

**POISON**

See antidote statement and other required warning statements on side and back panels.

ACTIVE INGREDIENTS:

Keep out of reach of children.

**DANGER!**

POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING. RAPIDLY ABSORBED THROUGH SKIN.

REPEATED EXPOSURE MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.

DO NOT GET ON SKIN, IN EYES, ON CLOTHING.

Wear heavy natural rubber gloves and goggles. Wear clean waterproof or freshly-laundered protective clothing (coveralls, rubber boots, cap, etc.). Destroy and replace gloves frequently.

Wash thoroughly with soap and warm water before eating or smoking.

Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use.

**POISON**

ATROPINE IS AN ANTIDOTE. CONSULT PHYSICIAN FOR EMERGENCY SUPPLY.

FIRST AID

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED METHYL PARATHION POISONING.

If symptoms or signs of poisoning include blurred vision, abdominal cramps, and tightness in the chest, do not wait for a doctor but give two atropine tablets (each 1/100 grain or 0.65 milligrams) at once (do not give atropine tablets to children under five years of age.)

Move patient immediately from the area where methyl parathion is present.

Remove contaminated clothing and wash thoroughly clean with plenty of soap and water to remove all traces of methyl parathion. If swallowed, induce vomiting by giving salty or soapy warm water. If victim lie down and keep quiet.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

PHYSICIANS NOTE: This product is a cholinesterase inhibitor. Warning symptoms include weakness, headache, tightness in the chest, blurred

**INCREASINGLY HAZARDOUS.**

- Do not get on skin, in eyes, on clothing.
- Do not breathe dust, vapor, or spray mist.
- Do not store near food or feed products.
- Do not use or store near heat or open flame.
- In case of fire, use water spray, foam, dry chemical or CO<sub>2</sub>.



**POISON**

See antidote statement and other required warning statements on side and back panels.

**ACTIVE INGREDIENTS:**

*O,O-dimethyl O-p-nitrophenyl phosphorothioate	45.7%
Aromatic petroleum derivative solvent	48.2%

<b>INERT INGREDIENTS:</b>	6.1%
	<hr/>
	100.0%

\*Equivalent to 4.0 lbs. of 100% methyl parathion per gallon.

**NET CONTENTS 5 U.S. GALLONS**

Wear heavy natural rubber gloves and goggles. Wear clean waterproof or freshly-laundered protective clothing (coveralls, rubber boots, cap, etc.). Destroy and replace gloves frequently.

Wash thoroughly with soap and warm water before eating or smoking.

Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water; for eyes, flush with water for 15 minutes.

**DO NOT BREATHE DUST, VAPOR, OR SPRAY MIST:** Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for METHYL PARATHION protection. Airplane pilots should wear full-

methyl parathion is present.

Remove contaminated clothing and wash thoroughly with clean water. Wash skin with plenty of soap and water to remove traces of methyl parathion. If vomiting occurs, induce vomiting by giving salty or soapy water. If victim lies down and keep quiet.

**NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

**PHYSICIANS NOTE:** This product is a cholinesterase inhibitor. Warning symptoms include headache, tightness in the chest, sweating, salivation, non-reactive pin-point pupils, nausea, vomiting, diarrhea, and ataxia.

**TREATMENT:** Large doses of atropine should be given every 10 to 15 minutes until pupils dilate. If 2-PAM Chloride (pralidoxime) is available, administer according to directions. Never give morphine.

# DIOSIN

IS AN ANTIDOTE.  
PHYSICIAN FOR  
EMERGENCY SUPPLY.

## AT ONCE IN ALL CASES OF PARATHION POISONING.

Signs of poisoning include blurred  
vision, dizziness, weakness, nausea,  
vomiting, diarrhea, sweating, tightness  
in the chest, and slow pulse. Give  
atropine (1 mg per 0.65 milligrams)  
at once. (One tablet under five  
years of age.)  
Remove the person immediately  
from the area where poisoning  
is present.

Remove contaminated clothing  
and wash the skin with soap  
and water to remove all traces  
of parathion. If swallowed, induce  
vomiting with salty or soapy  
warm water. Have the person  
keep quiet.

EVERYTHING BY MOUTH TO AN  
PERSON

The buyer and all users are deemed to have accepted the terms of this notice which may not be varied by any verbal or written agreement.

## TO PREVENT PERSONAL INJURY AND POSSIBLE FATALITIES:

Keep all persons and animals out of treated areas for 48 hours.

Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated.

Toxic to fish and wildlife. Birds and other wildlife in treated area may be harmed. Do not contaminate lakes, ponds, streams, woodlands, and other non-crop areas.

## DIRECTIONS FOR USE.

Be sure to read the precautionary statements before

of handpicking cotton.

**GRASS (Forage)**—For control of armyworms up to third instar, ~~crested wheat bugs, false chinch bugs, grasshoppers and leafhoppers~~, use 1½ pints per acre.

**HOPS**—For control of ~~and~~ spider mites, use 1 quart per acre. Do not use closer than 15 days before harvest.

**SOYBEANS**—For control of cutworms and three-cornered alfalfa hoppers, use ½ pint per acre. For control of garden webworms, use ¼ to 1 pint per acre. For control of aphids, stink bugs, velvet bean caterpillars, Mexican bean beetles, blister beetles and two-spotted mites, use 1 pint per acre. For bollworms, cabbage loopers, fall armyworms up to third instar, and green cloverworms, use 1 quart per acre. Do not apply closer than 20 days before harvest or grazing.

**SUGAR BEETS**—For aphids, armyworms up to third instar, flea beetles, leafhoppers, Lygus bugs, stink bugs and webworms, use ¾ pint per acre. Do not treat closer than 15 days before harvest.

**TOBACCO**—For control of green peach aphids, use ½ pint per acre. For control of surface feeding or climbing cutworms, use ¼ to 1 pint per acre.

**PEACHES, PEARS AND PRUNES**—For control of aphids, peach tree borers, plum curculio and Oriental fruit moths, use ½ pint to 1 quart per 100 gallons of water. Spray tree trunks at time of moth flight to control peach tree borers. For plum curculio and Oriental fruit moths, apply at petal fall or shuck split and 3 to 4 applications, 8-14 days thereafter, to maintain control. Do not use more than 1 gallon of this product per acre.

**STRAWBERRIES**—For control of aphids, use 1 to 1½ pints per acre.

## VEGETABLES

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 5 DAYS BEFORE HARVEST.

**POTATOES**—For control of aphids, armyworms up to third instar, cabbage loopers, false chinch bugs, flea beetles, leafhoppers, mites and shield bugs, use 1½ quarts per acre.

**SWEET POTATOES**—For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites,

hoppe  
crown  
pint sh  
have

**TOMATOES**  
third  
pers,  
Rates  
15 da

**TURNIPS**  
third  
pers,  
acre.  
than

**APPL**  
**FOLL**  
**CLOS**

**IF TO**  
**ARE**  
**INTE**

**HARV**  
**DAYS**  
**BEAN**