

RESTRICTED USE PESTICIDE

For restricted use only. Do not use on crops or plants unless specifically labeled for that use. Do not use on crops or plants unless specifically labeled for that use.

METHYL PARATHION 4 E

INSECTICIDE

Emulsifiable insecticide for controlling certain insects on the listed field, forage, fruit and vegetable crops.

Use only according to label directions.

READ LIMIT OF WARRANTY AND LIABILITY BEFORE USING. IF TERMS ARE NOT ACCEPTABLE, DO NOT USE.

POISON

See side panel for antidote and precautions




DANGER

Keep out of reach of children

ACTIVE INGREDIENT

40% Dimethyl 4-methylphosphorothioate 1.00%
Aromatic petroleum solvents 1.00%
INERT INGREDIENTS 1.00%

*Equivalent to 40% by weight methyl parathion per gallon



Keep out of reach of children

DANGER!

POISONOUS BY SKIN CONTACT
INHALATION OR SWALLOWING
RAPIDLY ABSORBED THROUGH SKIN

REPEATED EXPOSURE MAY WITHOUT SYMPTOMS BE INCREASINGLY HAZARDOUS

DO NOT GET ON SKIN, IN EYE, OR ON CLOTHING. Wash thoroughly with soap and water before eating, drinking, or using the toilet.

DO NOT BREATH DUST, VAPOR, OR SPRAY MIST. Keep container closed when not in use.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

DO NOT BREATHE DUST, VAPOR, OR SPRAY MIST. Keep container closed when not in use.

DO NOT USE NEAR FOOD OR FEED PRODUCTS

NET GAL.

POISON

ATROPINE IS AN ANTIDOTE
CONSULT PHYSICIAN FOR
EMERGENCY SUPPLY

FIRST AID: If swallowed, give 1-2 glasses of water. If inhaled, get fresh air. If on skin, wash with soap and water.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

DISPOSAL OF EMPTY CONTAINER

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

100

DII

POISON
ATROPINE IS AN ANTIDOTE
CONSULT PHYSICIAN FOR
EMERGENCY SUPPLY

FIRST AID
In case of poisoning, call your physician or the nearest hospital. Do not induce vomiting unless directed to do so by a physician. If necessary, give 1-2 glasses of water. Do not give anything by mouth to an unconscious person.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON

Keep this and all other medicines out of the reach of children. Do not use if the cap is missing or the seal is broken. Do not use if the medicine is discolored or has an unusual odor.

Do not use if the medicine is discolored or has an unusual odor. Do not use if the cap is missing or the seal is broken.

POISON OF EMERGENCY USE

ATROPINE

ATROPINE IS A FAST ACTING POISON. The following table gives complete directions for use.

OIL SEED CROPS
ATROPINE SUNFLOWER

FIELD AND FORAGE CROPS

BARLEY
BARNYARD
CORN
GRASS
HAY
RICE
SUGAR BEET
WHEAT

APPLES
CHAPPEL
PEACHES

FRUIT


PLUM
PRUNE
STRAWBERRIES

VEGETABLES

ARTICHOKES
BEANS
Dry and Green
BEETS Raw
BROCCOLI
BRUSSELS
SPROUTS
CABBAGE
CARROTS
CAULIFLOWER
CELERY
CUCUMBERS

ASPARAGUS
KOHLEWURST
LETTUCE
MUSTARD
ONIONS
PEAS
PEPPERS
POTATOES
RUTABAGAS
SPINACH
SWEET POTATOES
TOMATOES
TURNIPS

IMMEDIATE ACTION
Color: Red and Black



DIRECTIONS FOR USE

POISONOUS TO WILDLIFE
This product is toxic to wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning equipment or disposing of wastes.

The poison is highly toxic to many species of fish. Do not use in or near water. Do not use in or near streams, rivers, or other bodies of water.

ATTENTION
Do not use in or near water. Do not use in or near streams, rivers, or other bodies of water. Do not use in or near water. Do not use in or near water. Do not use in or near water.

Do not use in or near water. Do not use in or near streams, rivers, or other bodies of water. Do not use in or near water. Do not use in or near water. Do not use in or near water.

Combustible. Do not store or use near heat or open flame. Do not use in or near water. Do not use in or near water.

IN CASE OF EMERGENCY, CALL 1-800-4-A-GRAMA OR YOUR NEAREST POISON CONTROL CENTER.

AT-1-15004-53 AT-1-15004-53

ATROPINE

RESTRICTED USE PESTICIDE

For restricted use and use only by trained applicators or persons under their direct personal supervision only for those uses covered by the Certified Applicator's Certificate above.

METHYL PARATHION 4 E

Resealable Insecticide for controlling certain insects on the listed field, vegetable, fruit and vegetable crops.

NOT FOR HOME USE

Complete Directions for Use.

Use only according to these label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

RESEALABLE BAG

Pull flaps apart to open. Press along ridge to close.

04-11 150 05/50

LIMIT OF WARRANTY AND LIABILITY

Monsanto Company warrants that this insecticide is reasonably fit for the purposes referred to in the directions for use. This product is sold on the basis of the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as, for example, incompatibility with other pesticides, the manner of its use or application, or the presence of other products or materials in or on the soil or crop. MONSANTO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY. The exclusive remedy of the user or buyer, and the limit of the liability of Monsanto Company or any other seller for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this notice which may not be varied by any verbal or written agreement.

ATTENTION

Do not use in any manner other than recommended on this label. To avoid excessive residues of methyl parathion on food or forage crops, always observe the statements found under "Directions for Use," limiting the time before harvest when methyl parathion may be applied.

VEGETABLE INSECTS

ARTICHOKE 16
BEANS 17
Dry and Green 19
BEETS Red 19
BROCCOLI 16
BRUSSELS 16
SPROUTS 17
CABBAGE 18
CARROTS 19
CAULIFLOWER 17
CELERY 19
COLLARDS 18
CUCUMBERS 19

FIELD AND FORAGE CROP INSECTS

ALFALFA 12
CLOVER 12
COTTON 12
GRASS (Hay) 12
HOPS 12

FRUIT INSECTS

APPLES 14
GRAPES 14
PEACHES 15

OIL SEED CROP INSECTS

SUNFLOWER 15
SUNFLOWER 15

POISONOUS TO WILDLIFE

This product is toxic to wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning equipment or disposal of wastes. Combustible: Do not store or use near heat or open flame. In case of fire, use water spray, foam, dry chemical or CO₂.

Keep out of reach of children

DANGER! POISON

See all the statements and other required warning statements on pages 6, 7 and 8.
POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING.
RAPIDLY ABSORBED THROUGH SKIN.
REPEATED EXPOSURE MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.
Do not get on skin, in eyes, on clothing.
Do not breathe dust, vapor, or spray mist.
Do not store near food or feed products.
Do not use or store near heat or open flame.
In case of fire, use water spray, foam, dry chemical or CO₂.

ACTIVE INGREDIENTS

*O,O dimethyl O-p-nitrophenyl phosphorothioate 45.2%
Aromatic petroleum derivative solvent 48.8%
INERT INGREDIENTS 5.7%
100.0%

*Equivalent to 4.0 lbs. of 100% methyl parathion per gallon.

Directions

ENVIRONMENTAL PRECAUTIONS

Hazards and Ingredients

Hazards and Precautions

Keep out of reach of children.

DANGER!

POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING.

RAPIDLY ABSORBED THROUGH SKIN.

REPEATED EXPOSURE MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS.

DO NOT GET ON SKIN, IN EYES, ON CLOTHING

Wear heavy natural rubber gloves and goggles. Wear clean, waterproof or freshly laundered protective clothing (coveralls, rubber boots, cap, etc.). Destroy and replace gloves frequently.

Wash thoroughly with soap and warm water before eating or smoking.

Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before reuse.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water. For eyes, flush with water for 15 minutes.

DO NOT BREATHE DUST, VAPOR, OR SPRAY MIST

Wear a mask or respirator of a type passed by the Mine Safety and Health Administration for METHYL PARATHION protection. Replace filters should wear to have another type mask.

It has been found that mechanical exhaust ventilation.

Do not apply or allow drift to areas occupied by unprotected humans or beneficial animals.

DO NOT STORE NEAR FOOD OR FEED PRODUCTS

Keep out of reach of children and domestic animals.

Not for use or storage in or around the home.

Do not spillage, clean up area with strong detergent.

POISON

ATROPINE IS AN ANTIDOTE
CONSULT PHYSICIAN FOR
EMERGENCY SUPPLY

FIRST AID

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED METHYL PARATHION POISONING.

If symptoms or signs of poisoning are observed, such as: abdominal cramps, and tightness in the chest, do not wait for a doctor but give two atropine tablets (each 1/100 grain or 0.65 mg. grams) at once. (One tablet to children under five years of age.)

Repeat every 10 to 15 minutes until pupils dilate. In addition to atropine, at 2 P.M. Chloroform (paraffin-free) is available, administer according to manufacturer's directions. Have good emphysema (free chest) for prompt drainage. Artificial respiration or oxygen administration may be necessary if severe patient continuously 48 hours. Repeat for persistent cholinesterase inhibitors may, without warning, cause profound susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Precautions: 1. If used in...

Directions for Use

See all the statements and other required warning statements on pages 6, 7 and 8.

DIRECTIONS FOR USE

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

See all the statements and other required warning statements on pages 6, 7 and 8.

CEREAL

BARLEY, OATS, RYE AND WHEAT — For leafhoppers use 1/2 to 1 pint per acre. For aphids, armyworms up to third instar, leafhoppers and winter grain moths use 1/2 to 1 1/2 pints per acre. For climbing cutworms use 1/2 to 1 pint per acre. For chinch bugs, fall chinch bugs and Say's plant bugs use 1 1/2 pints per acre. For western or brown wheat moths use 1 pint per acre. Do not use more than 1/2 pint per acre within 15 days of harvest.

CORN — For control of armyworms up to third instar, climbing cutworms, corn leaf aphids, corn rootworm adults, fall armyworms and stink bugs use 1/2 pint per acre. Do not apply within 12 days of harvest.

RICE — For rice stink bugs use 1/2 to 1 pint per acre. Do not apply within 15 days of harvest. Stink bugs and crickets may be killed. Do not apply where there are important resources.

SORGHUM — For corn leaf aphids and moths use 1 pint per acre. For sorghum midges use 1 pint to 1 quart per acre. 2 applications, 3 to 5 days apart when approximately 90% of the heads have completely emerged from the husk or not later than start of blooming. Do not apply closer than 21 days before harvest. Leaf injury may occur on some hybrid varieties of sorghum. Spray a few rows a week or so before heading to test effects upon plants.

Cereal Insects

11

FIELD AND STORAGE CROPS

FIELD AND STORAGE CROPS

ALFALFA AND CLOVER — For control of alfalfa and clover aphids (including spotted alfalfa aphid) and climbing cutworms use 1/2 to 1 pint per acre. For alfalfa caterpillars, alfalfa cutworms, armyworms up to third instar, clover leaf beetles and webworms use 1 pint per acre. For Egyptian alfalfa weevils, leafhoppers, eggs bugs and spider mites use 1 to 2 pints per acre. Do not treat closer than 15 days before harvest, cutting or grazing. In California and Nevada do not use more than 1/2 pint per acre.

COTTON — For control of thrips use 1/2 to 1 pint per acre. For cotton leafworms use 1/2 to 1 pint per acre. For grasshoppers and fall armyworms up to third instar use 1/2 pint per acre. For spider mites (does not control all species) use 1/2 to 1 pint per acre. For control of leafhoppers, eggs bugs and other moths use 1/2 to 1 quart per acre. For boll weevils, aphids and garden webworms use 1/2 to 1 pint per acre. For false chinch bugs use 1 pint per acre. For cabbage loopers and cutworms use 1 to 2 pints per acre. For bollworms, salt marsh caterpillars, armyworms up to third instar, cotton leaf perforators and stink bugs use 1 pint per acre. Applications should be made at 4 to 5 day intervals until control is obtained. At above dosages, application may be made up to the day before harvest, if harvest will be with mechanical pickers.

CAUTION — When necessary to enter treated cotton fields within 24 hours after application, protective clothing should be worn. Do not apply within 5 days of handpicking cotton.

FIELD AND STORAGE CROPS

For control of alfalfa and clover aphids (including spotted alfalfa aphid) and climbing cutworms use 1/2 to 1 pint per acre.

For control of alfalfa caterpillars, alfalfa cutworms, armyworms up to third instar, clover leaf beetles and webworms use 1 pint per acre.

For control of Egyptian alfalfa weevils, leafhoppers, eggs bugs and spider mites use 1 to 2 pints per acre. Do not treat closer than 15 days before harvest, cutting or grazing. In California and Nevada do not use more than 1/2 pint per acre.

For control of thrips use 1/2 to 1 pint per acre. For cotton leafworms use 1/2 to 1 pint per acre. For grasshoppers and fall armyworms up to third instar use 1/2 pint per acre. For spider mites (does not control all species) use 1/2 to 1 pint per acre.

For control of leafhoppers, eggs bugs and other moths use 1/2 to 1 quart per acre. For boll weevils, aphids and garden webworms use 1/2 to 1 pint per acre.

For false chinch bugs use 1 pint per acre. For cabbage loopers and cutworms use 1 to 2 pints per acre.

For bollworms, salt marsh caterpillars, armyworms up to third instar, cotton leaf perforators and stink bugs use 1 pint per acre.

Applications should be made at 4 to 5 day intervals until control is obtained. At above dosages, application may be made up to the day before harvest, if harvest will be with mechanical pickers.

Field and Storage Insects cont'd

12

Vegetable Insects cont'd

13

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS MAKE NO APPLICATIONS CLOSER THAN 10 DAYS BEFORE HARVEST

CABBAGE — For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and moths use 1 to 3 pints per acre. For cabbage loopers and stink bugs use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

COLLARDS, KALE AND MUSTARD — For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and moths use 1 to 3 pints per acre. For cabbage loopers and stink bugs use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

PEAS — For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, eggs bugs and moths use 1 to 2 pints per acre. For cabbage loopers, corn earworms, cutworms, feeding and climbing cutworms and stink bugs use 2 pints per acre. Rates above 1 pint should not be applied closer than 15 days before harvest.

TOMATOES — For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, moths and eggplants use 1 to 3 pints per acre. For cabbage loopers use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 15 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS MAKE NO APPLICATIONS CLOSER THAN 14 DAYS BEFORE HARVEST

ANS (Dry and Green) — For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, eggs bugs and moths use 1 to 3 pints per acre. For grasshoppers, surface feeding and climbing cutworms and stink bugs use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

BEETS (Red) — For control of aphids, armyworms up to third instar, cabbage loopers, flea beetles, leafhoppers, eggs bugs, moths and stink bugs use 1 to 3 pints per acre. If tops are to be used for food use 1/2 pint per acre. Do not apply closer than 21 days before harvest.

CARROTS — For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers and moths use 1 to 3 pints per acre. Do not apply closer than 21 days before harvest.

CITRUS — For control of aphids, armyworms up to third instar, cabbage loopers, flea beetles, leafhoppers, eggs bugs, moths and stink bugs use 1 to 3 pints per acre.

CUCUMBERS — For control of aphids and two-spotted spiny-legged beetles use 1/2 to 1 pint per acre. Do not apply before harvest.

PEPPERS — For control of aphids use 1/2 to 1 pint per acre.

PLUMS AND PRUNES — For control of aphids, peach tree borers, plum curculion and Oriental fruit moths use 1/2 pint to 1 quart per 100 gallons of water. Spray tree trunks at time of moth flight for control of peach tree borers. For plum curculion and Oriental fruit moths apply at petal fall or shortly after and 3 to 4 applications, 8 to 10 days thereafter to maintain control. Do not use more than 1 gallon of this product per acre.

STRAWBERRIES — For control of aphids use 1 to 1 1/2 pints per acre.

OIL SEED CROPS

SUNFLOWER — For aphids and eggs bugs use 1 quart per acre before flowering. Do not apply after flowering.

SUNFLOWER — For control of sunflower moth use 1 quart per acre. Make no more than 3 applications at 5 day intervals. Do not apply within 30 days of harvest.

524-273

Field and Oil Seed Insects

15

Vegetable Insects

16

VEGETABLES

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS MAKE NO APPLICATIONS CLOSER THAN 5 DAYS BEFORE HARVEST

POTATOES — For control of aphids, armyworms up to third instar, cabbage loopers, false chinch bugs, flea beetles, leafhoppers, moths and shield bugs use 1 1/2 quarts per acre.

SWEET POTATOES — For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers and moths use 1/2 quart per acre.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS MAKE NO APPLICATIONS CLOSER THAN 7 DAYS BEFORE HARVEST

RATICHOLE — For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers, moths and milk-hole plume moths use 1 quart per acre.

BROCCOLI — For control of aphids, armyworms up to third instar, flea beetles, leafhoppers and moths use 1 to 3 pints per acre. For cabbage loopers and stink bugs use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

BEST DOCUMENT AVAILABLE

10-10-10-10-10-10

Vegetable Insects cont'd

20

PEPPERS — For control of aphids, armyworms up to third instar, surface feeding and climbing cutworms, flea beetles, leafhoppers and moths use 1 quart per acre.

SPIRACHE — For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, moths, seed corn maggots in crown and crown moths use 1 to 2 pints per acre. For cabbage loopers use 2 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

TURRIPS — For control of aphids, armyworms up to third instar, flea beetles, leafhoppers, leaf miners and moths use 1 to 1 1/2 pints per acre. For cabbage loopers use 1 1/2 pints per acre. If tops are to be used for food or feed do not apply closer than 21 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS MAKE NO APPLICATIONS CLOSER THAN 21 DAYS BEFORE HARVEST

LETTUCE — For control of aphids, armyworms up to third instar, cabbage loopers, flea beetles, leafhoppers and moths use 1 quart per acre.

996 11 548 69-31

EPA Reg No 524-273

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63166 U.S.A.