


41



HERBICIDE BY **Monsanto**

Emulsifiable herbicide for  
weed control in corn, soybeans,  
peanuts, dry beans, lima beans,  
red kidney beans, potatoes and  
crops listed.

**Complete Directions for Use**

U.S. Pat. No. 3,429,945 - EPA Reg. No. 524-285-AA

Read the entire label.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before  
buying or using. Terms are not acceptable return  
at once unopened.

**RESEALABLE BAG**

Flip flaps to open.  
Fold along ridge to close.

1977-3

\*Lasso EC is referred to throughout these  
instructions as "Lasso"

Precautionary Statements

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals

Keep out of reach of children

**DANGER!**

CAUSES EYE BURNS AND SKIN IRRITATION

HARMFUL IF SWALLOWED

MAY CAUSE ALLERGIC REACTION

Do not get in eyes, on skin or on clothing  
Wear goggles or face shield and gloves when handling

Wash thoroughly after handling

FIRST AID: In case of eye contact, flush eyes or skin immediately with water for 15 minutes. Remove contaminated clothing. For eye contact, call a physician immediately before reuse.

Physical or Chemical

**COMBUSTIBLE**

Do not use or store near heat or open flame  
Use only with adequate ventilation

In case of  
FIRE: use water spray to smother chemical or CO  
SPILL or LEAK: flush area with water spray

Preplant Incorporated Treatment — Apply 2.5 to 3 quarts of Lasso plus one half quart of Trellan in

Storage and Disposal

STORE ABOVE 32 F TO KEEP PRODUCT IN SOLUTION

Below 32 F crystals may form and settle to the bottom. If crystals form, place in a warm room (72 F) and roll and shake the can frequently for several days to re-suspend before use.

Avoid contamination of feed and foodstuffs

Do not reuse containers. Destroy when empty.

ACTIVE INGREDIENT

\*Atrazine 43.0%

INERT INGREDIENTS 57.0%

\*Contains 4 pounds of active ingredient per gallon (methoxy methyl atrazine per gallon)

GENERAL INFORMATION

Lasso herbicide is for the control of most annual grasses and certain broadleaf weeds. Except for minor weeds, treatment on tillage is best. The seed bed should be firm and free of clods and trash. Application should be made within 7 days after the last tillage for weed control. Do not apply when conditions favor drift.

Moisture is required to activate this product. Although the amount required depends primarily upon the amount of soil moisture in the weed germination

Storage and Disposal, Ingredients and General Information




POUR FROM OPENING

CUT OFF CORNER

MINIATURE FACSIMILE  
Smallest Type size on package 10 point.



Granular



HERBICIDE BY

Monsanto

ACCEPTED  
May 1, 1975  
Monsanto Company  
St. Louis, Missouri

Selective herbicide for preemergence weed control  
in corn, soybeans and peanuts

Read the entire label.  
Use only according to label instructions.

Read "TIME OF APPLICATION AND ABILITY"  
before buying or using. Do not use if not ac-  
ceptable in form at time of purchase.

Keep out of reach of children.

**CAUTION:**  
IRRITATING TO EYES.  
MAY BE DANGEROUS IF SWALLOWED.  
MAY CAUSE ALLERGIC SKIN REACTION.

Avoid contact with eyes, skin and clothing.  
Avoid breathing dust.

In case of contact with skin, wash with plenty  
of water.

Wash thoroughly after handling.  
Laundry clothing before use.  
Avoid contact with animal feed and food  
stuffs.

NET WEIGHT **50** LBS

ACTIVE INGREDIENT  
Algarol 2-(4-chlorophenyl)-2-methylpropane  
(methylamethyl) sulfonamide 10%  
INERT INGREDIENTS 90%  
100%

U.S. Patent 3,442,045

EPA Reg. No. 514-097 AA

In case of emergency, call this product  
"CAPTAIN" and give number 514-097 AA.  
Monsanto Company, Agricultural Division  
ST. LOUIS, MISSOURI 63166 U.S.A.