

DLU - 1035

Dr. Nancy Yates-Parker  
Monsanto Company  
700 14th Street, N.W., Suite 1100  
Washington, D.C. 20005

Dear Dr. Yates-Parker:

Subject: Fargo Granular Herbicide (Restriction of Label to Certain States)  
EPA Registration No. 524-292  
Your Application Dated May 26, 1993

The labeling submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable. Please submit five (5) copies of your final printed labeling before you release the product for shipment.

Note that this labeling does not bear the required Worker Protection statements as outlined in PR Notices 93-7 and 93-11. We note that you have submitted labels in accordance with the instructions in those Notices. The new protective language in those submissions must be melded into this label for any shipments released after the deadlines that have been established. You may opt to do so immediately if you are electing the self-verification option, or after acceptance by the special Agency Worker Protection Standard review group. Finished printed labels should be submitted after acceptance by the WPS group and in accordance with their instructions.

Sincerely,

*UKW*  
Robert J. Taylor  
Product Manager 25  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

**BEST AVAILABLE COPY**

2 of 10

**GRANULAR**

**FAR-GO®**

Herbicide by Monsanto

Granular selective herbicide for fall or spring application to control wild oats in winter wheat, spring and durum wheat, barley, triticale, green peas, field dried peas and lentils, and for fall application to suppress *Bromus* species (*B. tectorum*, *B. secalinus* and *B. japonicus*) in winter wheat and in winter barley.

**Complete Directions for Use**

Far-Go® is a registered trademark of Monsanto Company.

EPA Reg. No. 524-292-AA

©MONSANTO COMPANY 1989

1989-1

843.53-000.45/53

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
JEC - 9 1993  
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

524-292

For use in Idaho, Kansas, Minnesota, Missouri, Nebraska, North Dakota, Oregon, South Dakota, Washington and Wyoming.

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

**LIMIT OF WARRANTY AND LIABILITY**

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in

any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

**PRECAUTIONARY STATEMENTS**

---

**Hazards to Humans  
and Domestic Animals**

---

Keep out of reach of children.

**CAUTION!**

MAY CAUSE EYE IRRITATION.

Avoid contact with skin, eyes or clothing.

Wash thoroughly with soap and water  
after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water for  
at least 15 minutes. Call a physician.

---

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

---

**Environmental Hazards**

---

Avoid direct application to any body of water. Do not contaminate  
water when disposing of equipment washwaters.

**ACTIVE INGREDIENT:**

S-(2,3,3-Trichloroallyl)-

diisopropylthiocarbamate ..... 10.0%

INERT INGREDIENTS: ..... 90.0%

100.0%

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

**Storage and Disposal**

Do not contaminate water, foodstuffs, feed or seed by storage and disposal.

**STORAGE:**

Keep bag closed to prevent spills and contamination.

**DISPOSAL:**

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

This herbicide is recommended for wild oat control in barley, peas, lentils, durum, spring and winter wheat and triticale only, and for suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*) in winter wheat and winter barley. Other crops should not be treated with this product because injury may occur. For barley, durum and spring wheat and triticale, this product may be applied on the soil either in the fall or in the spring before wild oats germinate. For lentils and peas, this product may only be applied in the spring before wild oats germinate. For winter wheat, this product may only be applied in the fall before wild oats germinate.

Application equipment must be properly calibrated: application of too much herbicide may injure the crop; application of too little may result in poor wild oat control. Recommended rates should be followed in order to:

1. Avoid crop injury
2. Avoid crop residue at harvest
3. Control wild oats
4. Suppress *Bromus* species

Application to a field which is wet, lumpy, rough or ridged will result in reduced wild oat control and promote crop thinning. Incorporation must be completed within 48 hours after application and before germination of the wild oats. If weeds, including wild oats, have emerged prior to treating and/or planting, they must be controlled. All deep tillage by cultivators or double disc implements must be completed prior to application. Do not plow following application of this herbicide.

Seeding may be done either before or after application, depending upon the crop that is to be sown. If seeding is delayed, shallow reworking of the treated area before seeding will not destroy the effects of the chemical.

When using this herbicide, a strip should be left untreated for proof of results. Weed control may be evaluated also by removing a surface inch or two of the soil at the time of germination to inspect the number of wild oats that were killed before emergence.

Wild oats are usually killed before emergence, but occasionally, and particularly under dry conditions, plants may reach the 3 to 4-leaf stage before they die.

Under conditions of prolonged high temperature at the time of germination, or extreme drought in the spring, this product may not maintain the usual high standard of wild oat control.

**ATTENTION**

DELAYED EMERGENCE, STAND REDUCTION, STUNTING AND YIELD LOSS MAY RESULT-DUE TO COLD OR WET CONDITIONS, IRRIGATION DURING GERMINATION OR EMERGENCE, DEEP PLANTING, SOIL CRUSTING, DISEASE, INSECTS, OR SELECTION OF A VARIETY GENETICALLY SUSCEPTIBLE TO STRESS. THESE ARE CONDITIONS BEYOND THE CONTROL OF MONSANTO, AND THE GROWER SHOULD CONSULT THE STATE UNIVERSITY EXTENSION SPECIALIST OR SEED PRODUCER OR SUPPLIER FOR LOCAL VARIETAL INFORMATION AND RECOMMENDED TILLAGE AND PLANTING PRACTICES AND DATES.

Domestic oats should not be seeded where this product was used the previous year.

Do not graze livestock on treated crops.

IN THE STATE OF MONTANA, do not use this product on fields to be seeded to hard red spring wheat with press drills, if the field is or will be irrigated in the current growing season.

Do not use postplant incorporation with hoe drills. This practice will result in unacceptable seeding depth and an excessive concentration of this product in the seed zone.

Use of this product not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

**HOW TO APPLY AND CALIBRATE EQUIPMENT**

This product must be applied through a specially designed ground applicator or airplane capable of applying small quantities of granules evenly.

**BEST AVAILABLE COPY**

**GROUND EQUIPMENT:** It is important that the applicator be calibrated properly to deliver the desired amount of this product to avoid applying too little, or too much, material. To give even distribution, scatter plates (similar to those used for applying granules in a band) must be attached to each delivery tube or outlet in such a manner to give overall coverage. To calibrate, attach a collector pan, or strong plastic bag, over each spreader plate or delivery tube. Operate over normal terrain to be treated at 4 to 5 miles per hour. Collect the granules from all outputs after covering the desired distance.

**For example:** If a 14-foot applicator is being used, stake off a distance of 324 feet in the field to be treated (distance equal to 1/10 acre treated). After attaching a bag, or collection pan, to each outlet, collect the granules while driving the desired distance. Check to see that each outlet disperses the same amount of granules. Combine all samples and weigh. For the above set of conditions, the quantity of granules that should be collected for the following recommended rates of broadcast treatment are:

RECOMMENDED RATE PER ACRE

10 lbs    12.5 lbs    15 lbs

|                                  |      |          |         |
|----------------------------------|------|----------|---------|
| Proper amount<br>to be collected | 1 lb | 1.25 lbs | 1.5 lbs |
|----------------------------------|------|----------|---------|

If more or less than the desired quantity is collected, adjust opening accordingly and again collect the granules while driving the staked-off distance. Continue this procedure until the proper amount is delivered.

**AIRPLANE:** For aerial applications, attachments designed for applying low volumes of granules must be used. In order to ensure uniform aerial application, it is recommended that the field distribution pattern is checked and any necessary gate and vent modifications are made to ensure an even pattern distribution. In order to ensure uniform application and to avoid overlapping and possible crop injury, it is recommended that two flagmen, one at each end of the field, be used.

**FIELD PREPARATION**

Before applying this product, be sure the soil is in good working condition. All deep tillage by cultivation, or double disc implements, must be completed prior to application of this product. If stubble ground is being treated, one or two passes with a field cultivator may be required before application and incorporation.

**INCORPORATED APPLICATIONS**

When summer fallow or plowed ground is being treated, a field cultivator will provide adequate incorporation. For proper incorporation of Far-Go, set incorporation implement to work the soil no deeper than 3 to 4 inches. Do not use disc implements for incorporation.

7 of 10

**FALL**—On summer fallow ground which is loose and free of lumps and trash, incorporate this product using equipment similar to a culti-harrow or a duckfoot with rod weeder attachment.

For applications to fields or standing stubble, fields may be worked once or twice with a field cultivator or chisel plow. Apply granules and incorporate with a field cultivator or culti-harrow.

If soil must be ridged after incorporating to prevent soil erosion by high winds, the depth of ridging should be kept to a minimum.

In preparing the herbicide-treated area for seeding in the spring, care must be taken to avoid working the soil any deeper than fall tillage.

When incorporating with a single pass in the fall, a second incorporation must be performed in the spring during seedbed preparation. If no spring work is anticipated, then both incorporations should be done in the fall.

For applications to winter wheat after seeding, apply and shallowly incorporate with a spike tooth, or spring tooth harrow set shallow as not to disturb wheat seed.

**SPRING—Before Seeding Incorporation:** For applications to fields of standing stubble, work fields once or twice with a disc, field cultivator or chisel plow, to provide soil in a good working condition. Apply granules and incorporate with equipment such as a culti-harrow or duck-foot cultivator. A second incorporation at right angles should provide best results.

**FOR SPRING AND DURUM WHEAT AND BARLEY IN MONTANA ONLY—**For suppression of Persian darnel (*Lolium temulentum*), apply this product before seeding and shallowly incorporate.

**After Seeding Incorporation:** Apply granules immediately after seeding and shallowly incorporate at right angles with equipment such as a flex multi-weeder or harrow. Adjust incorporation equipment to a depth so as not to disturb the seed.

**WHEN TO USE  
FALL TREATMENT INCORPORATED**

| <b>CROP</b>                        | <b>RATE (lbs/a)</b> | <b>WHEN TO APPLY</b>  |
|------------------------------------|---------------------|---|
| Spring and Durum Wheat, and Barley | 12.5 to 15          | *Within 3 weeks of normal freeze-up or until snow cover occurs. |
| Winter Wheat and Winter Barley**†  | 12.5 to 15          | Before or after seeding.  |

Apply lower rates on light soils and apply higher rates on heavy soils.

**BEST AVAILABLE COPY**

Incorporate within 48 hours. For methods, see "FIELD PREPARATION" and "INCORPORATED APPLICATIONS" sections.

- \* Average soil temperature at the 2 inch depth must be 40° F or less.
- \*\* For suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*) apply 15 pounds per acre prior to planting and shallow incorporate. In Colorado, Kansas, Nebraska, Oklahoma and South Dakota plant with hoedrills only.
- † For winter wheat in the Pacific Northwest, some crop thinning may occur, especially on clay knobs, where seed is "dusted in" or "floated on" and emergence is delayed. Thinning is usually more than offset by tillering and increased yields.

#### SPRING TREATMENT INCORPORATED

| CROP                      | RATE (lbs/a) | WHEN TO APPLY   |
|---------------------------|--------------|---|
| Spring and Durum<br>Wheat | 10 to 12.5   | After seeding before<br>wild oats germinate.                                |
| Spring and Durum<br>Wheat | 10 to 12.5*  | Before seeding and before<br>wild oats germinate.                           |
| Barley and<br>Triticale   | 12.5 to 15   | Just before or immediately<br>after seeding, before wild<br>oats germinate. |
| Lentils                   | 12.5 to 15   | Just before or immediately<br>after seeding, before wild<br>oats germinate. |
| Peas                      | 12.5 to 15   | Just before or immediately<br>after seeding, before wild<br>oats germinate. |

Apply lower rates on light soils and apply higher rates on heavy soils.

Incorporate within 48 hours. For methods, see "FIELD PREPARATION" and "INCORPORATED APPLICATIONS" sections.

- \* For press drills use the 10 pound rate. For HOEDRILLS, use the 12.5 pound rate.

Best spring and durum wheat results will be obtained with at least a 3-day delay between herbicide incorporation and seeding.

#### SURFACE APPLICATION WITH DELAYED INCORPORATION

7 of 10

**USE ONLY IN IDAHO, MINNESOTA, MONTANA,  
NORTH DAKOTA, AND UTAH**

This product may be applied as a surface application with a delayed incorporation prior to planting spring wheat, durum wheat, and barley at a rate of 15 pounds per acre for all soil types.

Surface applications of this product which will be followed by a delayed incorporation in the spring may be made beginning 3 weeks prior to soil freeze-up in the fall and ending before spring thaw. (Average soil temperature at the 2-inch depth must be 40° F or less).

Surface applications must be incorporated in the spring; two passes are recommended. If surface applications are made because soil conditions do not permit incorporation and these conditions change, making incorporation possible, incorporate even if several weeks after application.

This surface application may be made in standing stubble or to fields with surface residues. Surface applications should not be made to fields covered with snow or with excessive crop residue, which will not allow granule contact with the soil.

Surface applications with delayed incorporation may not provide the same level of wild oat control as fall incorporated applications. Wild oat control resulting from delayed incorporation may be reduced if drought or abnormally warm temperatures occur between application and crop emergence in the spring. To achieve the most reliable wild oat control, this product should be incorporated within 48 hours of application as described in the "FIELD PREPARATIONS" and "INCORPORATED APPLICATIONS" sections of this label.

**NO-TILL APPLICATION**

**FOR WILD OAT CONTROL IN SPRING PLANTED CROPS  
IN MONTANA ONLY**

This product may be applied prior to planting no-till spring wheat, durum wheat, and barley at a rate of 15 pounds per acre for all soil types.

No-till applications of this product may be made 3 weeks prior to soil freeze-up and ending before spring thaw. (Average soil temperature at the 2-inch depth must be 40° F or less).

No-till applications may be made in standing stubble or to fields with surface residue. No-till applications should not be made under conditions which will not allow granule contact with the soil.

No-till applications are to be made in reduced tillage/no-till systems only. If this product is used in any other system, it must be applied and incorporated as described elsewhere in this label.

**FOR FALL PLANTED CROPS**

10 4 10

FOR USE IN COLORADO, KANSAS, NEBRASKA, OKLAHOMA AND SOUTH DAKOTA - This product can be surface-applied prior to seeding of winter wheat without incorporation when seeding is done with a hoeddrill. Apply 15 pounds per acre for suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*).

FOR USE IN IDAHO, MONTANA, OREGON, UTAH and WASHINGTON - Apply this product at a rate of 15 pounds per acre as a surface application ahead of a Yielder™ type no-till drill to obtain suppression of wild oats and downy brome in winter wheat or barley. A flex-tine harrow attached to the drill or used shallowly following planting at a right angle to the drill rows may improve herbicide performance.

Yielder™ is a trademark of Yielder Drill, Inc.

843.53-000.45/53

If you have any questions concerning this label pamphlet or the proper application or use of Far-Go herbicide, contact your local Monsanto Sales Representative.

---

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

---

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.

**BEST AVAILABLE COPY**