

524-292

4/28/2003

1/11



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
524-292

Date of Issuance:
APR 28 2003

NOTICE OF PESTICIDE:
 Registration
 x Reregistration

Term of Issuance:
Unconditional

Name of Pesticide Product:
Granular Far-Go
Herbicide

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Monsanto Company
600 13th Street, Suite 660
Washington, DC 20005

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above with the provision that you make the following changes to your label before you release the product for shipment.

1. An "EPA Establishment Number" must be added to your container label.
2. ON page 10, delete "In case of emergency involving this product," as this information appears elsewhere on the label.
3. Under Directions for Use, the heading "Recommended Rate Per Acre" should be revised to "Application Rates per Acre", since the current statement may imply that the application rates are only recommendations and that they do not have be followed.

Enclosed is a copy of your label stamped "Acceptable With Comments". This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain registration of your product.

Please submit three (3) copies of your labeling bearing the above changes before you release the product for shipment.

Signature of Approving Official:

Date:

Vukob Kuznetsov for James A. Tompkins

4/28/03

2/11

MASTER LABEL
29006

GRANULAR



Herbicide by Monsanto

Granular selective herbicide for fall or spring application to control wild oats in winter wheat, spring and durum wheat, barley, triticale, peas (green, field dried, chickpeas, garbanzo beans), lentils, and sugarbeet; and for fall application to suppress *Bromus* species (*B. tectorum*, *B. secalinus* and *B. japonicus*) in winter wheat and in winter barley.
For use in Colorado, Idaho, Kansas, Minnesota, Montana, Nebraska, Nevada, North Dakota, Oregon, South Dakota, Utah, Washington, and Wyoming.

Complete Directions for Use

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Container Label Statements:

(For Bulk Bags):

THIS IS AN END-USE PRODUCT.. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A MONSANTO REPACKAGING OR TOLL REPACKAGING AGREEMENT.

(For Multiwall Bags):

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING.

ACTIVE INGREDIENT:

Triallate, S-(2,3,3-Trichloroallyl)-diisopropylthiocarbamate..... 10.0%

OTHER INGREDIENTS:..... 90.0%

..... 100%

3/11

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

Causes moderate eye irritation.
Avoid contact with skin, eyes or clothing.

Wash thoroughly with soap and water after handling.

Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ACCEPTED
with COMMENTS
in EPA Letter Dated:
APR 28 2003
524-292

FIRST AID

If in eyes

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses. If present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

- Have the product container or label with you when calling a poison control center or physician, or going for treatment.
- In case of an emergency involving this product, Call Collect, day or night, (314)694-4000.
- This product is identified as **Granular Far-Go Herbicide, EPA Reg. No. 524-292.**

Personal Protective Equipment (PPE)

Loaders, applicators*, flaggers*, and other handlers must wear: Long-sleeved shirt and long pants, shoes plus socks.

In addition, loaders must wear: A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter.

*See engineering controls below for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Engineering Controls

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for Agricultural Pesticides [40 CFR 170.240(d)(6)],

Flaggers must be in an enclosed cab that meets the definition in the Worker Protection Standard for Agricultural Pesticides for dermal protection and in addition to wearing the required PPE specified above, have immediately available for use in case they must leave the cab: coveralls, chemical-resistant gloves and chemical-resistant footwear, such as nitrile rubber, neoprene, neoprene rubber or polyethylene.

4/11

Environmental Hazards

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Ground Water Advisory

Triallate has a degradation product TCPSA (trichloropropene sulfonic acid) with properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Surface Water Advisory

Under some conditions, the triallate degradate TCPSA may have a high potential for runoff into surface water (primarily via dissolution in runoff water). These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Monsanto Supplemental Labeling.

Do not allow this product to drift. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete following application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls; Shoes plus socks; Chemical-resistant gloves, such as nitrile rubber, neoprene rubber or polyethylene. For more options, follow the instructions for category A (dry and water-based formulations) on an EPA chemical resistant category selection chart.

For Product Information or Assistance in Using This Product, Call TollFree 1-800-332-3111

5/11

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage and disposal.

PESTICIDE STORAGE: Keep bag closed to prevent spills and contamination.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed. Completely empty container into transfer or application equipment. Do not reuse container. Return emptied container per the Monsanto container return program. If not returned, dispose of empty sack in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Container Label Statements:

(For Bulk Containers):

Triple rise emptied bulk container. Then offer for recycling or reconditioning, or dispose of in a manner approved by state and local authorities.

(For Multiwall Bags):

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This herbicide is recommended for wild oat control in barley, peas (green, field dried, chickpeas, garbanzo beans), lentils, durum, spring and winter wheat, triticale, and sugarbeet only, and for suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*) in winter wheat and winter barley. Other crops should not be treated with this product because injury may occur. For barley, durum, spring wheat, triticale and sugarbeet, this product may be applied on the soil either in the fall or in the spring before wild oats germinate. For lentils and peas, this product may only be applied in the spring before wild oats germinate. For winter wheat, this product may only be applied in the fall before wild oats germinate.

Application equipment must be properly calibrated: application of too much herbicide may injure the crop; application of too little may result in poor wild oat control. Recommended rates should be followed in order to:

1. Avoid crop injury
2. Avoid crop residue at harvest
3. Control wild oats
4. Suppress *Bromus* species

General Precautions and Restrictions

Application to a field which is wet, lumpy, rough or ridged will result in reduced wild oat control and promote crop thinning. Incorporation must be completed within 48 hours after application and before germination of the wild oats. If weeds, including wild oats, have emerged prior to treating and/or planting, they must be controlled. All deep tillage by cultivators or double disc implements must be completed prior to application. Do not plow following application of this herbicide.

Seeding may be done either before or after application, depending upon the crop that is to be sown. If seeding is delayed, shallow reworking of the treated area before seeding will not destroy the effects of the chemical.

6/11

When using this herbicide, a strip should be left untreated for proof of results. Weed control may be evaluated also by removing a surface inch or two inches of the soil at the time of germination to inspect the number of wild oats that were killed before emergence.

Wild oats are usually killed before emergence, but occasionally, and particularly under dry conditions, plants may reach the 3 - 4-leaf stage before they die.

Under conditions of prolonged high temperature at the time of germination, or extreme drought in the spring, this product may not maintain the usual high standard of wild oat control.

ATTENTION

DELAYED EMERGENCE, STAND REDUCTION, STUNTING AND YIELD LOSS MAY RESULT DUE TO COLD OR WET CONDITIONS, IRRIGATION DURING GERMINATION OR EMERGENCE, DEEP PLANTING, SOIL CRUSTING, DISEASE, INSECTS, OR SELECTION OF A VARIETY GENETICALLY SUSCEPTIBLE TO STRESS. THESE ARE CONDITIONS BEYOND THE CONTROL OF MONSANTO, AND THE GROWER SHOULD CONSULT THE STATE UNIVERSITY EXTENSION SPECIALIST OR SEED PRODUCER OR SUPPLIER FOR LOCAL VARIETAL INFORMATION AND RECOMMENDED TILLAGE AND PLANTING PRACTICES AND DATES.

LATE PLANTING OF WINTER WHEAT OR PLANTING WINTER WHEAT UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS THAT RESULT IN FAILURE OF THE WHEAT TO EMERGE IN THE FALL MAY RESULT IN STAND REDUCTION, STUNTING AND YIELD LOSS.

Domestic oats should not be seeded where this product was used the previous year.

[Optional Voluntary Text]

Do not rotate to crops other than winter wheat, spring and durum wheat, triticale, barley, peas (green, field dried, chickpeas, garbanzo beans), lentils, and sugarbeets for 12 months after a Far-Go application.

Do not graze livestock on treated crops.

When this product is used on clay knobs, crop thinning may occur due to uneven seeding depth and poor crop establishment conditions. Thinning is usually offset by increased tillering and by the reduction of wild oat competition.

IN THE STATE OF MONTANA, do not use this product on fields to be seeded to hard red spring wheat with press drills, if the field is or will be irrigated in the current growing season.

Do not use postplant incorporation with hoe drills. This practice will result in unacceptable seeding depth and an excessive concentration of this product in the seed zone.

Use of this product not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

APPLICATION EQUIPMENT AND TECHNIQUES

This product must be applied through a specially designed ground applicator or airplane capable of applying small quantities of granules evenly.

Application is limited to one per growing season and must not exceed 15 pounds of Granular Far-Go herbicide per acre.

7/11

GROUND EQUIPMENT:

It is important that the applicator be calibrated properly to deliver the desired amount of this product to avoid applying too little, or too much, material. To give even distribution, scatter plates (similar to those used for applying granules in a band) must be attached to each delivery tube or outlet in such a manner to give overall coverage. To calibrate, attach a collector pan, or bag, over each spreader plate or delivery tube. Operate over normal terrain to be treated at 4 to 5 miles per hour. Collect the granules from all outputs after covering the desired distance.

For example: If a 14-foot applicator is being used, stake off a distance of 324 feet in the field to be treated (distance equal to 1/10 acre treated). After attaching a bag, or collection pan, to each outlet, collect the granules while driving the desired distance. Check to see that each outlet disperses the same amount of granules. Combine all samples and weigh. For the above set of conditions, the quantity of granules that should be collected for the following recommended rates of broadcast treatment are:

	RECOMMENDED RATE PER ACRE		
	10 lbs	12.5 lbs	15 lbs
Amount to be collected	1 lb	1.25 lbs	1.5 lbs

If more or less than the desired quantity is collected, adjust accordingly and again collect the granules while driving the staked-off distance. Continue this procedure until the required amount is delivered.

AIRPLANE: For aerial applications, attachments designed for applying low volumes of granules must be used. In order to ensure uniform aerial application, it is recommended that the field distribution pattern is checked and any necessary gate and vent modifications are made to ensure an even pattern distribution. In order to ensure uniform application and to avoid overlapping and possible crop injury, it is recommended that two flaggers, one at each end of the field, be used.

FIELD PREPARATION

Before applying this product, be sure the soil is in good working condition. All deep tillage by cultivation, or double disc implements, must be completed prior to application of this product. If stubble ground is being treated, one or two passes with a field cultivator may be required before application and incorporation.

INCORPORATED APPLICATIONS

When summer fallow or plowed ground is being treated, a field cultivator will provide adequate incorporation. For proper incorporation of this product, set incorporation implement to work the soil no deeper than 3 to 4 inches. Do not use disc implements for incorporation.

FALL-- On summer fallow ground which is loose and free of lumps and trash, incorporate this product using equipment similar to a culti-harrow or a duck-foot with rod weeder attachment.

For applications to fields or standing stubble, fields may be worked once or twice with a field cultivator or chisel plow. Apply granules and incorporate with a field cultivator or culti-harrow.

If soil must be ridged after incorporating to prevent soil erosion by high winds, the depth of ridging should be kept to a minimum.

In preparing the herbicide-treated area for seeding in the spring, care must be taken to avoid working the soil any deeper than fall tillage.

When incorporating with a single pass in the fall, a second incorporation must be performed in the spring during seedbed preparation. If no spring work is anticipated, then both incorporations should be done in the fall.

8/11

For applications to winter wheat after seeding, apply and shallowly incorporate with a spike tooth, or spring tooth harrow set shallow as not to disturb wheat seed.

SPRING-- Before Seeding Incorporation: For applications to fields of standing stubble, work fields once or twice with a disc, field cultivator or chisel plow, to provide soil in a good working condition. Apply granules and incorporate with equipment such as a culti-harrow or duck-foot cultivator. A second incorporation at right angles should provide best results. A delay of at least 3 days between the first and second incorporation is recommended for optimum performance.

FOR SPRING AND DURUM WHEAT AND BARLEY IN MONTANA ONLY-- For suppression of Persian darnel (*Lolium temulentum*), apply this product before seeding and shallowly incorporate.

After Seeding Incorporation: Apply granules immediately after seeding and shallowly incorporate at right angles with equipment such as a flex multi-weeder or harrow. Adjust incorporation equipment to a depth so as not to disturb the seed.

FALL TREATMENT INCORPORATED

CROP	RATE (lbs/A)	WHEN TO APPLY
Spring and Durum Wheat, and Barley	12.5 to 15	*Within 3 weeks of normal freeze-up or until snow cover occurs.
Winter Wheat and Winter Barley**†	12.5 to 15	Before or after seeding.
Sugarbeet	15	Fall application – Within three weeks of normal freeze-up or until snow cover occurs.

Apply lower rates on light soils and apply higher rates on heavy soils.

Incorporate within 48 hours. For methods, see "FIELD PREPARATION" and "INCORPORATED APPLICATIONS" sections of this label.

- * Breakdown of this product is minimal at soil temperatures of 40° F or less.
- ** For suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*) apply 15 pounds per acre prior to planting and shallowly incorporate. In Colorado, Kansas, Nebraska and South Dakota plant with hoeddrills only.
- † For winter wheat in the Pacific Northwest, some crop thinning may occur, especially on clay knobs, where seed is "dusted in" or "floated on" and emergence is delayed. Thinning is usually more than offset by tillering and increased yields.

9/11

SPRING TREATMENT INCORPORATED

CROP	RATE (lbs/A)	WHEN TO APPLY
Spring and Durum Wheat	10 to 12.5	After seeding before wild oats germinate.
Spring and Durum Wheat	10	Before seeding and before wild oats germinate- Disc/Press drills.
Spring and Durum Wheat	10 to 12.5	Before seeding and before wild oats germinate - Hoedrills.
Barley and Triticale	12.5 to 15	Just before or immediately after seeding, before wild oats germinate.
Lentils	12.5 to 15	Just before or immediately after seeding, before wild oats germinate.
Peas (green, field dried, chickpeas, garbanzo beans)	12.5 to 15	Just before or immediately after seeding, before wild oats germinate.
Sugarbeet	15	Before seeding and before wild oats germinate

Apply lower rates on light soils and apply higher rates on heavy soils.

Incorporate within 48 hours. For methods, see "FIELD PREPARATION" and "INCORPORATED APPLICATIONS" sections of this label.

Best spring and durum wheat results will be obtained with at least a 3-day delay between herbicide incorporation and seeding.

SURFACE APPLICATION WITH DELAYED INCORPORATION USE ONLY IN IDAHO, MINNESOTA, MONTANA, NORTH DAKOTA, AND UTAH

This product may be applied as a surface application with a delayed incorporation prior to planting spring wheat, durum wheat, and barley at a rate of 15 pounds per acre for all soil types.

Surface applications of this product which will be followed by a delayed incorporation in the spring may be made beginning 3 weeks prior to soil freeze-up in the fall and ending before spring thaw. (Average soil temperature at the 2-inch depth must be 40° F or less).

Surface applications must be incorporated in the spring; two passes are recommended. If surface applications are made because soil conditions do not permit incorporation and these conditions change, making incorporation possible, incorporate even if several weeks after application.

This surface application may be made in standing stubble or to fields with surface residues. Surface applications should not be made to fields covered with snow or with excessive crop residue, which will not allow granule contact with the soil.

10/11

Surface applications with delayed incorporation may not provide the same level of wild oat control as fall incorporated applications. Wild oat control resulting from delayed incorporation may be reduced if drought or abnormally warm temperatures occur between application and crop emergence in the spring. To achieve the most reliable wild oat control, this product should be incorporated within 48 hours of application as described in the "FIELD PREPARATIONS" and "INCORPORATED APPLICATIONS" sections of this label.

**NO-TILL APPLICATION
FOR WILD OAT CONTROL IN SPRING PLANTED CROPS**

This product may be applied prior to planting no-till spring wheat, durum wheat, and barley at a rate of 15 pounds per acre for all soil types.

For best results no-till applications of this product may be made 3 weeks prior to soil freeze-up and ending before spring thaw (average soil temperature at the 2-inch depth must be 40° F or less). Spring applications can also be made if applications are made at least 10 to 14 days prior to seeding.

No-till applications may be made in standing stubble or to fields with surface residue. No-till applications should not be made under conditions which will not allow granule contact with the soil.

No-till applications are to be made in reduced tillage/no-till systems only. Best results should be expected when utilizing drills equipped with high disturbance openers. When seeding with drills equipped with low disturbance openers, or with Spring applications, use of heavy harrows or similar minimal disturbance implements prior to seeding may improve control. It is recommended that seeding be delayed until fields begin to green up in the spring and an application of a Roundup Family Brand herbicide has been made to control emerged weeds. If this product is used in any other system, it must be applied and incorporated as described elsewhere in this label.

**SURFACE APPLICATION
FOR FALL PLANTED CROPS**

FOR USE IN COLORADO, KANSAS, NEBRASKA AND SOUTH DAKOTA-- This product can be surface-applied prior to seeding of winter wheat without incorporation when seeding is done with a hoed drill. Apply 15 pounds per acre for suppression of downy brome (*Bromus tectorum*), cheat (*Bromus secalinus*) and Japanese brome (*Bromus japonicus*).

FOR USE IN IDAHO, MONTANA, OREGON, UTAH and WASHINGTON -- Apply this product at a rate of 15 pounds per acre as a surface application ahead of a Yielder™ type no-till drill to obtain suppression of wild oats and downy brome in winter wheat or barley. A flex-tine harrow attached to the drill or used shallowly following planting at a right angle to the drill rows may improve herbicide performance.

If you have any questions concerning this label pamphlet or the proper application or use of Granular Far-Go herbicide, contact your local Monsanto Sales Representative.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. **NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE.** This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

11/11

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

In case of an emergency involving this product,

Far-Go and Roundup are registered trademarks of Monsanto Technology LLC.

Yielder is a registered trademark of Yielder Drill Company.

©2001 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167 U.S.A.
Formulated in the United States Using
Active Ingredient Made in Belgium.

EPA Reg. No. 524-292

29006