

**RESTRICTED USE PESTICIDE**  
due to oncogenicity. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**LASSO® II**

HERBICIDE BY MONSANTO

POWER  
GRANULES

Selective herbicide for preemergence weed control in corn, soybeans, peanuts and woody ornamentals.

The use of this product may be hazardous to your health. This product contains alachlor, which has been determined to cause tumors in laboratory animals.

In case of an emergency involving this product or for user safety information on this product, call collect, day or night (314) 694-4000.

**COMPLETE DIRECTIONS FOR USE**

EPA Reg. No. 524-296

®Lasso is a registered trademark of Monsanto Company

1994-1 1996-1

1202891 1/5300

**ACCEPTED**  
JUN 19 1996  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-296

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

**LIMIT OF WARRANTY AND LIABILITY**

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OF MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS  
AND DOMESTIC ANIMALS

Keep out of reach of children.

**WARNING! AVISO!**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

~~CAUSES EYE IRRITATION.~~

CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY

MAY CAUSE RESPIRATORY TRACT IRRITATION.

MAY CAUSE ALLERGIC SKIN REACTION.

Do not get in eyes, on skin or on clothing.

~~Avoid breathing dust.~~

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Call a physician immediately.

IF SWALLOWED, do NOT induce vomiting. Give large quantities of water. Get medical attention. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. Avoid alcohol.

IF ON SKIN, immediately flush with plenty of water while removing contaminated clothing. Sensitized persons should avoid further contact and reuse of contaminated clothing. As soon as soap is available, wash skin thoroughly with soap and water. Get medical attention.

IF INHALED, remove individual to fresh air. Get medical attention if breathing difficulty develops.

**Personal Protective Equipment (PPE)**

**Applicators and other handlers must wear:** long-sleeved shirt and long pants; ~~chemical resistant gloves, such as neoprene rubber; chemical resistant footwear~~ waterproof gloves; shoes plus socks; and protective eyewear.

~~Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.~~ Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should:

- Wash hands before eating drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

In case of emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

**ENVIRONMENTAL HAZARDS**

This chemical has been identified in limited ground water sampling and there is the possibility that it can leach through the soil to ground water, especially where soils are coarse and ground water is near the surface. Following application, and during rainfall events that cause runoff, this chemical may reach surface water bodies including streams, rivers and reservoirs.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when disposing of equipment washwaters.

ACTIVE INGREDIENT:

Alachlor: 2-chloro-2',6'-diethyl-N-(methoxymethyl)acetanilide.....	15%
INERT INGREDIENTS.....	85%
	100%

Contains petroleum distillates.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; chemical resistant gloves, such as neoprene rubber; chemical resistant footwear plus socks; waterproof gloves; shoes plus socks; and protective eyewear.

For more product information, call toll-free 1-800-332-3111

**Storage and Disposal**

Do not contaminate water, foodstuffs, seed or feed by storage or disposal.

**STORAGE:**

Below 32°F this product can cake. If product cakes, place in warm, dry area until product uncakes.

**DISPOSAL:**

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable federal, state or local procedures. Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned or destroyed.

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**GENERAL INFORMATION**

This product is recommended for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label. This product may be applied either as a surface application prior to or after planting or shallowly incorporated prior to planting to blend the herbicide treatment into the upper 1/2 to 2 inches of soil. Except for minimum or conservation tillage systems, the seedbed should be fine, firm and free of clods and trash.

Performance is improved when moisture occurs within one week after application. The amount required depends primarily upon the amount of soil moisture in the weed germination zone at the time of application, soil type and organic matter content, but 1/3 to 3/4 inch is generally sufficient for performance of this product. High intensity or excessive rainfall or excessive irrigation after application may reduce control. This product will not control emerged seedlings.

Use of this product not consistent with this label may result in injury to persons, animals or crops or other unintended consequences. Keep container closed to prevent spills and contamination.

**SOIL TEXTURE**

Applicators should evaluate soil and weed conditions carefully to assure that they choose the lowest effective label rate.

The recommended use rates of this product generally vary with soil texture. Unless soil texture is specifically named, rate tables throughout this label refer to only three soil textural groups: coarse, medium and fine. The following is a listing of the soil textures included in each of these three soil textural groups:

SOIL TEXTURAL GROUP	SOIL TEXTURE
COARSE:	sand, loamy sand, sandy loam
MEDIUM:	loam, silt loam, silt, sandy clay loam
FINE:	silty clay loam, clay loam, sandy clay, silty clay, clay

Refer to the above table to determine the corresponding soil textural group for the soil to be treated.

DO NOT BAND apply this product on COARSE textured soils.

#### HANDLING INSTRUCTIONS

##### NOTE

Direct contact or exposure to this product should be minimized. The following instructions for loading and repairing equipment should be followed in order to minimize this exposure. Review the protective clothing requirements as listed in the Precautionary Statements section of this label and do not use this product until you have the necessary protective clothing.

##### Equipment Cleaning and Repair

Cleaning and repair of application equipment is a source of exposure to this product. Care should be taken to minimize exposure during cleaning and repair of application equipment. Whenever possible, this equipment should be rinsed before being cleaned or repaired. When repairs must be made during application, the equipment should be shutdown, and special care should be taken to avoid contact with the pesticide.

#### APPLICATION INFORMATION APPLICATION SYSTEMS

**Ground Broadcast Treatment**--Apply the recommended rates of this product per acre. Use broadcast applicators that can apply small amounts of granules evenly.

**Band Treatment** (Behind Press Wheel only)--Use applicators designed for this purpose. Calculate the amount of granules per acre needed for band treatment as follows:

$$\begin{array}{rcl}
 \text{Band width} & & \text{Broadcast} \\
 \text{in inches} & \times & \text{Rate} \\
 \text{Row width} & & \text{per acre} \\
 \text{in inches} & & \text{per acre} \\
 & = & \text{Band Rate}
 \end{array}$$

For example: If 26 pounds per acre are recommended for broadcast treatment and you wish to treat a band of 14 inches wide in crop rows spaced 38 inches apart:

$$\begin{array}{rcl}
 14 & & 9.6 \text{ pounds of granules needed} \\
 38 \times 26 & = & \text{for the band treatment}
 \end{array}$$

**NOTE:** For application of this product do not rely on settings used for other granular herbicides.

**Calibration**--The following settings are only guidelines for calibration, they do not allow for equipment variation or wear. Check settings after initial calibration so that the proper rate is applied.

GRANULAR APPLICATOR SETTING

APPLICATOR	Broadcast Rate Per Acre			
	16 Pounds		20 Pounds	
	4 mph	5 mph	4 mph	5 mph
Allis Chalmers	2	2	3	3
Gandy (bolt-on)	12	13	13	15
(snap-on)	12	13	14	15
I.H. before 1981	1/9	2/2	2/2	2/6
1981 -- 1 gauge/hopper	1/9	2/1	2/1	2/4
2 gauges/hopper	2/0	2/2	2/2	2/6
John Deere --				
metal hopper	1/27	1/31	1/31	2/1
plastic hopper,				
-10, 20 etc. increment	13	15	15	19



-5, 15 etc. increment	11	13	13	15
Knob meter	18	21	21	27
Noble-hopper				
bottom hole ●	10	13	13	15
Noble and White-				
bottom hole ■	7	8	9	10
Buffalo				
Fluted-row feed	2.0	2.0	2.5	2.5
Plateless planter	8	10	10	13

GRANULAR APPLICATOR SETTING

Broadcast Rate Per Acre  
23 Pounds                      26 Pounds

APPLICATOR	4 mph	5 mph	4 mph	5 mph
Allis Chalmers	4	4	4	4
Gandy (bolt-on)	14	17	16	18
(snap-on)	15	16	16	17
I.H. before 1981	2/4	2/9	2/6	3/5
1981 -- 1 gauge/hopper	2/3	2/6	2/5	2/8
2 gauges/hopper	2/4	2/8	2/6	3/1
John Deere --				
metal hopper	1/33	2/4	2/2	2/6
plastic hopper,				
-10, 20 etc. increment	17	21	19	24
-5, 15 etc. increment	14	16	15	17
Knob meter	25	31	28	35
Noble-hopper				
bottom hole ●	14	17	16	19
Noble and White-				
bottom hole ■	10	12	11	13
Buffalo				
Fluted-row feed	3.0	3.0	3.5	3.5
Plateless planter	12	15	13	18

It is important that the applicator be calibrated properly to deliver the desired amount of this product. To calibrate, set the applicator and attach a bag over the spreader plate or remove the delivery tube and attach a plastic

nursing bottle over the spout. Operate at a constant speed on terrain to be planted. Four to 5 mph is suggested. For a 14-inch band, collect the granules for 500 linear feet (583 sq. ft. area) and measure the VOLUME or WEIGHT collected. Tap plastic nursing bottle or other measuring container gently to settle the granules when taking a reading.

Refer to the following table to determine the broadcast rate per acre for the amount collected. If necessary adjust the application equipment so that each spout dispenses the proper amount.

Broadcast Rate Per Acre	Volume Collected in 500 linear feet for a 14 inch band	Weight Collected in 500 linear feet for a 14 inch band
16 lb.	4.4 fl. oz.	3.4 oz.
20 lb.	5.6 fl. oz.	4.3 oz.
23 lb.	6.4 fl. oz.	4.9 oz.
26 lb.	7.2 fl. oz.	5.6 oz.

For other band widths calculate the calibration distance as follows:

$$\frac{14 \text{ inches}}{\text{Band width in inches}} \times 500 = \text{Calibration distance in linear feet.}$$

**APPLICATION TIMING AND METHODS**

**Preemergence Surface Application** - This product may be applied to the soil surface after planting and prior to either crop or weed emergence. Apply within 5 days of last preplant tillage. If weeds emerge after treatment, or if treatment is applied more than 5 days after last preplant tillage, rotary hoe or shallowly cultivate immediately to improve performance. Precipitation or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone. The amount of precipitation or overhead sprinkler irrigation required depends on existing soil moisture, soil type and percent organic matter content, but 1/3 to 3/4 inch is normally adequate. Performance is improved when moisture is received within 7 days after application and prior to weed emergence. High intensity or excessive rainfall or excessive irrigation after application may reduce control.

**CULTIVATION INFORMATION**

If grass or weed seedlings emerge before it rains, rotary hoeing will improve performance.

If cultivation is necessary because of soil crusting or compaction, tillage should be shallow to minimize dilution of this herbicide.

Except for the above, as long as the grasses and weeds are controlled, delay cultivation as long as possible.

Only after weeds emerge in the treated band, set cultivator to throw soil into the row covering the band. On peanuts, do not throw soil into the row.

WEEDS CONTROLLED

Applicators should evaluate soil and weed conditions carefully to assure that they choose the lowest effective label rate.

When applied as directed under the conditions described, this product CONTROLS the following weeds:

ANNUAL GRASSES

**Barnyardgrass**

*Echinochloa crus-galli*

**Crabgrass**

*Digitaria ischaemum*

*Digitaria sanguinalis*

**Foxtail, robust**

*Setaria viridis*

**Foxtail, giant**

*Setaria faberi*

**Foxtail, green**

*Setaria viridis*

**Foxtail, yellow**

*Setaria lutescens*

**Goosegrass**

*Eleusine indica*

**Panicum, browntop**

*Panicum fasciculatum*

**Panicum, fall**

*Panicum dichtomifloru*

**Signalgrass, broadleaf**

*Brachiaria spp.*

**Sprangletop, red**

*Leptochloa filiformis*

**Witchgrass**

*Panicum capillare*

**ANNUAL BROADLEAVES**

---

**Carpetweed**

*Mollugo verticillata*

**Galinsoga**

*Galinsoga spp.*

**Groundcherry, cutleaf<sup>1</sup>**

*Physalis angulata*

**Nightshade, black<sup>1</sup>**

*Solanum nigrum*

**Nightshade, hairy<sup>1</sup>**

*Solanum sarrachoides*

**Pigweed, Carelessweed**

*Amaranthus spp.*

**Purslane**

*Portulaca oleracea*

**Pursley, Florida**

*Richardia scabria*

**Waterhemp**

*Amaranthus tuberculatos*

<sup>1</sup>Use a minimum of 17 pounds of this product per acre on coarse soils, or 20 to 26 pounds of this product on medium and fine soils for control of this weed.

When applied immediately after planting, under the conditions described, this product at a rate of 17 to 26 pounds per acre on a broadcast basis REDUCES COMPETITION from the following hard-to-control weeds:

**ANNUAL GRASSES**

---

Cupgrass, woolly  
*Eriochloa villosa*

Johnsongrass, seedling  
*Sorghum halepense*

Panicum, Texas  
*Panicum texanum*

Sandbur, Grassbur  
*Cenchrus incertus*

Shattercane, wild cane  
*Sorghum bicolor*

**ANNUAL BROADLEAVES**

---

Beggarweed, Florida  
*Desmodium tortuosum*

Lambsquarters  
*Chenopodium album*

Morningglory, smallflower  
*Jacquemontia tamnifolia*

Ragweed, common  
*Ambrosia artemisiifolia*

Smartweed  
*Polygonum pensylvanicum*  
*Polygonum persicaria*

Sida, spiny; Teaweed  
*Sida spinosa*

Starbur, bristly  
*Acanthospermum hispidum*

**SEDGES**

---

**Nutsedge, yellow**  
*Cyperus esculentus*

**CORN**

(Field corn, Hybrid Seed Corn, Silage Corn, Sweet Corn)

When making applications on corn, do not exceed a total of 26 pounds of this product per acre per year.

**LASSO® II**

Apply this product for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

**APPROVED APPLICATION SYSTEMS**

Ground - Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

**APPROVED APPLICATION METHOD**

**Preemergence Surface** - Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation. See the following tables for recommended broadcast and banded rates per acre for preemergence surface applications.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

Applicators should evaluate soil and weed conditions carefully to assure that they choose the lowest effective label rate.

For control of listed annual grasses and broadleaf weeds apply after planting and before crop or weeds emerge. For band treatment apply granules behind the press wheel. See table below for recommended rates on various soil textures.

**Preemergence Surface Application Rates**

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse***	13 to 15	13 to 15
Medium	13 to 18	13 to 18
Fine	13 to 18	17 to 22

\*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate in the recommended range in areas of heavy weed infestation. Use a minimum of 17 pounds per acre on coarse soils and 20 to 26 pounds on medium and fine soils to control black or hairy nightshade.

\*\*\*When applied under sprinkler irrigation on coarse soils, use a minimum of 20 pounds per acre of this product.

**For Lister or Deep Furrow Planted Corn**

**Broadcast Treatment**--Apply at the rate of 16 pounds per acre.

**Band Treatment (Behind Press Wheel)**--For a 10 inch band treatment over the row (40 inch spacing) apply 4 pounds per acre. For a 7 inch band treatment over the row (40 inch spacing), apply 2.8 pounds per acre.

If heavy rains follow application, weed control on the top and sides of the furrow may not be as good as that obtained in the bottom of the furrow.

Where deep lister planting is practiced, only partial weed control can be expected on steep furrow sides. It is suggested that a 7 inch band be applied at the bottom of the furrow. Better results can be expected where the granules lie on an even flat surface which is not subject to shifting by wind or rain.

**For Furrow Irrigated Corn**

This product must be incorporated into the soil by harrow, finger weeder or rotary hoe. Shallow incorporation of at least 1/2 inch but not more than 2 inches may result in unsatisfactory weed control. Incorporation deeper than 2 inches may result in unsatisfactory weed control. Incorporation depth must be kept above the seed. Soil moisture at application must be sufficient for crop and weed emergence. Irrigation within 10 days following planting may improve weed control. See the table above for recommended rates.

**SOYBEANS**

When making applications on soybeans, do not exceed a total of 20 pounds of this product per acre per year.

**LASSO® II**

Do not feed forage, hay or straw from soybeans treated with this product. Do not ensile soybeans treated with this product.

Apply this product for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

**APPROVED APPLICATION SYSTEMS**

Ground—Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

**APPROVED APPLICATION METHODS**

**Preemergence Surface**—Apply this product after planting before crop and weed emergence and within 5 days after last preplant tillage operation.

**NOTE:** The highest levels of weed control will be obtained when applications are made prior to the emergence of the crop and weeds.

Weeds which are emerged at time of application are not controlled by this product. If weeds are emerged at application shallowly cultivate or rotary hoe immediately to improve performance.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and methods selected.



**RECOMMENDED RATES:** Refer to the following tables for the recommended broadcast treatment rates for this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

**Preemergence Surface Application Rates**

SOIL TEXTURAL GROUP*	<u>BROADCAST RATE PER ACRE (POUNDS)**</u>	
	Less than 3% organic matter	3% or more organic matter
Coarse***	13 to 15	13 to 15
Medium	13 to 18	13 to 18
Fine	13 to 18	17 to 20

\*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate in the recommended range in areas of heavy weed infestation. Use a minimum of 17 pounds per acre on coarse soils and 20 pounds on medium and fine soils to control black or hairy nightshade.

\*\*\*When applied under sprinkler irrigation on coarse soils, use 20 pounds per acre of this product.

**For Furrow Irrigated Soybeans**

This product must be incorporated into the soil by harrow, finger weeder or rotary hoe. Shallow incorporation of at least 1/2 inch but not more than 2 inches may result in unsatisfactory weed control. Incorporation deeper than 2 inches may result in unsatisfactory weed control. Incorporation depth must be kept above the seed. Soil moisture at application must be sufficient for crop and weed emergence. Irrigation within 10 days following planting may improve weed control. See the table above for recommended rates.

**PEANUTS**

**Broadcast**--Apply 20 pounds of this product per acre, from immediately after planting and until either crop or weeds emerge. NOTE: Hard-to-control weeds require 23 to 26 pounds per acre.

NOTE: Hard-to-control weeds require 23 to 26 pounds per acre.

NOTE: Do not feed forage or hay from treated crop.

**WOODY ORNAMENTALS**

**Field and Liner Grown Ornamentals**

Use in **Cotoneaster** (Cotoneaster spp.), **Crabapple** (Malus spp.), **Dogwood** (Cornus spp.), **Euonymus** (Euonymus spp. shrub spp. only), **Holly** (Ilex spp.), **Juniper** (Juniperus spp.) and **Yew** (Taxus spp.).

**Container Grown Ornamentals**

Use in **Holly** (Ilex spp.) and **Juniper** (Juniperus spp.).

DO NOT USE IN GREENHOUSES OR OTHER ENCLOSED STRUCTURES.

NOT RECOMMENDED FOR DOMESTIC APPLICATION, EXCEPT BY CERTIFIED APPLICATORS.

When applied as directed under the conditions described, this product controls the weeds listed on this label in newly planted or established field and container grown woody ornamentals.

DO NOT USE THIS PRODUCT ON SEEDBEDS, UNROOTED CUTTINGS OR PRIOR TO TRANSPLANTING. DO NOT APPLY THIS PRODUCT TO PLANTS WHOSE FOLIAGE IS MOIST OR WET AS GRANULES MAY STICK TO THIS FOLIAGE AND CROP INJURY MAY RESULT. OVERHEAD IRRIGATION APPLIED WITHIN ONE HOUR AFTER APPLICATION WILL WASH GRANULES OFF THE FOLIAGE AND INJURY WILL BE MINIMIZED.

**Broadcast Treatment**--Apply this product at a broadcast rate of 27 pounds per acre anytime after transplanting and before weeds emerge or after existing weeds have been removed. See the "WEED CONTROL" and "CULTIVATION INFORMATION" sections of this label for further information. Repeat applications of this product may be required for sustained long-term weed control. Do not exceed 2 applications or a total of 40 pounds of this product per year.

**NOTE:** Plant injury symptoms (leaf tip burn, discoloration, distortion, etc.) may occur with applications of this product to actively growing woody ornamentals. Do not retreat within twenty-one (21) days.

---

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

19/19

---

©MONSANTO COMPANY 1994

1202861-1/53EG

MONSANTO COMPANY ° AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.