

KEEP PEOPLE AND PETS OFF TREATED AREAS UNTIL SPRAY SOLUTION HAS DRIED

**DOMESTIC ANIMALS:** This product is considered to be relatively nontoxic to dogs and other domestic animals. However, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.). If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

#### ADDITIONAL PRECAUTIONS

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Roundup is a nonselective herbicide that is diluted and applied to the foliage of actively growing weeds as a spot or broadcast application. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms may require a week or more to appear, with burn-down usually occurring in 2 to 4 weeks. Symptoms are a gradual wilting and yellowing of the sprayed plant followed by deterioration of both shoots and roots. This product has no herbicidal activity in the soil and will not wash or leach to affect nearby vegetation.



This product controls many weeds around buildings, vacant lots, storage and recreational areas, gravel or ground bark mulch, shade trees, ornamentals and shrubs; along fences, sidewalks, driveways and patios; and prior to planting turfgrasses, shrubs, flower beds or vegetables listed on the label. When used as a spot treatment in lawns, all vegetation contacted will be damaged.

**THIS PRODUCT CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. AVOID SPRAY DRIFT CONTACT WITH DESIRABLE LAWN GRASSES, FLOWERS, VEGETABLES, SHRUBS OR TREES. DO NOT CONTACT GREEN BARK OF TREES OR SHRUBS IF DESIRABLE VEGETATION IS CONTACTED, WASH IMMEDIATELY WITH WATER.**

For more product information, call toll-free 1-800-225-2883

#### Storage and Disposal

Do not contaminate water, food, feed or seed by storage or disposal.

Store this product in its original container and in a secured storage area.

In case of SPILL or LEAK, soak up with paper towels and discard in trash.

To dispose of unused product, securely wrap original container in several layers of newspaper and discard in trash.

Empty container retains vapor and product residue. Observe all labeled safeguards until container is discarded.

Do not reuse container. Rinse thoroughly before discarding in trash.

#### WEEDS CONTROLLED

When applied as recommended under the conditions described, Roundup controls many common weeds including the following:

##### Annuals

Barnyardgrass	Lettuce, prickly	Ryegrass, annual
Bluegrass, annual	Mustard, fancy	Sandbur, field
Chickweed	Pennycress, field	Shepherdspurse
Crabgrass	Pigweed, redroot, smooth	Sowthistle
Foxtail		
Lambsquarter, common	Ragweed, common	

##### Perennials

Bahiagrass	Dallisgrass	Orchardgrass
Bermudagrass	Dandelion	Quackgrass
Bindweed, field	Fescue species	Ryegrass, perennial
Bluegrass, Kentucky	Fescue, tall	Thistle, Canada
Bromegrass, smooth	Johnsongrass	Timothy
Clover, red, white	Nutsedge, purple, yellow	Vaseygrass

##### Brush

Alder	Honeysuckle	Raspberry
Blackberry	Kudzu	Rose, multiflora
Cherry	Oak	Sumac
Dewberry	Poison Ivy	Trumpet creeper
Elderberry	Poison Oak	Willow

#### MIXING AND SPRAYING

Do not apply this product through any type of irrigation system.

**Hand-held Sprayers** - Add 1-1/2 to 2-1/2 fluid ounces (3 to 5 tbsp.) of this product to 1 gallon of clean water. Use the higher recommended rate for control of listed perennials and brush. Use pump-up-type or hand-trigger-type sprayers. USE OF HOSE-END SPRAYERS OR SPRINKLER-TYPE DEVICES MAY RESULT IN POOR AND/OR ERRATIC RESULTS.

Ensure uniform and complete spray coverage, but do not spray plants to the point that the spray solution runs off. Use a coarse spray. Do not spray during windy conditions, as spray drift may cause damage to any vegetation contacted.

#### WEEDS TO APPLY

Apply to vigorously growing plants. Apply to annual weeds anytime after emergence. For best results, apply to perennial weeds after seedheads, flowers or berries appear. On woody brush or trees, apply after leaves have fully expanded. Applications made at less than the recommended stage of growth may result in reduced control. After mowing, clipping or tilling, allow weeds to regrow to the recommended growth stage before spraying this product.

mowing, clipping, tilling, planting or sodding of treated areas for at least 7 days after application. This allows time for this product to move within the plant for most effective results. **SINCE THIS PRODUCT DOES NOT WORK IN SOIL, RETREATMENTS WILL BE NECESSARY TO CONTROL NEW WEEDS EMERGING FROM SEED.**

One application will control most weeds listed. Symptoms on certain tough perennial weeds and woody brush and tree species may be slow to develop. If control is not achieved within 4 weeks, allow time for the sprayed weeds to regrow and make a repeat application. In some cases, control of woody brush and trees treated in the fall may not be fully apparent until the following season.

#### LAWN RENOVATION

Best results are obtained when at least one mowing is skipped prior to treatment. This provides more leaf surface to intercept the spray. For areas where renovation and reseeding is desired, thorough spray coverage of all existing vegetation is necessary. Delay planting or sodding of treated areas for at least 7 days after application.

#### ORNAMENTALS

Any ornamental species may be planted in treated areas 7 days or more after application. Apply this product as recommended to control undesirable vegetation growing around ornamentals, trees or shrubs. Avoid spray contact with any desirable plants. To avoid injury on desirable plants, prevent spray or drift contact with foliage, stems or green bark. Do not spray into, or over the top of, growing ornamentals.

#### TREES (Apple, Citrus, Nut, Palm) AND GRAPES

Apply this product to control undesirable plants growing around these trees and vines. Spray contact, other than with mature bark of the main trunk, can injure the tree or vine. Allow 21 days between application and harvest.

▲ Cut or tear along perforation ▲

File # Form 13 117 80 432288  
AMERICAN AG. MADE IN U.S.A.



**BERRIES, SMALL FRUIT & VEGETABLE GARDENS**

The following vegetables may be planted in treated areas 7 days or more after application: asparagus, beans, beet greens, beets, blackberry, blueberry, boysenberry, broccoli, cabbage, carrot, cauliflower, celery, chicory, corn, cranberry, cucumber, currant, dewberry, elderberry, eggplant, garlic, gourds, gooseberry, horseradish, huckleberry, Jerusalem artichoke, kale, lentils, lettuce, loganberry, melons, mustard greens, okra, ollalieberry, onion, parsnips, peas, peppers, Irish and sweet potatoes, pumpkin, radish, raspberry (black, red), rutabagas, spinach, squash (summer, winter), strawberry, tomatillo, tomato\*, watercress, watermelon and turnips. In the treated area, do not plant any vegetables not listed on this label for one year following application. Do not apply this product in or around growing vegetables.

\*Use on tomatoes is restricted to direct seeded tomatoes only

**CLEANING EQUIPMENT**

Triple-rinse spray tank and flush all sprayer components with water to remove residues of this product. After thorough cleaning, equipment may be used to apply other products.



This page is intentionally left blank.

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

▲ Cut or tear along perforation ▲

Printed in the U.S.A. by  
AMERICAN LABEL MANUFACTURING CO., INC.

BEST AVAILABLE COPY

# Roundup Super Concentrate

Nonselective herbicide for lawn and garden use. Avoid spray on desirable plants as severe injury or destruction may result.

Read "NOTICE" in next paragraph before buying or using. If terms are not acceptable, return at once unopened.

**NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS IN THIS LABELING.**

REFORMULATION OR REPACKAGING IS PROHIBITED.

#### ACTIVE INGREDIENT:

Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.	41.0%
INERT INGREDIENTS:	59.0%
	100.0%

Contains 480 grams per litre or 4 pounds per U.S. gallon of glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt.

This product is protected by U.S. Patents No. 3,799,758, No. 4,405,584 and No. 4,840,650. Other patents pending. No license granted under any non-U.S. patent(s).

© MONSANTO COMPANY 1990

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.



Carefully read and follow detailed instructions in pamphlet.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

**WARNING! CAUSES EYE IRRITATION. HARMFUL IF SWALLOWED OR INHALED. MAY CAUSE SKIN IRRITATION.** Do not get in eyes, on skin or on clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling.

**FIRST AID: IF IN EYES,** immediately flush with plenty of water for at least 15 minutes. Get medical attention.

**IF ON SKIN,** immediately flush with plenty of water. Remove contaminated clothing. Wash clothing before reuse.

**IF SWALLOWED,** this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

**IF INHALED,** remove individual to fresh air. Get medical attention if breathing difficulty develops.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

#### Environmental Hazards

Do not apply directly to water or wetland (swamps, bogs, marshes or potholes). Do not contaminate water when disposing of equipment washwaters.

#### Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers. Do NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. (See additional Precautions inside Pamphlet.)

#### Storage and Disposal

Do not contaminate water, food, feed or seed by storage or disposal.

**STORAGE:** Store this product in its original container and in a secured storage area.

In case of SPILL or LEAK, soak up with paper towels and discard in trash.

**DISPOSAL:** To dispose of unused product, securely wrap original container in several layers of newspaper and discard in trash.

Empty container retains vapor and product residue. Observe all labeled safeguards until container is discarded.

Do not reuse container. Rinse thoroughly before discarding in trash.

This product is protected by U.S. Patents No. 3,799,758, No. 4,405,531 and No. 4,840,650. Other patents pending. No license granted under any non-U.S. patent(s).

© Trademark of Monsanto Company

© MONSANTO COMPANY 1990

MONSANTO COMPANY • AGRICULTURAL PRODUCTS • ST. LOUIS, MISSOURI 63167 U.S.A.  
EPA Est. 524 LA 1      EPA Reg. No. 524-308 AA      897 17 000 191 3

File # 70000 U.S. Pat. 3,799,758 and 4,405,531 and 4,840,650

▲ Cut or tear along perforation