

Keep out of reach of children.
See back panel for telephone number.

Keep out of reach of children.

Read "LIMIT OF WARRANTY AND LIABILITY" before
buying or using. If terms are not acceptable, return
at once unopened.

Read precautions on back panel.

ACTIVE INGREDIENT:	
"Glyphosate	85.0%
INERT INGREDIENTS:	15.0%
	100.0%

*N,N-bis(phosphonomethyl)glycine

U.S. Pat. No. 3,550,762 LIA Reg. No. 521-311 AA

NET 25 LBS., NET 11.34 KG.

ACCEPTED
Oct 26 1974
 U.S. DEPT. OF AGRICULTURE
 PHOSPHATE DIVISION
 WASHINGTON, D.C. 20250
 01311

MINIATURE FACSIMILE
 Smallest Type also available



PLANT GROWTH
PROMOTOR BY

Monsanto

Use as a wettable powder for foliar application to hasten ripening and increase the level of sucrose in sugar cane.

PRELIMINARY STATEMENTS

CAUTION! ...

Preparation of Concentrate

... Do not mix with other products ...

Storage and Disposal

... Avoid contact with skin, eyes, nose, and mouth ...

GENERAL INFORMATION


This product is a foliar applied plant growth regulator to hasten ripening and increase the level of sucrose in sugarcane ...

... Within 2 to 3 weeks after application, this product may produce a slight yellowing ...

... Do not contaminate water by leakage of equipment ...



224-311



NOTE

This product may not increase the sucrose content of sugarcane under conditions of good natural ripening ...

Application of Polaris may initiate development of shooting eyes.

AERIAL APPLICATION

AVOID DRIFT. Do not apply when winds are gusty or under other conditions which will allow drift ...

INSURE UNIFORM APPLICATION. To avoid streaked, uneven or overlapped application, use 2 flagmen, 1 at each end of the field ...

ADDITION OF DRIFT CONTROL ADDITIVES When the addition of a drift control additive is desired, Nalco-trol™ may be added at 8 to 16 ounces per 100 gallons of spray solution ...

Read and carefully observe the label on any state label and all other information appearing on the container ...

MIXING INSTRUCTIONS

To achieve the best spray mix, proceed as follows:

- 1. Place a wetting basket over the filling port and make all additions through it.
2. Fill the sprayer tank one-half full with water.
3. Add the recommended amount of this product.
4. If Nalco-trol is used, add the measured amount as a pencil thin stream to the agitating tank.
5. Wash through the wetting basket and fill the remaining portion of the tank with water.

DIRECTIONS FOR USE

Apply in 5 to 10 gallons of water per acre. Do not use on very light sandy soils. Equipment should be thoroughly flushed with water after each use to avoid possible corrosion.

NOTE — Use the higher recommended rate under adverse ripening conditions or on less responsive varieties.

Florida — Apply 2.5 to 3.5 pounds of this product per acre 5 to 8 weeks before harvest.

Louisiana — Apply 2.5 to 3.5 pounds of this product per acre 4 to 8 weeks before harvest.

Hawaii — Apply 4.7 pounds of this product per acre 6 to 10 weeks before harvest.

Texas — Apply 3.5 to 4 pounds of this product per acre 4 to 6 weeks before harvest.


EPA Est. 34661 1634-53 7 23 70

LIMIT OF WARRANTY AND LIABILITY

This company warrants that this material conforms to the characteristics shown on the label and is reasonably fit for the uses referred to in the directions for use ...

CRM — B. n. o. s. — KEEP DRY

In case of an emergency involving this product, call Chemagro Corp., Dept. 1000, 654 1000



MONSANTO COMPANY, Agricultural Promotions, St. Louis, MO, U.S.A.