



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

524-314

Date of Issuance:

NOV 19 2003

Term of Issuance:

Unconditional

Name of Pesticide Product:

Lasso Herbicide

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Monsanto Company
600 13th Street, N.W. Suite 660
Washington, DC 20005

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above, with following provisions.

1. Under "Storage and Disposal" revise "Storage" to read "Pesticide Storage", and revise "Disposal" to read "Pesticide Disposal".
2. Incorporate the following bulk container language into your Container Disposal section. Alternatively you may incorporate the language recently approved for EPA Registration No. 524-537 or 524-539.
3. Container Disposal (Bulk and Minibulk)

When the container is empty, replace the cap and seal all openings that have been opened during use, and return the container to the point of purchase, or to an alternate location designated by the registrant at the time of purchase of this product. If not returned to the point of purchase or to a designated location, triple rinse or pressure rinse the empty container and offer for recycling if available.

Instructions for Users and Refillers

The container must only be refilled with this pesticide product. **Do not Reuse the Container for Any other Purpose.** Do not transport if this container is damaged or leaking if the container is damaged, leaking, or obsolete, or to obtain information about recycling refillable containers,

Signature of Approving Official:

Date:

11-19-03

contact **(insert company name)** at **(insert phone number)**. Cleaning is not necessary prior to refilling with the same product. Clean container before final disposal. Disposal of this container must be in compliance with state and local regulations.

Instructions for Refillers

Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. If the container can not be refilled, triple rinse or pressure rinse the empty container and offer for recycling if available.

3. Refer to the attachment entitled Spray Drift Management for statements required for all products that permit aerial application.

Enclosed is a copy of your label stamped "Acceptable With Comments". This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain registration of your product



5/68

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Attachment-Spray Drift Management

Under the heading **Spray Drift Management** the text should read as follows:

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $3/4$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

Controlling Droplet Size

Volume-Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure-Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

4/68

Number of nozzles-Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation-Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length-For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application-Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

5/68

Temperature Inversions

Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun set and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

6/68

RESTRICTED USE PESTICIDE
due to groundwater concerns. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

LASSO®
Herbicide by Monsanto


Emulsifiable herbicide for weed control in corn, soybeans, peanuts, grain sorghum (milo) and crops listed.

The use of this product may be hazardous to your health. This product contains alachlor, which has been determined to cause tumors in laboratory animals.

In case of an emergency involving this product or for user safety information on this product, Call Collect, day or night, (314) 694-4000.

Complete Directions for Use

EPA Reg. No. 524-314
For use of this product in California see page 56.
Not for use in Nassau and Suffolk counties of New York.



Read each of these sections of this label for essential product performance information.

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OF MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING

7/68

FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

FLUID FERTILIZER COMPATIBILITY TEST

GENERAL INFORMATION

INGREDIENTS

LIMIT OF WARRANTY & LIABILITY

MIXING, SPRAYING & HANDLING INSTRUCTIONS

PRECAUTIONARY STATEMENTS

SOIL TEXTURE

STORAGE & DISPOSAL

WEEDS CONTROLLED WITH LASSO

WOODY ORNAMENTALS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep out of reach of children.

DANGER! PELIGRO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

CORROSIVE. CAUSES IRREVERSIBLE EYE DAMAGE.

CAUSES SKIN IRRITATION. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. This product may cause skin sensitization reactions in some people.

HARMFUL IF SWALLOWED.

HARMFUL IF INHALED. Avoid breathing vapor or spray mist.

Do not get in eyes, on skin or on clothing.

ACCEPTED
WITH COMMENTS
In EPA Letter Dated
NOV 19 2003

S24-314

8/68

FIRST AID
<p>If in eyes</p> <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
<p>If on skin or clothing</p> <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.• Sensitized persons should avoid further contact and reuse of contaminated clothing.
<p>If swallowed</p> <ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
<p>If inhaled</p> <ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
<ul style="list-style-type: none">• Have the product container or label with you when calling a poison control center or physician, or going for treatment.• In case of an emergency involving this product, Call Collect, day or night (314) 694-4000.• This product is identified as Lasso herbicide, EPA Registration No. 524-314.
<p>NOTE TO PHYSICIAN</p> <ul style="list-style-type: none">• This product may post an aspiration pneumonia hazard. Contains petroleum distillate.• Probable mucosal damage may contraindicate the use of gastric lavage.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are barrier laminate or viton. If you want more options, follow the instruction for category H on an EPA chemical-resistance category selection chart.

Mixers and loaders using engineering controls and applicators must wear: long-sleeved shirt and long pants; shoes and socks; and goggles or face shield.

9/68

256 **In addition to the above, mixers and loaders and other persons must wear** chemical-resistant gloves and
257 chemical-resistant apron when cleaning equipment.

258
259 **Mixers and loaders not using engineering controls and all other handlers exposed to the concentrate**
260 **must wear:** coveralls over short sleeved shirt and short pants, socks and chemical resistant footwear, goggles or
261 face shield, chemical-resistant gloves, and chemical resistant apron.

262
263 Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this
264 product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no
265 such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

266
267 Engineering Control Statement: Mixers and loaders supporting aerial applications, chemigation, or impregnation of
268 dry bulk fertilizer are required to use a mechanical transfer system designed by the manufacturer to transfer liquid
269 pesticide in a manner that prevents the liquid (but not necessarily any vapor) from contacting handlers and other
270 people during the transfer. The closed system must be used in a manner that meets the requirements listed in the
271 Worker Protection Standard (WPS for agricultural pesticides [40 CFR 170.240 (d) (4)]).

272
273 **IMPORTANT:** Mixers and loaders using closed systems must: (1) wear PPE specified above for mixers and
274 loaders using engineering controls, and (2) must be provided and have immediately available for use in an
275 emergency, such as a spill or equipment breakdown, the following PPE: Coveralls, chemical resistant footwear.

276
277
278
279 **User Safety Recommendations**

280
281 User should:

- 282 • Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- 283 • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- 284 • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As
285 soon as possible, wash thoroughly and change into clean clothing.

286
287 In case of an emergency involving this product,
288 Call Collect, day or night, (314) 694-4000.

289
290 **Environmental Hazards**

291
292
293
294
295 This chemical is toxic to terrestrial and aquatic plants, fish, and aquatic invertebrates. Do not apply directly to
296 water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff
297 may be hazardous to aquatic organisms in neighboring areas.

298
299 Alachlor can contaminate surface water through spray drift. Under some conditions, alachlor may also have a
300 high potential for runoff into surface water (primarily via dissolution in runoff water), for several weeks post-
301 application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters,
302 frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches
303 that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and
304 areas over-laying tile drainage systems that drain to surface water. Do not apply to water or to areas where
305 surface water is present, or to intertidal areas below the mean high water mark.

306
307 Care must be taken when using this product to prevent back siphoning into wells, spills or improper disposal of
308 excess pesticide, spray mixtures or rinsates.

309
310 Check valves or anti-siphoning devices must be used on all mixing and/or irrigation equipment. Disposal of excess
311 pesticide, spray mixtures or rinsate should be according to label use instructions or according to the state pesticide
312 or environmental control agency or the hazardous waste representative at the nearest EPA regional office.

10/68

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical and/or its metabolites are known to leach through soil into ground water under certain conditions as a result of registered uses. Use of this chemical in areas where soils are permeable, particularly where the water table is too shallow, may result in ground water contamination.

Do not apply to highly permeable soils (as classified by the USDA Natural Resources Conservation Service) where the depth to ground water is 30 feet or less.

Physical or Chemical Hazards

COMBUSTIBLE.

Do not use or store near heat or flame.

Use only with adequate ventilation.

In case of: FIRE, use water spray, foam, dry chemical or CO₂.

SPILL or LEAK, soak up, remove and keep away from humans and domestic animals.

11/68

ACTIVE INGREDIENT:

*Alachlor, 2-chloro-2',6'-diethyl-N-(methoxymethyl)acetaniide	45.1%
INERT INGREDIENTS	54.9%
	100.0%

Contains petroleum distillates.

*Contains 480 grams/litre or 4 pounds/gallon of 2-chloro-2',6'-diethyl-N-(methoxymethyl)acetaniide.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirement

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted- entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical-resistant gloves, made of any waterproof material, and shoes plus socks.

This product may not be mixed or loaded within 50 feet of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be mixed/loaded or used within 50 feet of all wells, including abandoned wells (unless the well has been properly capped or plugged), drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash-water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain a minimum of 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size capacity to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

Rotation to crops not specified on this label is prohibited.

12/68

For more information, call toll-free 1(800) 332-3111.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

STORE ABOVE 32°F TO KEEP PRODUCT IN SOLUTION.

Below 32°F crystals may form and settle to the bottom. If crystals form, place in a warm room and roll and/or shake the container frequently to redissolve before using. For bulk containers, see the container label for alternate storage information.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

DISPOSAL:

Wastes of this pesticide may cause eye burns and skin irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL OPTIONS:

(FOR REFILLABLE PORTABLE CONTAINERS)

Do not reuse this container except for refill in accordance with a valid Monsanto Repackaging or Toll Repackaging Agreement. If not refilled or returned to the authorized repackaging facility, triple rinse containers, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(FOR BULK CONTAINERS)

Triple rinse emptied bulk container. Then offer for recycling or reconditioning, or dispose of in a manner approved by state and local authorities.

(FOR PLASTIC 1-WAY CONTAINERS & BOTTLES)

Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(FOR DRUMS)

Do not reuse container. Return container per the Monsanto container return program. If not returned, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Lasso herbicide is recommended for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label. This product may be applied either as a surface application before or after planting or shallowly incorporated prior to planting to blend the herbicide treatment into the upper 1 to 2 inches of soil. The seedbed should be fine, firm and free of clods when tillage is performed.

Read and carefully observe precautionary statements and all other information appearing on the labeling of all products used in mixtures and sequential treatments.

13/68

Do not flood irrigate following application of this product or tank mixtures. This product will not control emerged seedlings.

Do not apply when conditions favor drift.

Applications which are not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

Flush sprayer with clean water after use.

SOIL TEXTURE

Applicators should evaluate soil and weed conditions carefully to ensure that they choose the lowest effective label rate.

The recommended use rates of this product and the other herbicides labeled for use in tank mixtures with this product generally vary with soil texture. Unless soil texture is specifically named, rate tables throughout this label refer to only three soil textural groups: coarse, medium and fine. The following is a complete listing of soil textures included in each of these three soil textural groups:

SOIL TEXTURAL GROUP	SOIL TEXTURE
COARSE	sand, loamy sand, sandy loam
MEDIUM	loam, silt loam, silt, sandy clay loam
FINE	silty clay loam, clay loam, sandy clay, silty clay, clay

Refer to the above table to determine the corresponding soil textural group for the soil to be treated.

MIXING, SPRAYING AND HANDLING INSTRUCTIONS

NOTE

The use of a mechanical transfer system is required by all mixer/loaders and/or applicators who treat 300 acres or more annually with pesticides containing alachlor as an active ingredient.

Direct contact or exposure to this product or spray mixtures of this product should be minimized. The following instructions for transfer, mixing, cleaning or repairing equipment should be followed in order to minimize this exposure. Review the protective clothing requirements as listed in the "PRECAUTIONARY STATEMENTS" section of this label and do not use this product until you have the necessary protective clothing.

2.5 Gallon Containers

Open pouring from these containers can result in exposure from splashing or spilling. Special care in lifting and pouring is strongly recommended.

55 Gallon and Bulk Containers

Open pouring from these containers can result in exposure from splashing or spilling and is not recommended. This product should be transferred from these containers to the mix or spray tank using pumps or transfer probes. The probe or pump should not be removed from the container or disconnected until the container is emptied and rinsed. Use the pump or probe system to rinse the empty container and transfer the rinsate directly to the mix or spray tank.

Equipment Cleaning & Repair

14/68

Cleaning and repair of transfer systems and application equipment is a source of exposure to this product. Care should be taken to minimize exposure during cleaning and repair of transfer systems and application equipment. Whenever possible, these systems or equipment should be rinsed before being cleaned or repaired.

When repairs must be made during transfer or application, the equipment should be shut down and special care taken to avoid contact with the pesticide.

Mixing Instructions

Always predetermine the compatibility of this product or labeled tank mixtures of this product with water carrier or sprayable fluid fertilizer carrier by mixing small proportional quantities in advance. See the "STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST" section in this label to determine the compatibility of this product and the labeled tank mixtures recommended for use with sprayable fluid fertilizer carrier. **NOTE:** This product and all tank mixtures listed for corn, sorghum and soybeans in the "DIRECTIONS FOR USE" section may be applied in sprayable fluid fertilizers. DO NOT apply this product or labeled tank mixtures of this product in sprayable fluid fertilizers on any other crops.

Mix this product or labeled tank mixtures of this product with the appropriate carrier as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with the appropriate carrier.
3. If a compatibility agent is necessary to improve mixing or to prevent the formation of undesirable and unsprayable gels or precipitates, while agitating add it to the carrier already in the tank. Use only compatibility agents, cleared by FDA for this use. Read and follow all directions for use, cautionary statements and all other information appearing on the selected compatibility agent label. Check for adequate agitation.
4. If a wettable powder or dry flowable formulation is used, make a slurry with water and add it slowly through the screen into the tank. Continue agitation.
5. If a flowable formulation is used, add slowly through screen into the tank. Mixing and compatibility may be improved when flowable is premixed one part flowable with one part water and added to the tank in diluted form.
6. Add this product slowly through the screen into the tank. Mixing and compatibility may be improved when this product is prediluted with two parts of water and added to the tank in diluted form.
7. Complete filling the sprayer tank with carrier. If a Roundup® brand labeled for agricultural uses or Gramoxone™ Extra is used, add the required amount near the end of the filling process. Remove hose from tank immediately after filling to avoid siphoning back into the carrier source.

Maintain good agitation at all times until the contents of the tank are sprayed. **NOTE:** If spray mixture is allowed to settle at any time, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. Check for even distribution of spray droplets. To reduce loss of the chemical due to drift of a fine mist, apply at nozzle pressures below 40 psi.

STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST

Herbicides may not always mix evenly throughout a sprayable fluid fertilizer or the components may separate too quickly to make their combined use of practical value. This may be due to certain characteristics of the different fluid fertilizers. A simple test using small quantities of the components is suggested to provide compatibility potential.

The test follows:

A. Materials Required For A Compatibility Test

1. Two one-quart jars with lid or stopper (marked "with" and "without").
2. TEAspoons (for a more exacting test, a five to ten milliliter (mL) pipette or graduated cylinder is desirable).
3. Sprayable fluid fertilizer to be tested.
4. The herbicide chemicals to be mixed.
5. A compatibility agent (the purpose of the adjuvant is to help keep the fertilizer and crop protection chemical in suspension, if this assistance is needed).

B. Procedure

1. Add one pint of the sprayable fluid fertilizer that will be used or other herbicide carrier to each jar marked "with" and "without".

Add One Pint Liquid Fertilizer
To Two Quart Jars.

WITH

WITHOUT

2. To the jar marked "with", add 1/4 TEAspoon or 1.2 milliliters of a suitable compatibility agent; shake gently for five to ten seconds to mix (1/4 TEAspoon in one pint is the equivalent of two pints per 100 gallons of liquid fertilizer).

To Jar Marked "With"
Add Compatibility Agent
And Shake to Mix.

WITH

WITHOUT

3. To each jar add the appropriate amount of herbicide(s) listed in the following tables. If more than one is used, add them separately with the wettable powders or dry flowables added first, flowables second and liquid last. Shake gently five to ten seconds after each addition.

Add Herbicide(s) To Both Jars
And Shake to Mix.

WITH

WITHOUT

Amount to Be
Added Per Pint Of
Sprayable Fluid Fertilizer
(Assuming Volume is 25 gallons/Acre)

HERBICIDE	RATE/ACRE	Level TEAspoons
-----------	-----------	--------------------

16/68

Wettable	1 pound	=	1.5
Powders or	2 pounds	=	3
Dry Flowables	3 pounds	=	4.5
	4 pounds	=	6
	5 pounds	=	7.5

HERBICIDE	RATE/ACRE	Level TEAspoons	Milliliters
Emulsifiable	1 pint	= 0.5	or 2.4
Concentrates or	1 quart	= 1	or 4.7
Flowables or	2 quarts	= 2	or 9.5
Liquids or	3 quarts	= 3	or 14.2
Solutions	1 gallon	= 4	or 19.0
	5 quarts	= 5	or 23.8

This compatibility test is designed for 25 gallons of spray per acre. For changes in spray volume or herbicide rate, make appropriate changes in the ingredients of the test. Regardless of spray volume, the amount of compatibility agent should be equal to two or three pints (two pints = 1/4 TEAspoon or 1.2 milliliters, three pints = 3/8 TEAspoon or 1.8 milliliters per pint of sprayable fluid fertilizer) per 100 gallons of liquid fertilizer.

C. Observations and Decisions

1. If the herbicide(s) and the sprayable fluid fertilizer are compatible.
2. If a compatibility agent is necessary.

Five minutes after the final addition and mixing, observe both jars for the formation of large flakes, sludge, gels or other precipitates. Observe if the herbicide(s) cannot be physically mixed with the liquid fertilizer but remains as small oily particles in the solution.

If incompatibility in any form described above occurs in the jar "with" the compatibility agent added, the liquid fertilizer and the herbicide(s) should not be used together in the same spray tank.

If incompatibility as described above occurs in the jar "without" the adjuvant but not in the jar "with" adjuvant, the use of a compatibility adjuvant is recommended.

Both jars should be allowed to stand and be observed periodically for one-half hour. If the separate layers of liquid fertilizer and additives can be resuspended by shaking, commercial application is possible. An emulsifiable concentrate normally will go to the top after standing; wettable powders will either settle to the bottom of the tank or jar, or float to the top, depending upon the density of the fertilizers.

If the herbicide(s) is compatible with fluid fertilizer in the foregoing test without having to use a compatibility agent, fluid fertilizer may be used for the premixing. If it is not compatible without the compatibility agent, the herbicide(s) should be premixed with water before adding to the spray tank.

APPLICATION INFORMATION

APPLICATION SYSTEMS

Spray Drift Management

17/68

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parts and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

For ground boom applications, apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site as measured by an anemometer. Use medium or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles.

For aerial applications, the boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3-10 mph as measured by an anemometer. Use medium or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or the crop canopy.

For overhead chemigation, apply only when wind speed is 10 mph or less.

The applicator also must use all other measures necessary to control drift.

Ground Broadcast Treatment-For application into CONVENTIONAL TILLAGE SYSTEMS, apply this product and the labeled tank mixtures in 10 or more gallons of solution per acre using broadcast boom equipment. For application into CONSERVATION OR MINIMUM TILLAGE SYSTEMS, apply in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre when adding a Roundup brand labeled for agricultural uses or 2,4-D to the mix. Apply in 20 to 60 gallons of water or nitrogen solution when adding Gramoxone herbicide to the mix. The carrier may be either water or sprayable fluid fertilizer as specified for the crop to be treated in the "DIRECTIONS FOR USE" section of this label. Do not apply during periods of gusty winds, when winds are in excess of 15 mph or when other conditions favoring drift exist.

Band Treatment-Apply a broadcast equivalent rate and volume per acre. To determine these:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per acre} = \text{Band RATE per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per acre} = \text{Band VOLUME per acre}$$

Chemigation—Center Pivot Irrigation - Apply this product and labeled tank mixtures only through center pivot irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Use only in systems that apply uniformly.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

18/168

Do not chemigate through systems connected to a public water system.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Do not apply when system connections or fittings leak, when nozzles do not provide uniform distribution or when the line containing the product must be dismantled and drained.

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a systems interlock. Pumps, injection equipment, agitation equipment, hoses and connections between supply tank and the point of injection must be constructed of materials which are resistant to this product.

Meter this product or a labeled tank mixture of this product into the center pivot irrigation system after planting and before crop emergence. Herbicide application should be made in 1/3 to 3/4 inch of water per acre. Do not apply in more than 3/4 inch of water per acre under any conditions or reduced performance may occur. On very sandy soils (more than 60 percent sand and less than 1 percent organic matter), apply these herbicides in a maximum of 1/2 inch water per acre. Good agitation must be maintained during the entire application period. Flush the system with water when application is complete. Refer to the "MIXING, SPRAYING AND HANDLING INSTRUCTIONS" section of the label for mixing procedures.

NOTE: Applications can be made by injection through center pivot irrigation equipment only when specified for the crop to be treated in the "DIRECTIONS FOR USE" section of this label. Do not apply this product or any labeled tank mixture through center pivot irrigation unless the treatment is specifically recommended in this label for the crop to be treated.

Posting of areas to be chemigated is required when any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day-care centers, hospitals, in-patient clinics, nursing homes or any public area such as schools, parks, playgrounds or other public facilities not including public roads. Posting must conform to the following requirement. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are not usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Posting signs may be obtained from your local farm chemical dealer. This sign is in addition to any sign posted to comply with the Worker Protection Standard.

19/68

Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 ½ inches tall, and all letters and the symbol shall be a color that sharply contrasts with the immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

Aerial-Fixed wing; helicopter-Apply the recommended rates of this product or approved tank mixtures in 3 to 10 gallons of water per acre. To avoid streaked, uneven or overlapped application, use mechanical marking systems only. HUMAN FLAGGERS PROHIBITED!

Application with Dry Bulk Fertilizer-The herbicide-dry fertilizer impregnation process (In-Plant and On-Board systems) must be completed only by commercial fertilizer or chemical dealerships properly equipped for this procedure. Contact Monsanto Company for additional information regarding recommended equipment and methods for herbicide-fertilizer impregnation applications.

Dry bulk fertilizer may be impregnated with this product or the tank mixtures of this product plus atrazine on corn, this product or the tank mixture of this product plus atrazine on grain sorghum, and this product or the tank mixtures of this product plus Sencor™ on soybeans. This product and these tank mixtures must be applied with 200 to 450 pounds of dry bulk fertilizer per acre and shallowly incorporated within 7 days prior to planting. On medium and fine textured soils in areas where soil incorporation is not planned; i.e., reduced tillage situations or in some conventional till situations, applications can be made up to 30 days before planting to allow moisture to move the herbicide-fertilizer mixture into the soil. On coarse-textured soils, applications can be made up to 14 days prior to planting. The herbicide must be applied as recommended in this label for the crop, weed and soil type. Refer to the table for broadcast rate per acre in the appropriate crop section of this label to determine the recommended rate per acre for the herbicide treatment to be applied.

The following table provides a reference to determine the amount of LIQUID herbicide to be mixed per ton of dry bulk fertilizer for a range of herbicide recommendations for fertilizer rates per acre:

Fertilizer Rate (Pounds/Acre)	Acres Covered (Per Ton)	RECOMMENDED QUARTS LIQUID HERBICIDE/ACRE			
		1.5	2 (Quarts of Herbicide/Ton Dry Bulk Fertilizer)	2.5	3
200	10	15	20	25	30
250	8	12	16	20	24
300	6.7	10	13.3	16.7	20
350	5.7	8.6	11.4	14.3	17.1
400	5	7.5	10	12	15
450	4.5	6.7	8.9	11.1	13.3

To determine the amount of herbicide needed for rates not included in the preceding table, use the following formula:

$$\frac{\text{Recommended herbicide rate (pounds or quarts)/acre} \times 2000}{\text{pounds fertilizer/acre}} = \text{Pounds or quarts herbicide per ton of dry bulk fertilizer}$$

With the In-Plant system, mix and blend the dry fertilizer and herbicide mixture in a closed rotary drum-type mixer allowing sufficient time to ensure uniform coverage. Use at least one ton of dry fertilizer per

20/68

mixing operation. Inject the herbicide into the drum over a minimum of a 2-minute period and allow at least 2 additional minutes mixing time to ensure uniformity. The nozzle used to spray the herbicide treatment must be placed inside the mixer to provide uniform spray coverage of the tumbling fertilizer.

If the dry fertilizer used has inadequate absorptive capacity, a highly absorptive material such as Agsorb™, MP-79™, or Microcel™ can be added to provide a free-flowing mixture. Contact Monsanto Company for specific guidelines with regard to the sequence of addition for the various components and the amount of drying agent to add to provide a free-flowing mixture.

The following table provides a partial list of dry fertilizers which may be impregnated with this product or tank mixtures of this product with other herbicides:

Fertilizer	Lasso + Lasso	Lasso + atrazine	Lasso + Sencor
Ammonium sulfate (21-0-0)	Yes	Yes	Yes
Ammonium phosphate-sulfate (16-20-0)	Yes	Yes	Yes
Diammonium phosphate (18-46-0)	Yes	Yes	Yes
Potassium chloride (0-0-60)	Yes	Yes	Yes
Potassium sulfate (0-0-52)	Yes	Yes	Yes
Single super-phosphate (0-20-0)	Yes	No	Yes
Treble super-phosphate (0-46-0)	Yes	No	Yes
*Urea (46-0-0)	Yes	Yes	Yes

*Some ureas may be phytotoxic when applied on corn. Use only ureas known to be safe to corn.

NOTE: DO NOT impregnate this product or tank mixtures of this product with other herbicides on fertilizers containing ammonium nitrate, potassium nitrate or sodium nitrate.

Spread the herbicide-dry fertilizer mixture uniformly with a properly calibrated dribble, pneumatic (air flow) or spin applicator. When using spin applicators, the herbicide-dry fertilizer mixture must be spread at half-rate and overlapped 100 percent to obtain full rate and uniform distribution. Non-uniform spreading of the fertilizer-herbicide mixture may result in unsatisfactory weed control or crop injury.



2/1/68

APPLICATION TIMING AND METHODS

Early Preplant Surface-Some labeled tank mixtures of this product may be applied in no-till and other conservation tillage systems before weeds emerge and up to 45 days before planting certain crops. Split applications can be made 30 to 45 days prior to planting, with 60 percent of the recommended broadcast rate applied initially and the remaining 40 percent applied at planting. Applications made less than 30 days prior to planting can be made either as a split or as a single application. Refer to the individual crop to determine if early preplant application is recommended. If weeds are present at the time of application, apply this product in tank mixtures with an appropriate contact herbicide. Observe directions for use, precautions and restrictions on the label of the contact herbicide. During the planting operation, be careful not to move untreated soil to the surface or move treated soil out of the row, as weed control may be reduced.

Preplant Incorporation-This product and many of the labeled tank mixtures may be mixed into the soil using shallow incorporation equipment any time within 7 days prior to planting. Refer to the crop to be treated in the "DIRECTIONS FOR USE" section of this label to determine if preplant incorporation is recommended. Apply the recommended treatment rate to the soil surface as a broadcast application. Precipitation or irrigation is required to bring incorporated herbicide treatments into contact with germinating weed seedlings. The amount of precipitation or irrigation required depends on existing soil moisture, soil type and percent organic matter content, but 1/3 to 3/4 inch is normally adequate. If weeds emerge after treatment, rotary hoe or shallowly cultivate immediately to improve performance.

NOTE: Do not incorporate unless specifically recommended in this label for the crop and herbicide treatment selected. Incorporation of this product and its labeled tank mixtures on coarse soil types may reduce length of weed control due to leaching with rainfall or irrigation.

1-Pass Incorporation (surface blend): Shallowly incorporate the treatment into the upper 1 to 2 inches of the soil. Equipment should be operated at manufacturer's designed speed for incorporation to ensure adequate mixing and distribution of the herbicide treatment in the soil. Equipment design, including any drag attachments, must be adequate to avoid soil ridging which may result in streaked or reduced weed control. Soil conditions, including moisture content and crop residue levels, must be suitable to allow thorough and uniform mixing with the equipment used for 1-pass incorporation.

Contact Monsanto Company for additional information regarding recommended equipment and methods for Surface Blend application.

2-Pass Incorporation: Shallowly incorporate the herbicide treatment into the upper 1 to 2 inches of the soil with equipment set to work the soil NO DEEPER THAN 4 INCHES. The second pass must be made at an angle to and no deeper than the first pass to ensure proper distribution of the herbicide treatment in the soil.

Preemergence Surface Application-This product and all labeled tank mixtures except this product plus CommandTM or EptamTM or TreflanTM may be applied to the soil surface after planting and prior to either crop or weed emergence. Apply within 5 days of last preplant tillage. If weeds emerge after treatment, or if treatment is applied more than 5 days after last preplant tillage, rotary hoe or shallowly cultivate immediately to improve performance. Precipitation or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone. The amount of precipitation or overhead sprinkler irrigation required depends on existing soil moisture, soil type and percent organic matter content, but 1/3 to 3/4 inch is normally adequate. Performance is improved when moisture is received within 7 days after application and prior to weed emergence. High-intensity or excessive rainfall or excessive irrigation after application may reduce control.

Cultivation Information-Delay cultivation after application for as long as possible unless weeds or grasses emerge. Shallowly cultivate or rotary hoe immediately if weeds or grasses emerge. If cultivation is

22/68

necessary because of soil crusting or compaction, set equipment shallow and minimize lateral soil movement to avoid dilution or displacement of the herbicide treatment. If a band application is used and weeds have emerged in the treated band, set cultivator to throw soil into the row covering the band. On peanuts, do not throw soil into the row.

**Conservation or Minimum Tillage Systems Tank Mixtures
AT-PLANTING APPLICATIONS**

When applied as directed under the conditions described, the recommended tank mixtures control many emerged annual weeds, suppress many emerged perennial weeds and give preemergence control of many annual grasses and broadleaf weeds when corn, grain sorghum or soybeans will be planted directly into a cover crop, stale seedbed, established sod, or in previous crop residues. Consult the WEEDS CONTROLLED section of this label to determine the weeds that are controlled or reduced prior to their emergence, and the WEEDS CONTROLLED section of the Roundup brands labeled for agricultural uses, Gramoxone or 2,4-D labels to determine the weeds that are controlled or reduced following postemergence treatment by these products.

Apply the recommended tank mixtures BEFORE CROP EMERGENCE. As density of stubble, crop residue or weeds increases, spray volume and rate should be increased within the recommended ranges to ensure complete coverage. In the absence of emerged vegetation, delete the Roundup brands labeled for agricultural uses; 2,4-D or Gramoxone Extra portion of these tank mixtures. Follow the recommendations in the preceding sections of this label and also the label of the contact herbicide that is included in the tank mixture to provide postemergence control of weeds.

APPLY ONLY BY GROUND APPLICATION SYSTEMS. DO NOT APPLY BY AIR.

ATTENTION

AVOID DRIFT - EXTREME CARE MUST BE USED WHEN APPLYING THESE TANK MIXTURES TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS. Do not allow spray mist to drift because even minute quantities of spray can cause severe damage or destruction to nearby crops, plants or other areas on which treatment is not intended. Do not apply when winds are gusty or in excess of 5 mph or when other conditions, including lesser wind velocities, will allow drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in fine particles (mist) that are more likely to drift.

For emerged annual or perennial weeds controlled or suppressed, see the "WEEDS CONTROLLED" section of the Roundup brands labeled for agricultural uses; 2,4-D or Gramoxone Extra labels. USE OF THESE MIXTURES FOR BERMUDAGRASS OR JOHNSONGRASS CONTROL IS NOT RECOMMENDED.

Ammonium Sulfate—The addition of ammonium sulfate in the spray solution may increase the performance of Roundup brands labeled for agricultural uses tank mixtures on emerged annual weeds under adverse growing conditions. When using ammonium sulfate, add 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water. Ammonium sulfate should be added to the water in the spray tank and completely dissolved prior to adding the herbicide or surfactant. Do not mix ammonium sulfate in fluid fertilizer solutions.

If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet lines. Nozzle tip plugging may result from the use of low-quality ammonium sulfate. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for one minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to adding to the spray tank.

Following use of ammonium sulfate, thoroughly rinse the spray system with clean water to reduce corrosion.

23/168

Surfactants—Nonionic surfactants which are labeled for use with herbicides may be used with some Roundup brands labeled for agricultural uses. Check specific label for restrictions. When allowed, do not reduce rates of Roundup brands labeled for agricultural uses when adding surfactant. Use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 50 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

WEEDS CONTROLLED

When applied as directed under conditions described, this product and tank mixtures of this product will control or reduce competition from the weeds listed. Refer to the crop on which treatment is intended to determine the rates and tank mixtures recommended.

For the scientific names of the weeds listed below, refer to the appendix at the end of this booklet.

The tank mixtures of this product plus atrazine or plus Princep™ may not control triazine-resistant weeds as indicated in the tables below for these mixtures.

NOTE: C=Control R=Reduced Competition

ANNUAL GRASSES

	Lasso plus				
	Lasso	atrazine	dicamba	Canopy™	Command™
Barnyardgrass	C	C	C	C	C
Crabgrass	C	C	C	C	C
Cupgrass, woolly ¹	R	R	•	R	R
Foxtail: giant; green; robust purple, robust white, yellow	C	C	C	C	C
Goosegrass	C	C	C	C	C
Johnsongrass, seedling ²	R	R	R	R	R
Millet, proso ³	R	R	•	•	•
Panicum: browntop, fall	C	C	C	C	C
Panicum, Texas ⁴	R	R	R	R	R
Rice, red ⁵	C	C	•	C	C
Sandbur, Grassbur ^{2,6}	R	C	•	•	•
Shattercane, wild cane ¹	R	R	•	R	R
Signalgrass, broadleaf	C	C	C	C	C
Sprangletop, red	C	C	C	C	C
Witchgrass	C	C	C	C	C

ANNUAL BROADLEAVES

24/68

	Lasso plus				
	Lasso	atrazine	dicamba	Canopy™	Command™
Beggarweed, Florida ²	R	C	•	C	•
Carpetweed	C	C	C	C	C
Cocklebur	•	R	R	C	•
Coffeeweed, Hemp sesbania	•	•	•	C	•
Copperleaf, hophornbeam	•	•	•	C	•
Galinsoga	C	C	C	C	C
Groundcherry, annual	•	C	•	C	•
Groundcherry, cutleaf ⁷	C	C	C	C	C
Jimsonweed	•	C	•	C	C
Kochia	•	C	•	•	C
Lambsquarters ²	R	C	C	C	C
Morningglory: tall ⁸ ,	•	R	R	C	•
pitted, ivyleaf ⁸ , entireleaf ⁸ ,	•	•	•	C	•
smallflower	R	C	•	C	•
Mustard	•	C	C	C	•
Nightshade: black ⁷ ; hairy ⁷	C	C	C	C	C
Pigweed, Carelessweed	C	C	C	C	C
Purslane	C	C	C	C	C
Pusley, Florida	C	C	C	C	C
Ragweed, common ²	R	C	C	C	C
Ragweed, giant	•	•	•	R	•
Sicklepod	•	C	•	C	•
Sida, prickly; Teaweed ²	R	•	•	C	C
Smartweed ²	R	C	C	C	R
Starbur, bristly	R	C	•	C	•
Sunflower, common	•	•	•	R	•
Velvetleaf, Buttonweed ⁸	•	R	R	C	C
Waterhemp	C	C	C	C	C
SEDGE					
Nutsedge, yellow ⁵	C	C	•	C	C

ANNUAL GRASSES

Lasso plus				
------------	--	--	--	--

25/68

	linuron	Lorox™Plus	Pursuit™	Scepter™	Sencor™	Treflan™
Barnyardgrass	C	C	C	C	C	C
Crabgrass	C	C	C	C	C	C
Cupgrass, woolly ¹	•	•	R	R	R	R
Foxtail: giant; green; robust purple; robust white; yellow	C	C	C	C	C	C
Goosegrass	C	C	C	C	C	C
Johnsongrass, seedling ²	R	R	C	C	R	R
Millet, proso ³	•	•	•	•	•	•
Panicum: browntop; fall Panicum, Texas ⁴	C	C	C	C	C	C
Rice, red ⁵	R	R	R	R	R	R
Sandbur, Grassbur ^{2,6}	•	•	•	•	•	•
Shattercane, wild cane ¹	•	•	R	R	R	R
Signalgrass, broadleaf	C	C	C	C	C	C
Sprangletop, red	C	C	C	C	C	C
Witchgrass	C	C	C	C	C	C

ANNUAL BROADLEAVES

	Lasso plus					
	linuron	Lorox™Plus	Pursuit™	Scepter™	Sencor™	Treflan™
Beggarweed, Florida ²	•	•	•	•	R	•
Carpetweed	C	C	C	C	C	C
Cocklebur	•	C	C	C	R	•
Coffeeweed	•	•	•	•	•	•
Copperleaf, hophornbeam	•	•	•	•	•	•
Galinsoga	C	C	C	C	C	C
Groundcherry, annual	C	C	C	C	C	•
Groundcherry, cutleaf ⁷	C	C	C	C	R	C
Jimsonweed	•	C	C	C	R	•

Lasso 2001-1-Sep1701

26/168

Kochia	•	•	•	•	•	C
Lambsquarters ²	C	C	C	C	C	•
Morningglory: tall ⁸ ,	•	R	R	R	•	•
pitted; ivyleaf ⁸ ,	•	R	R	C	•	•
entireleaf ⁸ ,	•	R	R	R	•	•
smallflower	•	•	C	•	C	•
Mustard	C	C	C	C	•	•
Nightshade: black ⁷ ; hairy ⁷	C	C	C	C	C	C
Pigweed, Carelessweed	C	C	C	C	C	C
Purslane	C	C	C	C	C	C
Pusley, Florida	C	C	C	C	C	C
Ragweed, common ²	C	C	R	C	C	•
Ragweed, giant	•	R	R	R	•	•
Sicklepod	•	•	•	R	R	•
Sida, prickly; Teaweed ²	C	C	C	C	C	•
Smartweed ²	C	C	C	C	C	•
Starbur, bristly	•	•	•	C	C	•
Sunflower, common	•	•	C	C	C	•
Velvetleaf, Buttonweed ⁸	R	C	C	C	R	•
Waterhemp	C	C	C	C	C	C

SEDGE

Nutsedge, yellow ⁵	•	•	C	C	C	C
-------------------------------	---	---	---	---	---	---

- 1 Use 4 quarts per acre of this product applied alone or in tank-mix combinations. Preplant incorporate only to reduce competition from this weed.
- 2 For reduced competition from this weed using this product alone, use 2.5 to 4 quarts of Lasso per acre.
- 3 Plant corn into freshly prepared seedbed. Apply 4 quarts of this product per acre preemergence surface immediately after corn planting to reduce competition from this weed.
- 4 Use 3 to 4 quarts of this product per acre and apply through center pivot irrigation or under sprinkler irrigation systems. DO NOT apply this product plus atrazine tank mixture through center pivot irrigation systems.
- 5 Use 3 to 4 quarts of this product per acre applied alone or in tank mixture and apply preplant incorporated only for control on medium-and fine-textured soils. When tank-mixed with Pursuit or Scepter, use 2 to 3 quarts of this product per acre and preplant incorporate only. On coarse-textured soils with less than 3 percent organic matter, use 3 to 4 quarts of this product per acre as a preemergence surface application to reduce competition from yellow nutsedge.
- 6 Apply 2.5 to 3 quarts of this product plus 1 to 2 quarts of atrazine 4L per acre. Preplant incorporated applications should be made immediately before planting. Preemergence surface application should be applied to a freshly prepared seedbed. For best control, rotary hoe or initiate overhead irrigation (1/3 to 1/2 inch) within 24 hours of application.
- 7 Use a minimum of 2.5 quarts of Lasso per acre on coarse soils, or 3 quarts of this product per acre on medium and fine soils when used alone or in tank mixture with Banvel™ or equivalent rates of other dicamba products registered for this use, Canopy, Command, Sencor or Treflan for control of this weed.
- 8 For best results on these species when using this product plus Scepter tank mixture, use preplant incorporated treatments.

27/68

Certain tank mixtures of this product and broadleaf herbicides may be combined with applications of Roundup brands labeled for agricultural uses or Gramoxone Extra for control of many emerged weeds prior to corn or soybean emergence. Refer to the Roundup brands labeled for agricultural uses or Gramoxone Extra label for a list of emerged weeds controlled by tank mixtures with these products.

TM

Read each of these sections of this label for essential product performance information.

CROPS

CORN, SORGHUM, SOYBEANS, PEANUTS, BEANS, WOODY ORNAMENTALS

CORN CONTAINING THE ROUNDUP READY® GENE

This product may be used in Roundup Ready corn, or corn that contains the Roundup Ready gene. A minimum of 1.25 quarts per acre of this product is required. Use the full labeled rate of this product when the application is 14 days or more prior to planting or when the following grasses may be expected: barnyardgrass, shattercane, seedling johnsongrass or sandbur.

Preemergence Sequential Program

This product may be used at a reduced rate preemergence in a sequential program with Roundup UltraMAX™ herbicide, or other Roundup brands labeled for use in Roundup Ready corn. Use 50% to 100% of the labeled rate of this product for the appropriate soil texture and organic matter, per the Application Rate Table. Apply a sequential application of Roundup UltraMAX, or other Roundup brand labeled for use in Roundup Ready corn, per the label instructions.

If weeds are present at the time of preemergence application, tankmix Roundup UltraMAX or other Roundup brand labeled for agricultural uses, to control emerged vegetation.

**CORN—CONVENTIONAL TILLAGE
(All Types)**

NOTE: Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the "GENERAL INFORMATION", the specific information should control.

When making applications on corn, do not make more than two applications per year or exceed a total of 4 quarts of this product per acre per year. Do not make more than one application or exceed 4 quarts of this product per acre per year on sweet corn.

28/168

LASSO

Apply this product in water or sprayable fluid fertilizer solution.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Center Pivot Irrigation; Dry Bulk Fertilizer Impregnation

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

Sequential Treatment-This product may be applied as two sequential treatments for improved control of hard-to-control weeds. The combined total application rate for the two treatments must not exceed 4 quarts per acre per year. The second treatment must be applied before the corn emerges.

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)*	
	Less than 3% organic matter	3% or more organic matter
Coarse	2.5	2.5
Medium	2.5 to 3	2.5 to 3
Fine	2.5 to 3	3 to 3.5

*Use 2.5 to 3.5 quarts per acre where furrow irrigation is used. Use a minimum of 3 quarts on coarse-textured soils for control of yellow nutsedge or red rice. Rates may be increased to a maximum of 4 quarts on any soil type when heavy infestation of yellow nutsedge or red rice are present.

On muck or peat or coarse soils with 10 percent or more organic matter use 4 quarts of this product per acre.

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)*	
	Less than 3% organic matter	3% or more organic matter
Coarse**	2 to 2.25	2 to 2.25
Medium	2 to 2.75	2 to 2.75
Fine	2 to 2.75	2.5 to 3.25

29/68

*Use a minimum of 2.5 quarts of this product per acre on coarse soils and 3 to 4 quarts on medium-or fine textured soils to control black or hairy nightshade.

**When applied through center pivot irrigation or under sprinkler irrigation systems on coarse soils, use a minimum of 3 quarts per acre of this product.

On muck or peat soils or on coarse soils with 10 percent or more organic matter use 4 quarts of this product per acre.

**LASSO plus ATRAZINE
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.
DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Dry Bulk Fertilizer Impregnation

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

When applied following Lasso or tank mixtures of Lasso, the total amount of this product applied alone or in combination must not exceed 4 quarts per acre per year. The total amount of atrazine 4L must not exceed 3 quarts per acre per year.

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)					
	Less than 3% organic matter			3% or more organic matter		
	Lasso	+	atrazine 4L*	Lasso	+	atrazine 4L*
Coarse	1.75 to 2.25		1.25 to 2	1.75 to 2.25		1.25 to 2
Medium	1.75 to 2.25		1.25 to 2	2.25 to 2.5		1.5 to 2
Fine	2.25 to 2.5		1.5 to 2	2.25 to 2.75		1.5 to 2

30/68

*Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)					
	Less than 3% organic matter			3% or more organic matter		
	Lasso	+	atrazine 4L*	Lasso	+	atrazine 4L*
Coarse	1.5 to 2		0.75 to 2	1.5 to 2		1 to 2
Medium	1.5 to 2		1 to 2	2 to 2.25		1.25 to 2
Fine	2 to 2.25		1.25 to 2	2 to 2.75		1.25 to 2

*Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

NOTE: When applied under sprinkler irrigation systems on coarse soils, use 1 quart per acre of atrazine 4L and a minimum of 2 quarts per acre of this product in this tank mixture.

**LASSO plus BALANCE PRO
(TANK MIXTURE)**

FOR USE IN FIELD CORN AND SILAGE CORN

APPROVED APPLICATION SYSTEMS

Ground – Broadcast boom; banded.

APPROVED APPLICATION METHOD

Preplant Incorporated, Preemergence Surface

Balance PRO is not registered in all states. Follow all Restrictions and Precautions on the Balance PRO label including planting depth, environmental precautions, and soil type restrictions.

Follow the Balance PRO Technical Bulletins, 24(c) labels and 2(ee) Recommendations for additional **use rate restrictions** based on soil textures and depth to groundwater in various states.

APPLICATION RATES:

31/68

Preplant Incorporate: BROADCAST RATE PER ACRE (quarts of Lasso and ounces of Balance PRO)

SOIL TEXTURE	< 3% ORGANIC MATTER		> 3% ORGANIC MATTER	
	Lasso (quarts)	Balance PRO (ounces)	Lasso (quarts)	Balance PRO (ounces)
COARSE *	2.0 to 2.25 +	1.0 to 1.88	2.0 to 2.25 +	1.0 to 1.88
MEDIUM	2.0 to 2.75 +	1.0 to 2.5	2.0 to 2.75 +	1.5 to 3.0
FINE	2.0 to 2.75 +	1.5 to 2.75	2.5 to 3.25 +	2.0 to 3.5

* It is not recommended to use Balance PRO on coarse soils < 1.5% organic matter.

Preemergence Surface: BROADCAST RATE PER ACRE (quarts of Lasso and ounces of Balance PRO)

SOIL TEXTURE	< 3% ORGANIC MATTER		> 3% ORGANIC MATTER	
	Lasso (quarts)	Balance PRO (ounces)	Lasso (quarts)	Balance PRO (ounces)
COARSE*	2.0 to 2.25 +	1.0 to 1.88	2.0 to 2.25 +	1.0 to 1.88
MEDIUM	2.0 to 2.75 +	1.0 to 2.5	2.0 to 2.75 +	1.5 to 3.0
FINE	2.0 to 2.75 +	1.5 to 2.75	2.5 to 3.25 +	2.0 to 3.5

* It is not recommended to use Balance PRO on coarse soils < 1.5% organic matter.

**LASSO plus BANVEL
(TANK MIXTURE)
Field and Silage Corn Only**

For use in ILLINOIS, IOWA, MINNESOTA and WISCONSIN on level or flat-planted field corn on fine-textured (silty clay loam, clay loam, sandy clay, silty clay or clay) soils with more than 4 percent organic matter.

Apply this tank mixture in water or sprayable fluid fertilizer solution.

NOTE: Use on coarse-or medium-textured soils or on fine textured soils with 4 percent or less organic matter may result in crop injury and/or destruction.

APPROVED APPLICATION SYSTEMS

Ground:Broadcast boom, banded

NOTE: DO NOT apply by air or injection through center pivot irrigation systems.

APPROVED APPLICATION METHODS

Preemergence Surface- Corn seeds must be planted 1-1/2 inches or deeper beneath the soil surface. Direct chemical contact with corn seed must be avoided since crop injury may result. Apply far enough behind planter equipment to avoid any incorporation by the planter wheel or other covering device. If corn seeds are planted less than 1-1/2 inches beneath the soil surface, delay application until corn has spiked.

NOTE: PREVENT DRIFT TO SOYBEANS OR OTHER DESIRABLE PLANTS. Do not use on furrow irrigated corn, or when corn is planted at the bottom of a furrow, utilizing lister, till or other similar planting methods. DO NOT incorporate prior to planting or corn emergence. If it is necessary to drag for leveling or rotary hoe to break soil crust, DO NOT disturb the soil more than 1/2 inch deep.

APPLICATION RATES

309/68

BROADCAST RATE PER ACRE (QUARTS)

SOIL TEXTURE	More than 4% organic matter	
	Lasso	Banvel ¹
Fine-silty clay loam through clay	2.5 to 3	0.5

¹ Banvel or equivalent rates of other dicamba products registered for this use.

**LASSO plus MARKSMAN™
(TANK MIXTURE)
Field and Silage Corn Only**

For use in ILLINOIS, IOWA, MINNESOTA and WISCONSIN on level or flat-planted field corn on fine-textured (silty clay loam, clay loam, sandy clay, silty clay or clay) soils with more than 4 percent organic matter.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

Apply this tank mixture in water or sprayable fluid fertilizer solution.

NOTE: Use on coarse- or medium-textured soils or on fine-textured soils with 4 percent or less organic matter may result in crop injury and/or destruction.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded

NOTE: DO NOT apply by air or injection through center pivot irrigation systems.

APPROVED APPLICATION METHODS

Preemergence Surface- Corn seeds must be planted 1-1/2 inches or deeper beneath the soil surface. Direct chemical contact with corn seed must be avoided since crop injury may result. Apply far enough behind planter equipment to avoid any incorporation by the planter wheel or other covering device. If corn seeds are planted less than 1-1/2 inches beneath the soil surface, delay application until corn has spiked.

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

NOTE: PREVENT DRIFT TO SOYBEANS OR OTHER DESIRABLE PLANTS. Do not use on furrow irrigated corn, or when corn is planted at the bottom of a furrow, utilizing lister, till or other similar planting methods. DO NOT incorporate prior to planting or corn emergence. If it is necessary to drag for leveling or rotary hoe to break soil crust, DO NOT disturb the soil more than 1/2 inch deep.

APPLICATION RATES:

BROADCAST RATE PER ACRE

Lasso 2001-1-Sep1701

33/168

SOIL TEXTURE	More than 4% organic matter	
	Lasso (quarts)	Marksman (pints)
Fine-silty clay loam through clay	2.5 to 3	3.5

LASSO plus PRINCEP (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution.
APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Dry Bulk Fertilizer Impregnation

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP, PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS.

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)					
	1% to 3% organic matter			3% or more organic matter		
	Lasso	+	Princep 4L*	Lasso	+	Princep 4L*
Coarse-sandy loam only	1.75 to 2.25		1.25 to 2	1.75 to 2.25		1.25 to 2
Medium	1.75 to 2.25		1.25 to 2	2.25 to 2.5		1.5 to 2
Fine	2.25 to 2.5		1.5 to 2	2.25 to 2.75		1.5 to 2

39/68

*Use rates listed in this label when using Princep 4L. Use equivalent rates when using Princep 80W or 90% dry flowable formulations. One quart of the 4L formulation equals 1.25 pounds of the 80W and 1.1 pounds of the 90% dry flowable formulations.

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)					
	1% to 3% organic matter			3% or more organic matter		
	Lasso	+	Princep 4L*	Lasso	+	Princep 4L*
Coarse-sandy loam only	1.5 to 2		0.75 to 2	1.5 to 2		1 to 2
Medium	1.5 to 2		1 to 2	2 to 2.25		1.25 to 2
Fine	2 to 2.25		1.25 to 2	2 to 2.75		1.25 to 2

*Use rates listed in this label when using Princep 4L. Use equivalent rates when using Princep 80W or 90% dry flowable formulations. One quart of the 4L formulation equals 1.25 pounds of the 80W formulation and 1.1 pounds of the 90% dry flowable formulations.

**LASSO plus ATRAZINE plus PROWL™
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of triazine-resistant lambsquarters and pigweed.
DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter

APPROVED APPLICATION METHOD

Preemergence Surface

DO NOT preplant incorporate this tank mixture as serious crop injury can result.

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

APPLICATION RATES:

Lasso 2001-1-Sep1701

35/168

BROADCAST RATE PER ACRE (QUARTS)

Less than 3% organic matter

SOIL TEXTURE	Lasso	+	atrazine 4L* +	Prowl
Coarse	1.5 to 2		0.75 to 2	0.75
Medium	1.5 to 2		1 to 2	0.75 to 1
Fine	2 to 2.25		1.25 to 2	1

BROADCAST RATE PER ACRE (QUARTS)

3% or more organic matter

SOIL TEXTURE	Lasso	+	atrazine 4L* +	Prowl
Coarse	1.5 to 2		1 to 2	0.75 to 1
Medium	2 to 2.25		1.25 to 2	1
Fine	2 to 2.75		1.25 to 2	1

*Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

**LASSO plus ATRAZINE plus LINURON
(TANK MIXTURE)
FOR USE ONLY EAST OF THE ROCKY MOUNTAINS.**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of triazine-resistant lambsquarters and pigweed.
DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded

APPROVED APPLICATION METHOD

Preemergence Surface

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

36/168

APPLICATION RATES:

BROADCAST RATE PER ACRE (QUARTS)

Less than 3% organic matter

SOIL TEXTURE	Lasso	+	atrazine 4L*	+	linuron4L**
Coarse-sandy loam only	1.5 to 2		0.75 to 2		0.33
Medium	1.5 to 2		1 to 2		0.5
Fine	2 to 2.25		1.25 to 2		0.5

BROADCAST RATE PER ACRE (QUARTS)

3% or more organic matter

SOIL TEXTURE	Lasso	+	atrazine 4L*	+	linuron 4L**
Coarse-sandy loam only	1.5 to 2		1 to 2		0.5
Medium	2 to 2.25		1.25 to 2		0.5
Fine	2 to 2.75		1.25 to 2		0.5

*Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

**Use the rates listed in this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50% dry flowable formulations. One quart of linuron 4L equals 2 pounds of linuron 50WP or 2 pounds of linuron 50% dry flowable.

CORN-CONSERVATION OR MINIMUM TILLAGE

(All Types)

NOTE: Important information regarding Conservation or Minimum Tillage appears in the Application Information section at the front of this label and should be carefully reviewed in conjunction with the information in this section.

Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

37/168

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the "GENERAL INFORMATION", the specific information should control.

This product and the following tank mixtures can be tank mixed with Roundup brands labeled for agricultural uses, Gramoxone Extra and/or 2,4-D.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom

ROUNDUP UltraMAX

Annual Weeds-Apply 13 to 20 ounces of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, in these tank mixtures when weeds are less than 6 inches tall. Use 26 ounces of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, for control of fall panicum, barnyardgrass, crabgrass and shattercane up to 2 inches tall; Pennsylvania smartweed up to 6 inches tall; other annual weeds 6 to 12 inches tall.

Perennial Weeds-At normal application dates in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. Use of 1.6 to 3.25 quarts of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, in the above mixtures under these conditions provides top kill and reduces competition from many emerged perennial grasses and broadleaf weeds.

NOTE: When using these tank mixtures, do not exceed 3.25 quarts of Roundup UltraMAX per acre or equivalent rates of other Roundup brands labeled for agricultural uses.

GRAMOXONE EXTRA

Broadcast Treatment-Apply 1.5 to 3 pints of Gramoxone Extra in these tank mixtures immediately before, during or after planting. Use 1.5 to 2 pints when weeds are 1 to 3 inches tall, 2 to 2.5 pints when weeds are 3 to 6 inches tall and 2.5 to 3 pints when weeds are 6 inches tall. This mixture may not control weeds taller than 6 inches. Add a nonionic spreader surfactant (approved for use on crops) containing at least 75 percent surfactant active agent at 8 ounces per 100 gallons of diluted spray.

2,4-D

Broadcast Treatment-Apply 1 to 2 pints of 2,4-D (amine or low-volatile ester) in these tank mixtures. DO NOT use 2,4-D on light, sandy soils, or where soil moisture is inadequate for normal weed growth.

PREEMERGENCE WEED CONTROL

LASSO

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)
	Lasso
Coarse	2.5

38/168

Medium	3
Fine	3 to 3.5

Use the higher rate of in the recommended ranges in areas where organic matter content is 3 percent or more.

LASSO plus ATRAZINE

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)	
	Lasso	+ atrazine 4L*
Coarse-sandy loam only	1.75 to 2	1 to 2
Medium	2 to 2.5	1.2 to 2
Fine	2.5 to 3	1.6 to 2

*Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

Use the higher rate of this product in the recommended ranges when fall panicum or crabgrass will be present. Use the higher rate of atrazine in the recommended ranges on soils with greater than 3 percent organic matter.

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

**LASSO plus BALANCE PRO
(TANK MIXTURE)**

FOR USE IN FIELD CORN AND SILAGE CORN

Balance PRO is not registered in all states. Follow all Restrictions and Precautions on the Balance PRO label including planting depth, environmental precautions, and soil type restrictions.

Follow the Balance PRO Technical Bulletins, 24(c) labels and 2(ee) Recommendations for additional **use rate restrictions** based on soil textures and depth to groundwater in various states.

APPLICATION RATES:

59/68

BROADCAST RATE PER ACRE (quarts of Lasso and ounces of Balance PRO)

SOIL TEXTURE	< 3% ORGANIC MATTER		> 3% ORGANIC MATTER	
	Lasso (quarts)	Balance PRO (ounces)	Lasso (quarts)	Balance PRO (ounces)
COARSE *	2 to 2.25	+ 1.0 to 1.88	2 to 2.25	+ 1.0 to 1.88
MEDIUM	2 to 2.75	+ 1.0 to 2.5	2 to 2.75	+ 1.5 to 3.0
FINE	2 to 2.75	+ 1.5 to 2.75	2.5 to 3.25	+ 2.0 to 3.5

* It is not recommended to use Balance PRO on coarse soils < 1.5% organic matter.

LASSO plus PRINCEP

APPLICATION RATES:

BROADCAST RATE PER ACRE (QUARTS)

SOIL TEXTURE	Lasso	+	Princep 4L*
Coarse-sandy loam only	1.75 to 2		1 to 1.2
Medium	2 to 2.5		1.2 to 1.6
Fine	2.5 to 3		1.6 to 2.2

*When using Princep 80W use equivalent rates. One quart of Princep 4L equals 1.25 pounds of Princep 80W.

Use the higher rate of this product in the recommended ranges when fall panicum or crabgrass will be present.

Use the higher rate of Princep in the recommended ranges on soils with greater than 3 percent organic matter.

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP, PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS.

LASSO plus ATRAZINE plus PRINCEP

For additional preemergence control of heavy weed infestation and/or hard-to-control weeds, a three-way tank mixture of this product herbicide plus atrazine plus Princep may be applied.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

40/68

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)*				
	Lasso	+	atrazine 4L***	+	Princep 4L***
Coarse	1.75 to 2		0.75 to 1		0.75 to 1
Medium	2 to 2.5		1 to 1.25		1 to 1.5
Fine	2.5 to 3		1 to 1.25		1 to 1.5

*Use the higher rates of atrazine and Princep in the recommended ranges on soils with greater than 3 percent organic matter.

**When using 4L formulations use the rates listed. Use equivalent rates when using 80W or 90% dry flowable formulations. One quart of the 4L formulation of these products equals 1.25 pounds of the 80W formulation. One quart of atrazine 4L or Princep 4L equals 1.1 pounds of the 90% dry flowable formulations of these products.

Use the higher rate of this product in the recommended ranges when fall panicum, crabgrass, or soils with greater than 3 percent organic matter will be present.

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP, PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDING ROTATIONAL CROPS. CORN, PEANUTS, SORGHUM OR SOYBEANS* CAN BE PLANTED THE YEAR FOLLOWING USE OF THIS MIXTURE.

*There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year.

DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

EARLY PREPLANT APPLICATION

**LASSO plus ATRAZINE
(TANK MIXTURE)**

For use in no-till and other conservation tillage systems in CONNECTICUT, DELAWARE, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA, MISSOURI, NEBRASKA, NEW JERSEY, NEW YORK, NORTH CAROLINA, OHIO, PENNSYLVANIA, SOUTH DAKOTA, VERMONT, VIRGINIA, AND WISCONSIN.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

This tank mixture, when applied in a single application (alone or in a 3-way combination with Princep), split application or as a sequential application to Princep in early preplant programs, will provide preemergence control or reduced competition of listed annual grasses and broadleaf weeds.

If emerged weeds are present at the time of treatment, Roundup brands labeled for agricultural uses, Gramoxone Extra or 2,4-D should be added to the product according to the directions for use on their respective product labels. If unsatisfactory weed control occurs (due to excessively dry or excessively wet

41/68

conditions) following the earlier application, a postemergence application of an appropriate labeled grass and/or broadleaf weed herbicide may be used. If a postemergence treatment includes the herbicide used early preplant, do not exceed the labeled rate for corn on a given soil texture. DO NOT apply tank mixtures containing Roundup brands labeled for agricultural uses, Gramoxone Extra or other contact herbicides by air.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom; Aerial: Fixed wing, helicopter; Dry Bulk Fertilizer Impregnation

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application-Application of this tank mixture should be made less than 30 days before planting but prior to weed emergence.

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)	
	Lasso	+ atrazine 4L
Coarse*	2.25 to 2.5	1.25 to 1.5
Medium	2.5 to 3	1.5 to 2
Fine	3 to 3.5	1.75 to 2.25

*Applications on coarse soils should not be made more than 2 weeks prior to planting.

This tank mixture may also be mixed in a 3-way combination with 1 to 1.25 quarts per acre of Princep to provide improved control of fall panicum and crabgrass.

Split application-Apply 60 percent of the recommended rate as a split treatment prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting.

FOR CONTROL OF FALL PANICUM, CRABGRASS OR BROADLEAF SIGNALGRASS, USE THE SEQUENTIAL APPLICATION OF THIS TANK MIXTURE FOLLOWING PRINCEP.

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)	
	Lasso	+ atrazine 4L
Coarse	2.25 to 2.5	1.25 to 1.5
Medium	2.5 to 3	1.5 to 2

42/68

Fine

3 to 3.5

1.75 to 2.25

Sequential application-Application of this tank mixture following Princep should be utilized for the control of fall panicum, crabgrass or broadleaf signalgrass. Apply 1.5 to 2 quarts per acre of 1 to 1.25 quarts per acre of Princep prior to weed emergence and no more than 45 days prior to planting. At or immediately following planting apply the recommended rate of this tank mixture.

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)		
	Lasso	+	atrazine 4L
Coarse		1.75 to 2.5	1 to 1.5
Medium		2.25 to 2.75	1.25 to 1.75
Fine		2.5 to 3	1.5 to 2

Use rates listed in this label when using the 4L formulation of atrazine or Princep. Use equivalent rates of the 80W or 90% dry flowable formulations of atrazine or Princep. One quart of the 4L formulation of atrazine or Princep equals 1.25 pounds of the 80W and 1.1 pounds of the 90% dry flowable formulations of atrazine or Princep.

CORN, PEANUTS, SORGHUM (MILO) OR SOYBEANS can be planted the year following the use of this program. There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year.

NOTE: Soil treated with Princep should not be planted to any crop except corn for one year following treatment as crop injury may occur. After harvest of the treated crop, plow and thoroughly till the soil in the fall or spring to minimize possible injury to spring seeded rotational crops.

GRAIN SORGHUM (MILO)—CONVENTIONAL TILLAGE

NOTE: Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding APPLICATION SYSTEMS and APPLICATION TIMINGS AND METHODS appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the "GENERAL INFORMATION", the specific information should control.

When making applications on grain sorghum, do not make more than two applications per year or exceed a total of 4 quarts of this product per acre per year.

43/68

Do not graze or harvest forage for 70 days following application of this product or tank mixtures of this product.

LASSO

Apply this product in water or sprayable fluid fertilizer solution. Applications must be made ONLY to grain sorghum planted with seed that has been properly treated with a seed protectant or safener.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Center Pivot Irrigation; Dry Bulk Fertilizer Impregnation

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)*
	Lasso
Coarse	2 to 2.5
Medium	2.5 to 2.75
Fine	2.5 to 3

*Rates may be increased to a maximum of 4 quarts on any soil type when heavy infestation of yellow nutsedge is present.

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)
	Lasso
Coarse	1.5 to 2
Medium	2 to 2.25
Fine	2 to 2.5

44/68

NOTE: When applied through center pivot irrigation or under sprinkler irrigation systems, use a minimum of 3 quarts per acre of this product on coarse soils.

**LASSO plus ATRAZINE
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution. Applications must be made **ONLY** to grain sorghum planted with seed that has been properly treated with a seed protectant or safener.

NOTE: In Texas, use only in the Panhandle area and the fine-textured soils of the Gulf Coast and Blacklands. In the Texas Panhandle and Oklahoma Panhandle, apply this tank mixture as a preemergence surface application only. In the Texas Panhandle, Oklahoma Panhandle and the fine-textured soils of the Gulf Coast and Blacklands of Texas, do not exceed 1.25 quarts of atrazine 4L per acre as crop injury may result due to atrazine.

Applications made to grain sorghum (milo) growing on alkali soils or where cuts, fills or erosion have exposed calcareous or alkali subsoils may result in crop injury.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom, banded; Aerial-Fixed wing, helicopter; Dry Bulk Fertilizer Impregnation

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. **DO NOT** plant soybeans the year following use of this tank mixture on furrow irrigated milo.

APPLICATION RATES: Preplant Incorporated

BROADCAST RATE PER ACRE (QUARTS)*

Less than 1.5%
organic matter

1.5% or more
organic matter

**SOIL
TEXTURE**

Lasso +

atrazine 4L**

Lasso +

atrazine 4L**

Medium
Fine

1.5 to 2
1.75 to 2

1 to 1.25
1 to 1.5

1.75 to 2
2 to 2.5

1 to 1.5
1.25 to 1.75

45/68

*Use the higher rates in the recommended ranges for fields under irrigation.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

NOTE: Use 3 to 4 quarts of this product per acre for control of yellow nutsedge or to reduce competition from the hard-to-control weeds listed on this label.

APPLICATION RATES: Preemergence Surface

BROADCAST RATE PER ACRE (QUARTS)*

Less than 1.5% organic matter	1.5% or more organic matter
-------------------------------	-----------------------------

SOIL TEXTURE	Lasso + atrazine 4L**	Lasso + atrazine 4L**
Coarse (except sand)	1.5 to 1.75	0.75 to 1
Medium	1.5 to 2	1 to 1.25
Fine	1.5 to 2	1 to 1.5

*Use the higher rates in the recommended ranges for fields under irrigation.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

NOTE: When applied under sprinkler irrigation systems use a minimum of 2 quarts per acre of this product and 1 quart per acre of atrazine 4L in this tank mixture on coarse soils.

In the Texas Panhandle, Oklahoma Panhandle and the fine-textured soils of the Gulf Coast and Blacklands of Texas, do not exceed 1.25 quarts of atrazine 4L per acre as crop injury may result due to atrazine.

GRAIN SORGHUM (MILO)—CONSERVATION OR MINIMUM TILLAGE

NOTE: Important information regarding Conservation or Minimum Tillage appears in the "APPLICATION INFORMATION" section at the front of this label and should be carefully reviewed in conjunction with the information in this section.

Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

46/68

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the "GENERAL INFORMATION", the specific information should control.

Lasso and the following tank mixtures can be tank mixed with Roundup brands labeled for agricultural uses or Gramoxine Extra.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom

CONTROL OR SUPPRESSION OF EMERGED WEEDS - BURNDOWN PRIOR TO CROP

ROUNDUP UltraMAX

Annual Weeds-Apply 13 to 20 ounces of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, in these tank mixtures when weeds are less than 6 inches tall. Use 26 ounces of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, for control of fall panicum, barnyardgrass, crabgrass and shattercane up to 2 inches tall; Pennsylvania smartweed up to 6 inches tall; other annual weeds 6 to 12 inches tall.

Perennial Weeds-At normal application dates in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. Use of 1.6 to 3.25 quarts of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, in the above mixtures under these conditions provides top kill and reduces competition from many emerged perennial grasses and broadleaf weeds.

NOTE: When using these tank mixtures, do not exceed 3.25 quarts of Roundup UltraMAX per acre or equivalent rates of other Roundup brands labeled for agricultural uses.

GRAMOXONE EXTRA

Broadcast Treatment-Apply 1.5 to 3 pints of Gramoxone Extra per acre in these tank mixtures immediately before, during or after planting. Use 1.5 to 2 pints when weeds are 1 to 3 inches tall, 2 to 2.5 pints when weeds are 3 to 6 inches tall and 2.5 to 3 pints when weeds are 6 inches tall. These mixtures may not control weeds taller than 6 inches. Add a nonionic spreader surfactant (approved for use on crops) containing at least 75 percent surfactant active agent at 8 fluid ounces per 100 gallons of diluted spray.

PREEMERGENCE WEED CONTROL

LASSO

Applications must be made ONLY to grain sorghum planted with seed that has been properly treated with a seed protectant or safener.

APPLICATION RATES:

BROADCAST RATE PER ACRE (QUARTS)

47/68

SOIL TEXTURE	Lasso
Coarse	2 to 2.5
Medium	2.5 to 2.75
Fine	2.5 to 3

Use the higher rate of in the recommended ranges when organic matter content is 3 percent or more.

LASSO plus ATRAZINE

Applications must be made ONLY to grain sorghum planted with seed that has been properly treated with a seed protectant or safener.

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)	
	Lasso	+ atrazine 4L*
Coarse-sandy loam only	1.5 to 2	0.75 to 1
Medium	1.75 to 2.25	1 to 1.25
Fine	2 to 2.5	1.25 to 1.5

*Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

Use the higher rate of this product in the recommended ranges when fall panicum or crabgrass will be present.

Use the higher rate of atrazine in the recommended ranges on soils with greater than 3 percent organic matter.

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS can be planted the year following use of this mixture.

There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

EARLY PREPLANT APPLICATIONS

Apply this tank mixture in water or sprayable fluid fertilizer as a single application within 30 days prior to planting. When making split applications, apply this tank mixture within 45 days prior to planting, followed by an additional application of this product at planting. If emerged weeds are present at the time of the early preplant application or at-planting application, a labeled contact herbicide or tillage should be used. The at-planting application should be made before crop or weeds emerge. Minimize movement of treated

48/68

soil out of the row or untreated soil to the surface during planting or weed control may be reduced. Do not use in coarse or fine-textured soils.

Applications must be made ONLY to grain sorghum planted with seed that has been properly treated with a seed protectant or safener.

APPLICATION RATES: Early Preplant

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)		
	Early application		At-planting application
	Lasso	+ atrazine 4L	Lasso
SINGLE APPLICATION			
Coarse (except sand)	2.25 to 3	1 to 1.25	•
Medium	2.75 to 3	1.5 to 2	•
SPLIT APPLICATION			
Coarse (except sand)	1.75 to 2	1 to 1.5	0.75 to 1
Medium	1.75 to 2	1.5 to 2	0.75 to 1

SOYBEANS—CONVENTIONAL TILLAGE

NOTE: Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the GENERAL INFORMATION, the specific information should control.

When making applications on soybeans, do not make more than one application per year or exceed a total of 3 quarts of this product per acre per year.

Do not feed forage, hay or straw from soybeans treated with this product or tank mixtures with this product. Do not ensile soybeans treated with this product.

LASSO

Apply this product in water or sprayable fluid fertilizer solution.

APPROVED APPLICATION SYSTEMS

49/68

Ground-Broadcast boom, banded; Aerial-Fixed wing, helicopter; Center Pivot Irrigation;
Dry Bulk Fertilizer Impregnation

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface .

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)*	
	Less than 3% organic matter	3% or more organic matter
Coarse	2.5	2.5
Medium	2.5 to 3	2.5 to 3
Fine	2.5 to 3	3

* Use a minimum of 3 quarts per acre on coarse textured soils for control of yellow nutsedge or red rice.

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)*	
	Less than 3% organic matter	3% or more organic matter
Coarse**	2 to 2.25	2 to 2.25
Medium	2 to 2.75	2 to 2.75
Fine	2 to 2.75	2.5 to 3

* Use a minimum of 2.5 quarts of this product per acre on coarse soils and 3 quarts on medium or fine textured soils to control black or hairy nightshade.

**When applied through center pivot irrigation or under sprinkler irrigation systems on coarse soils, use 3 quarts per acre of this product.

**LASSO plus CANOPY
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.
DO NOT plant rice or peanuts within 18 months after application of this tank mixture.

APPROVED APPLICATION SYSTEM

Ground: Broadcast boom

50/68

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

Preplant Incorporated- Preplant incorporation may cause slight soybean injury early, but soybeans should recover with no subsequent effect on growth or yield.

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE*	BROADCAST RATE PER ACRE**			
	1/2% to 3% organic matter		3% to 5% organic matter	
	Lasso (quarts)	Canopy (ounces)	Lasso (quarts)	Canopy (ounces)
Coarse-loamy sand, sandy loam only	2 to 2.5	6	2 to 2.5	8
Medium***	2 to 2.5	8	2.25 to 2.75	10
Fine	2.25 to 2.75	10	2.5 to 3	12

*DO NOT APPLY TO SOIL WITH LESS THAN 1/2 percent organic matter.

**Use the higher rate of this product in the recommended ranges when broadleaf signalgrass is present.

***In southeast KS, southwest MO and northeast OK on soils with 1/2 to 3 percent organic matter, use 5 to 6 ounces Canopy per acre. On silt loam soils in TN and KY, use 6 to 8 ounces of Canopy per acre.

NOTE: In the states of IA, IL, IN, KS, MI, MO, MN, NE, OK, OH, and WI on soils with 1/2 to 4 percent organic matter, reduced rates of Canopy are recommended for broadleaf weed control. Use 5 ounces of Canopy per acre on coarse soils (loamy sand and sandy loam only); 6 ounces on medium soils; 6 to 7 ounces on fine soils. Do not apply to soils with greater than 6.8 pH or to soils with less than 1/2 percent organic matter or greater than 4 percent organic matter.

For improved control of heavy velvetleaf infestation, add 1/2 pint of Command herbicide per acre to this tank mixture and preplant incorporate.

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE*	BROADCAST RATE PER ACRE**			
	1/2% to 3% organic matter		3% to 5% organic matter	
	Lasso (quarts)	Canopy (ounces)	Lasso (quarts)	Canopy (ounces)
Coarse-loamy sand, sandy loam only	1.75 to 2.25	6	1.75 to 2.25	8
Medium***	1.75 to 2.25	8	2 to 2.5	10

51/68

Fine 2 to 2.5 10 2.25 to 2.75 12

* DO NOT APPLY TO SOIL WITH LESS THAN 1/2 percent organic matter.

**Use the higher rate of this product in the recommended ranges when broadleaf signalgrass is present.

***In southeast KS, southwest MO and northeast OK on soils with 1/2 to 3 percent organic matter, use 5 to 6 ounces Canopy per acre. On silt loam soils in TN and KY, use 6 to 8 ounces of Canopy per acre.

NOTE: In the states of IA, IL, IN, KS, MI, MO, MN, NE, OK, OH, and WI on soils with 1/2 to 4 percent organic matter, reduced rates of Canopy are recommended for broadleaf weed control. Use 5 ounces of Canopy per acre on coarse soils (loamy sand and sandy loam only); 6 ounces on medium soils; 6 to 7 ounces on fine soils. Do not apply to soils with greater than 6.8 pH or to soils with less than 1/2 percent organic matter or greater than 4 percent organic matter.

**LASSO plus COMMAND
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.
APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom, banded

APPROVED APPLICATION METHODS

Preplant Incorporated

APPLICATION RATES: Apply 2 to 2.5 quarts of this product plus 1 pint of Command per acre. In areas of heavy pigweed and lambsquarter infestation, use 2.5 to 3 quarts of this product per acre in the tank mixture.

**LASSO plus SENCOR
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.

NOTE: DO NOT use on muck soils or on alkaline soils with greater than 7.4 pH. Crop injury may result if atrazine was applied the year preceding use of this tank mixture.

DO NOT REPLANT CROPS OTHER THAN SOYBEANS FOR 120 DAYS AFTER APPLICATION OF THESE TANK MIXTURES.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Dry Bulk Fertilizer Impregnation

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

52/68

Preplant Incorporated- Preplant incorporation may cause slight soybean injury early, but soybeans should recover with no subsequent effect on growth or yield.

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE	BROADCAST RATE PER ACRE**			
	Less than 3% organic matter		3% or more organic matter	
	Lasso + (quarts)	Sencor 4F* (pint)	Lasso + (quarts)	Sencor 4F** (pint)
Coarse-sandy loam only	2 to 2.5	0.5	2 to 2.5	0.75
Medium	2 to 2.5	0.5 to 0.75	2.25 to 2.75	0.75
Fine	2.25 to 2.75	0.75 to 1	2.5 to 3	1

*Use the rates listed in this label when using Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

**Use the 3 quart per acre rate of this product in areas of heavy weed infestation. Use 3 quarts of this product when black or hairy nightshade, yellow nutsedge or red rice is present.

NOTE: Do not use on coarse textured soils. Do not use on medium or fine textured soils with less than 1 percent organic matter. These mixtures, preplant incorporated, may cause slight soybean injury early, but the soybeans should recover and subsequent growth or yield should not be affected.

For improved control of heavy velvetleaf infestation, add 1/2 pint of Command per acre to this tank mixture and preplant incorporate.

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE	BROADCAST RATE PER ACRE			
	Less than 3% organic matter		3% or more organic matter	
	Lasso + (quarts)	Sencor 4F* (pint)	Lasso + (quarts)	Sencor 4F* (pint)
Coarse-sandy loam only	1.75 to 2.25	0.5	1.75 to 2.25	0.75
Medium	1.75 to 2.25	0.5 to 0.75	2 to 2.5	0.75
Fine	2 to 2.5	0.75 to 1	2.25 to 2.75	1

*Use the rates listed in this label when using or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

NOTE: DO NOT use on sand or loamy sand soil. Use 1.5 pints of or Sencor 4F per acre in this tank mixture when applying on heavy clay soil of the Mississippi Delta. Use a minimum of 2.5 quarts of this product on coarse soils and 3 quarts in this tank mixture on medium or fine-textured soils to control black or hairy nightshade.

**LASSO plus LINURON
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.

NOTE: Plant seed at least 1-3/4 inches deep and DO NOT use on sand or loamy sand or on soils with less than 1 percent organic matter as crop injury may occur. DO NOT spray over the top of emerged soybeans.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded

APPROVED APPLICATION METHODS

Preemergence Surface

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)			
	Less than 3% organic matter		3% or more organic matter	
	Lasso +	linuron 4L**	Lasso +	linuron 4L**
Coarse-sandy loam only	1.75 to 2.25	0.5	1.75 to 2.25	0.5 to 0.75
Medium	1.75 to 2.25	0.5 to 0.75	2 to 2.5	0.75 to 1
Fine	2 to 2.5	1	2.25 to 2.75	1.25 to 1.5

**Use the rates listed on this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50% dry flowable formulations. One quart of linuron 4L equals 2 pounds of linuron 50WP or 2 pounds of linuron 50% dry flowable.

**LASSO plus LOROX PLUS
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded

54/68

APPROVED APPLICATION METHODS

Preemergence Surface

APPLICATION RATES: Preemergence Surface

SOIL TEXTURE	BROADCAST RATE PER ACRE		
	1/2% to 3% organic matter		
	Lasso (quarts)	+	Lorox Plus (ounces)
Coarse-sandy loam only	1.75 to 2.25		12 to 14
Medium	1.75 to 2.25		14 to 16
Fine	2 to 2.5		16 to 18

**Use the higher rate of this product in the recommended ranges when broadleaf signalgrass is present. Use the higher rate of Lorox Plus in the recommended ranges on soils with higher organic matter or heavy pressure from large, deep germinating weed seeds.

LASSO plus PURSUIT (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE	BROADCAST RATE PER ACRE*		
	Less than 3% organic matter	3% or more organic matter	
	Lasso (quarts)	Lasso + Pursuit (quarts)	Pursuit (ounces)
Coarse-sandy loam only	1.5 to 2	1.5 to 2	4
Medium	1.5 to 2	1.75 to 2.25	4

Lasso 2001-1-Sep1701

55/68

Fine	1.75 to 2.25	2 to 2.5	4
------	--------------	----------	---

*Use the higher rate of this product in the recommended ranges when the following weeds are present: fall panicum, barnyardgrass, broadleaf signalgrass, seedling johnsongrass, red rice or yellow nutsedge.

NOTE: Preplant incorporate this tank mixture for more consistent control of common cocklebur, common lambsquarters, jimsonweed, prickly sida, velvetleaf and sunflower and for reduced competition of morningglory and giant ragweed.

**LASSO plus SCEPTER
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE*		
	Less than 3% organic matter		3% or more organic matter
	Lasso (quarts)	Lasso (quarts)	Scepter** (pints)
Coarse	1.5 to 2	1.5 to 2	1/2 to 2/3
Medium	1.5 to 2	1.75 to 2.25	1/2 to 2/3
Fine	1.75 to 2.25	2 to 2.5	1/2 to 2/3

*Use the higher rate of this product in the recommended ranges when the following weeds are present: fall panicum, barnyardgrass, broadleaf signalgrass, seedling johnsongrass, red rice or yellow nutsedge.

**Use the higher rate of Scepter in the recommended range when red rice, sicklepod, beggarweed or velvetleaf are present.

For improved control of heavy velvetleaf infestation, add 1/2 pint of Command herbicide per acre to this tank mixture and preplant incorporate.

**LASSO plus TREFLAN
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded

Lasso 2001-1-Sep1701

5/6/68

APPROVED APPLICATION METHODS

Preplant Incorporated-Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil within eight (8) hours after application.

APPLICATION RATES: Apply 2.5 to 3 quarts of this product plus 0.5 quart of Treflan 4EC per acre.

SOYBEANS—CONSERVATION OR MINIMUM TILLAGE

NOTE: Important information regarding Conservation or Minimum Tillage appears in the Application Information section at the front of this label and should be carefully reviewed in conjunction with the information in this section

Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the GENERAL INFORMATION, the specific information should control.

APPROVED APPLICATION SYSTEMS

Ground: —Broadcast boom

CONTROL OR SUPPRESSION OF EMERGED WEEDS – BURNDOWN PRIOR TO CROP

ROUNDUP UltraMAX

Annual Weeds-Apply 13 to 20 ounces of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, in these tank mixtures when weeds are less than 6 inches tall. Use 26 ounces of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, for control of fall panicum, barnyardgrass, crabgrass and shattercane up to 2 inches tall; Pennsylvania smartweed up to 6 inches tall; other annual weeds 6 to 12 inches tall.

Perennial Weeds-At normal application dates in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. Use of 1.6 to 3.25 quarts of Roundup UltraMAX per acre, or equivalent rates of other Roundup brands labeled for agricultural uses, in these tank mixtures under these conditions provides top kill and reduces competition from many perennial grasses and broadleaf weeds. **NOTE:** When using these tank mixtures, do not exceed 3.25 quarts of Roundup UltraMAX per acre or equivalent rates of other Roundup brands labeled for agricultural uses.

57/68

GRAMOXONE EXTRA

Broadcast Treatment-Apply 1.5 to 3 pints of Gramoxone Extra in this tank mixture immediately before, during or after planting. Use 1.5 to 2 pints when weeds are 1 to 3 inches tall, 2 to 2.5 pints when weeds are 3 to 6 inches tall and 2.5 to 3 pints when weeds are 6 inches tall. This mixture may not control weeds taller than 6 inches. Add a nonionic spreader surfactant (approved for use in crops) containing at least 75 percent surface active agent at 8 fluid ounces per 100 gallons of diluted spray.

PREEMERGENCE WEED CONTROL

LASSO

APPLICATION RATES:

BROADCAST RATE PER ACRE (QUARTS)	
SOIL TEXTURE	Lasso
Coarse	2.5
Medium	3
Fine	3

LASSO plus CANOPY

APPLICATION RATES:

BROADCAST RATE PER ACRE		
SOIL TEXTURE	Lasso (quarts)	+ Canopy (ounces)
Coarse-sandy loam only	1.75 to 2.25	6
Medium	2 to 2.5	8
Fine	2.5 to 3	10

LASSO plus SENCOR

APPLICATION RATES:

BROADCAST RATE PER ACRE		
SOIL TEXTURE	Lasso (quarts)	+ or Sencor 4F* (pints)

Lasso 2001-1-Sep1701

58/68

Coarse-sandy loam only	1.75 to 2.25	0.5 to 0.75
Medium	2 to 2.5	0.75 to 1
Fine	2.5 to 3	1 to 1.5**

*Use the rates listed in this label when using or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

**On the heavy clay soil of the Mississippi Delta, use 1.5 pints of or Sencor 4F per acre.

Use the higher rate of this product in the recommended ranges when fall panicum or crabgrass will be present.

Use the higher rate of Sencor in the recommended ranges on soils with greater than 3 percent organic matter.

DO NOT use this mixture on sand or loamy sand soils as crop injury from Sencor may occur.

DO NOT use on muck soils.

DO NOT apply on alkaline soils with a pH greater than 7.4.

Crop injury may occur if atrazine was applied the year before use of this tank mixture.

DO NOT REPLANT CROPS OTHER THAN SOYBEANS FOR 120 DAYS AFTER APPLICATION OF THESE TANK MIXTURES.

LASSO plus LINURON

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)	
	Lasso	+ linuron 4L*
Coarse-sandy loam only	1.75 to 2.25	0.5 to 0.75
Medium	2 to 2.5	0.75 to 1
Fine	2.5 to 3	1 to 1.5

*Use the rates listed on this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50% dry flowable formulation. One quart of linuron 4L equals 2 pounds of linuron 50WP or 2 pounds of linuron 50% dry flowable.

Use the higher rate of this product in the recommended ranges when fall panicum or crabgrass will be present. Use the higher rate of linuron in the recommended ranges on soils with greater than 3 percent organic matter.

59/68

DO NOT use this tank mixture on sand or loamy sand or on soil with less than 1 percent organic matter as crop injury from linuron may occur.

LASSO plus LOROX PLUS

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE	
	Lasso (quarts)	Lorox Plus (ounces)
Coarse-sandy loam only	1.75 to 2.25	12 to 14
Medium	2 to 2.5	14 to 16
Fine	2.5 to 3	16 to 18

LASSO plus PURSUIT

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE	
	Lasso (quarts)	Pursuit (ounces)
Coarse	1.75 to 2.25	4
Medium	2 to 2.5	4
Fine	2.5 to 3	4

LASSO plus SCEPTER

APPLICATION RATES:

SOIL TEXTURE	BROADCAST RATE PER ACRE**	
	Lasso (quarts)	Scepter (pints)
Coarse	1.75 to 2.25	2/3
Medium	2 to 2.5	2/3
Fine	2.5 to 3	2/3

PEANUTS

60/68

NOTE: Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the GENERAL INFORMATION, the specific information should control.

When making applications on peanuts, do not make more than two applications per year or exceed a total of 4 quarts of this product per acre per year.

Do not feed forage or hay from treated crop.

LASSO

Apply this product in water.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Center Pivot Irrigation

APPROVED APPLICATION METHODS

Preplant Incorporated, Preemergence Surface, Ground-crack Surface

Preplant Incorporated*- For control of Texas panicum, this application should be made as a tank mixture with a labeled rate of Prowl or SonolanTM DF herbicide.

*Incorporation on coarse soils may result in reduced length of weed control due to leaching with rainfall or irrigation. For improved control of yellow nutsedge and labeled broadleaf weeds, a labeled sequential ground-crack herbicide application should be made.

APPLICATION RATES:

APPLICATION METHOD	LASSO (QUARTS)*
Preplant Incorporated	3 to 4
Preemergence or Ground-crack Surface	3 to 4

DRY BEANS

6/1/68

NOTE: Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the GENERAL INFORMATION, the specific information should control.

When making applications on dry beans, do not make more than one application per year or exceed a total of 3 quarts of this product per acre per year.

Do not feed forage or hay from treated crop.

Dry Beans (All Types)

FOR USE ONLY IN NEW YORK AND WEST OF THE MISSISSIPPI RIVER,
EXCEPT IN KERN COUNTY, CALIFORNIA.

LASSO

Apply this product in water.

NOTE: DO NOT APPLY ON DRY BEANS AFTER PLANTING AS CROP INJURY MAY OCCUR. This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom

APPROVED APPLICATION METHODS

Preplant Incorporated

APPLICATION RATES: Apply 2.5 to 3 quarts of this product per acre and shallowly incorporate prior to planting. Use the higher rate when black or hairy nightshade or yellow nutsedge are present.

**LASSO plus EPTAM
(TANK MIXTURE)
Dry Beans (All Types)**

FOR USE ONLY IN MICHIGAN AND
WEST OF THE MISSISSIPPI RIVER,
EXCEPT IN CALIFORNIA

Apply this tank mixture in water.

NOTE: Read the label for Eptam thoroughly for rotational crop restrictions. This tank mixture may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

APPROVED APPLICATION SYSTEMS

02/68

Ground: Broadcast boom

APPROVED APPLICATION METHODS

Preplant Incorporated

APPLICATION RATES: Apply 2 to 3 quarts of this product plus 2 to 3 pints of Eptam 7-E per acre. Use the higher recommended rate of this product for this tank mixture when black or hairy nightshade or yellow nutsedge are present. USE ONLY THE 2 QUART RATE OF LASSO IN MICHIGAN.

**LASSO plus TREFLAN
(TANK MIXTURE)**

DRY BEANS (ALL TYPES)

FOR USE ONLY WEST OF THE
MISSISSIPPI RIVER, EXCEPT IN KERN
COUNTY, CALIFORNIA.

Apply this tank mixture in water.

NOTE: This tank mixture may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom

APPROVED APPLICATION METHODS

Preplant Incorporated

APPLICATION RATES: Apply 2.5 to 3 quarts of this product plus 1/2 quart of Treflan per acre. Use the higher rate of this product for this tank mixture when black or hairy nightshade or yellow nutsedge are present.

RED KIDNEY BEAN TYPES ONLY

FOR USE IN ILLINOIS,
INDIANA AND WISCONSIN ONLY
LASSO

Apply this product in water.

NOTE: This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

Only one application of this product may be made per cropping season.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom, banded

APPROVED APPLICATION METHODS

Preplant Incorporated

63/68

APPLICATION RATES: Apply 2.5 to 3 quarts of this product per acre. Use the higher recommended rate when black or hairy nightshade or yellow nutsedge are present.

LIMA BEANS (GREEN)

NOTE: Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the GENERAL INFORMATION, the specific information should control.

When making applications on lima beans, do not make more than one application per year or exceed a total of 3 quarts of this product per acre per year.

Do not feed forage or hay from treated crop.

FOR USE IN ALL STATES, EXCEPT
IN KERN COUNTY, CALIFORNIA

LASSO

Apply this product in water.

NOTE: This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom

APPROVED APPLICATION METHODS

Preplant Incorporated

APPLICATION RATES: Preplant Incorporated

SOIL TEXTURE	BROADCAST RATE PER ACRE (QUARTS)
	LASSO
Coarse	2
Medium and Fine**	2.5 to 3

**Use the higher rate in the recommended range when black or hairy nightshade or yellow nutsedge are present.

64/68

WOODY ORNAMENTALS

NOTE: Each crop section of this label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appears in the "GENERAL INFORMATION" section and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the GENERAL INFORMATION, the specific information should control.

LASSO

FOR USE ON JUNIPER (JUNIPERUS SPP.) AND YEW (TAXUS SPP.)

Apply this product in water.

NOTE: Topical application or contact of this product with green foliage of the woody ornamental may cause injury and is not recommended. In case green foliage is accidentally sprayed, overhead irrigation within 1 hour after application will minimize injury symptoms (i.e., leaf tip burn, discoloration, distortion, etc.). This product is recommended for this use for application by certified, professional applicators only.

DO NOT USE THIS PRODUCT ON SEEDBEDS, UNROOTED CUTTINGS OR PRIOR TO TRANSPLANTING. DO NOT USE THIS PRODUCT IN GREENHOUSES OR OTHER ENCLOSED STRUCTURES, APPLIED THROUGH IRRIGATION SYSTEMS OR MIXED WITH SPRAYABLE FLUID FERTILIZERS. APPLICATIONS DURING PERIODS WHEN TEMPERATURE EXCEEDS 90°F MAY CAUSE INJURY TO ORNAMENTALS.

APPROVED APPLICATION SYSTEMS

Ground: Broadcast boom banded

APPROVED APPLICATION METHODS

Post-Transplant Directed-Apply this product as a directed spray after transplanting or to established plantings prior to weed emergence.

APPLICATION RATE: Apply 4 quarts of this product per acre as a broadcast treatment.

DO NOT exceed a total of 4 quarts of Lasso per year.

APPENDIX

LIST OF COMMON AND SCIENTIFIC NAMES OF ANNUAL GRASSES AND BROADLEAF WEEDS.

ANNUAL GRASSES

Barnyardgrass

Echinochloa crus-galli

Crabgrass

Digitaria ischaemum

Digitaria sanguinalis

Cupgrass, woolly

65/68

Eriochloa villosa

Foxtail, giant
Setaria faberi

**Foxtail; green, robust
purple, robust white**
Setaria viridis

Foxtail, yellow
Setaria lutescens

Goosegrass
Eleusine indica

Johnsongrass, seedling
Sorghum halepense

Millet, proso
Panicum miliaceum

Panicum, browntop
Panicum fasciculatum

Panicum, fall
Panicum dichotomiflorum

Panicum, Texas
Panicum texanum

Rice, red
Oryza sativa

Sandbur, Grassbur
Cenchrus incertus

Shattercane, wild cane
Sorghum bicolor

Signalgrass, broadleaf
Brachiaria platyphylla

Sprangletop, red
Leptochloa filiformis

Witchgrass
Panicum capillare

ANNUAL BROADLEAVES

Beggarweed, Florida
Desmodium tortuosum

Carpetweed

Lasso 2001-1-Sep1701

66/68

Mollugo verticillata

Cocklebur

Xanthium strumarium

Coffeeweed,

Hemp sesbania
Sesbania exaltata

Copperleaf, hophornbeam

Acalypha spp.

Galinsoga

Galinsoga spp.

Groundcherry, annual

Physalis spp.

Groundcherry, cutleaf

Physalis angulata

Jimsonweed

Datura stramonium

Kochia

Kochia scoparia

Lambsquarters

Chenopodium album

Morningglory:

Tall

Ipomoea purpurea

Pitted

Ipomoea lacunosa

Ivyleaf

Ipomoea hederacea

Entireleaf

Ipomoea hederacea
var. integruscula

Smallflower

Jacquemontia tamnifolia

Mustard

Brassica spp.

Nightshade, black

Solanum nigrum

Nightshade, hairy

Solanum sarrachoides

Pigweed, Carelessweed

Amaranthus spp.

67/68

Purslane
Portulaca oleracea

Pusley, Florida
Richardia scabra

Ragweed, common
Ambrosia artemisiifolia

Ragweed, giant
Ambrosia trifida

Sicklepod
Cassia obtusifolia

Sida, prickly; Teaweed
Sida spinosa

Smartweed
Polygonum pensylvanicum
Polygonum persicaria

Starbur, bristly
Acanthospermum hispidum

Sunflower, common
Helianthus annuus

Velvetleaf, Buttonweed
Abutilon theophrasti

Waterhemp
Amaranthus tuberculatos

SEDGE

Nutsedge, yellow
Cyperus esculentus

®Lasso, Roundup, Roundup Ready, and Roundup UltraMAX are registered trademarks of Monsanto Technology LLC.

™Agsorb is a trademark of Oil-Dri Corporation.

™Marksman, Prowl, Pursuit and Scepter are trademarks of BASF Corporation.

™Canopy is a trademark of E. I. duPont de Nemours and Company.

™Command is a trademark of FMC Corporation.

™Eptam, Gramoxone and Princep are trademarks of Syngenta Group Company.

™Microcel is a trademark of Celite Corporation.

™MMP-79 is a trademark of Eagle Picher Industries, Inc.

68/168

TMSencor is a trademark of the parent company of Farbenfabriken Bayer GmbH.

TMTreflan and Sonalan are trademarks of Dow AgroSciences LLC. TMBanvel is the trademark of Micro Flo Company LLC.

TMLorox is the trademark of Griffin LLC

TMBalance Pro is a trademark of the Aventis Group.

CALIFORNIA

This product has been approved by the U.S. Environmental Protection Agency for the uses, crops and sites listed in this label. Approval of the items listed below is pending under the State of California registration requirements.

These use conditions, crops and sites may not be treated with this product in California until approval is received:

- Aerial applications.
- Lasso plus atrazine in early preplant applications on grain sorghum (milo).
- Tank mixtures with Canopy, Command, Lorox Plus and Scepter in soybeans.
- Tank mixtures with Banvel, Marksman, Prowl, and atrazine 90DF in corn.
- Tank mixtures with Gramoxone Extra in grain sorghum (milo).
- Control of *Ipomoea hederacea*.
- Control of *Ipomoea hederacea* var.
- Control of *Ipomoea integruscula*.
- Control of *Ipomoea lacunosa*.
- Control of *Jacquemontia tamnifolia*.
- Control of *Leptochloa filiformis*.
- Control of *Panicum fasciculatum*.

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000.

©2002 MONSANTO COMPANY
St. Louis, Missouri, 63167 U.S.A.