

524-314

06-01-2004

1/7

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval Expires 2-28-95

	United States <b>Environmental Protection Agency</b> Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number <b>296801</b>
--	---	--	--

**Application for Pesticide - Section I**

1. Company/Product Number <b>524-314</b>	2. EPA Product Manager <b>James A. Tompkins</b>	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) <b>Intrro Preemergent herbicide</b>	PM # <b>25</b>	
5. Name and Address of Applicant (Include ZIP Code) <b>Monsanto Company 600 13<sup>th</sup> St., N.W., Suite 660 Washington, DC 20005</b> <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name <b>JUN - 1 2004</b>	

**NOTIFICATION**

**Section - II**

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter
<input type="checkbox"/> Resubmission in response to Agency letter dated	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)  
Notification of additional uses on field and silage corn in the states of Hawaii and California via supplemental labeling.

**Section - III**

Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per Container	If "Yes" Package wgt. No. per Container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name <b>Russell P. Schneider Ph.D.</b>	Title <b>Director, Regulatory Affairs</b>	Telephone No. (Include Area Code) <b>(202) 383-2866</b>
Certification <i>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</i>		6. Date Application Received (Stamped)
2. Signature 	3. Title <b>Registration Manager</b>	
4. Printed Name <b>Dawn M. Fee-White</b>	5. Date <b>4 May 2004</b>	

**RESTRICTED USE PESTICIDE**  
due to groundwater concerns. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

**SUPPLEMENTAL LABELING**

READ THE ENTIRE LABEL FOR INTRRO™ HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING. FOLLOW ALL APPLICATION DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL.

"Label" as used in the supplemental labeling refers to the label booklet for INTRRO herbicide and this supplement.

**NOTIFICATION**  
JUN - 1 2004



EPA Reg. No. 524-314

INTRRO is a trademark of Monsanto Technology LLC.

**FOR USE IN FIELD CORN AND SILAGE CORN IN HAWAII ONLY**

Keep out of reach of children.

**DANGER! PELIGRO!**

The use of this product may be hazardous to your health. This product contains alachlor, which has been determined to cause tumors in laboratory animals.  
In case of an emergency involving this product or for user safety information on this product, Call Collect, day or night, (314) 694-4000.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

**CORN CONTAINING THE ROUNDUP READY® GENE**

This product may be used in Roundup Ready corn, or corn that contains the Roundup Ready gene. A minimum of 1.25 quarts per acre of this product is required. Use the full labeled rate of this product when the application is 14 days or more prior to planting or when the following grasses may be expected: barnyardgrass, shattercane, seedling johnsongrass or sandbur.

**Preemergence Sequential Program**

This product may be used at a reduced rate preemergence in a sequential program with Roundup UltraMAX® herbicide, or other Roundup agricultural herbicides labeled for use in Roundup Ready corn. Use 50% to 100% of the labeled rate of this product for the appropriate soil texture and organic matter, per the Application Rate Table. Apply a sequential application of Roundup UltraMAX, or other Roundup agricultural herbicides labeled for use in Roundup Ready corn, per the label instructions.

If weeds are present at the time of preemergence application, tank-mix Roundup UltraMAX or other Roundup agricultural herbicides, to control emerged vegetation.

**CORN—CONVENTIONAL TILLAGE (All Types)**

**NOTE:** This label provides application rates for this product and tank mixtures including this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appear in the "GENERAL INFORMATION" section of the INTRRO label booklet and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the "GENERAL INFORMATION", this specific information should control.

When making applications on corn, do not make more than two applications per year or exceed a total of 4 quarts of this product per acre per year. Do not make more than one

application or exceed 4 quarts of this product per acre per year on sweet corn.

**INTRRO**

Apply this product in water or sprayable fluid fertilizer solution.

**APPROVED APPLICATION SYSTEMS**

**Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Center Pivot Irrigation; Dry Bulk Fertilizer Impregnation**

**APPROVED APPLICATION METHODS**

**Preplant Incorporated, Preemergence Surface**

**Sequential Treatment**—This product may be applied as two sequential treatments for improved control of hard-to-control weeds. The combined total application rate for the two treatments must not exceed 4 quarts per acre per year. The second treatment must be applied before the corn emerges.

**APPLICATION RATES: Preplant Incorporated**

SOIL TEXTURE	BROADCAST RATE PER ACRE	
	INTRRO (quarts)*	
	Less than 3% organic matter	3% or more organic matter
Coarse	2.5	2.5
Medium	2.5 to 3	2.5 to 3
Fine	2.5 to 3	3 to 3.5

\* Use 2.5 to 3.5 quarts per acre where furrow irrigation is used. Use a minimum of 3 quarts on coarse-textured soils for control of yellow nutsedge or red rice. Rates may be increased to a maximum of 4 quarts on any soil type when heavy infestation of yellow nutsedge or red rice is present.

On muck or peat or coarse soils with 10 percent or more organic matter use 4 quarts of this product per acre.

**APPLICATION RATES: Preemergence Surface**

SOIL TEXTURE	BROADCAST RATE PER ACRE	
	INTRRO (quarts)*	
	Less than 3% organic matter	3% or more organic matter
Coarse**	2 to 2.25	2 to 2.25
Medium	2 to 2.75	2 to 2.75
Fine	2 to 2.75	2.5 to 3.25

\* Use a minimum of 2.5 quarts of this product per acre on coarse soils and 3 to 4 quarts on medium- or fine-textured soils to control black or hairy nightshade.

\*\* When applied through center pivot irrigation or under sprinkler irrigation systems on coarse soils, use a minimum of 3 quarts per acre of this product.

On muck or peat soils or on coarse soils with 10 percent or more organic matter use 4 quarts of this product per acre.

**INTRRO plus ATRAZINE (TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

**APPROVED APPLICATION SYSTEMS**

**Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Dry Bulk Fertilizer Impregnation**

**APPROVED APPLICATION METHODS Preplant Incorporated, Preemergence Surface**

**NOTE:** CORN, PEANUTS, SORGHUM OR SOYBEANS\* can be planted the year following use of this mixture.

\* There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow-irrigated corn.

When applied following INTRRO or tank mixtures of INTRRO, the total amount of this product applied alone or in combination must not exceed 4 quarts per acre per year. The total amount of atrazine 4L must not exceed 3 quarts per acre per year.

**APPLICATION RATES: Preplant Incorporated**

SOIL TEXTURE	BROADCAST RATE PER ACRE			
	Less than 3% organic matter		3% or more organic matter	
	INTRRO + atrazine 4L* (quarts)	INTRRO + atrazine 4L* (quarts)	INTRRO + atrazine 4L* (quarts)	INTRRO + atrazine 4L* (quarts)
Coarse	1.75 to 2.25	1.25 to 2	1.75 to 2.25	1.25 to 2
Medium	1.75 to 2.25	1.25 to 2	2.25 to 2.5	1.5 to 2
Fine	2.25 to 2.5	1.5 to 2	2.25 to 2.75	1.5 to 2

\* Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

**APPLICATION RATES: Preemergence Surface**

SOIL TEXTURE	BROADCAST RATE PER ACRE			
	Less than 3% organic matter		3% or more organic matter	
	INTRRO + atrazine 4L* (quarts)	INTRRO + atrazine 4L* (quarts)	INTRRO + atrazine 4L* (quarts)	INTRRO + atrazine 4L* (quarts)
Coarse	1.5 to 2	0.75 to 2	1.5 to 2	1 to 2
Medium	1.5 to 2	1 to 2	2 to 2.25	1.25 to 2
Fine	2 to 2.25	1.25 to 2	2 to 2.75	1.25 to 2

\* Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

**NOTE:** When applied under sprinkler irrigation systems on coarse soils, use 1 quart per acre of atrazine 4L and a minimum of 2 quarts per acre of this product in this tank mixture.

**INTRRO plus PRINCEP® (TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution.

**APPROVED APPLICATION SYSTEMS**

**Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter; Dry Bulk Fertilizer Impregnation**

**APPROVED APPLICATION METHODS**

**Preplant Incorporated, Preemergence Surface**

**NOTE:** LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP, PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS.

**APPLICATION RATES: Preplant Incorporated**

SOIL TEXTURE	BROADCAST RATE PER ACRE			
	1% to 3% organic matter		3% or more organic matter	
	INTRRO (quarts)	+ Princep 4L* (quarts)	INTRRO (quarts)	+ Princep 4L* (quarts)
Coarse-sandy loam only	1.75 to 2.25	1.25 to 2	1.75 to 2.25	1.25 to 2
Medium	1.75 to 2.25	1.25 to 2	2.25 to 2.5	1.5 to 2
Fine	2.25 to 2.5	1.5 to 2	2.25 to 2.75	1.5 to 2

\* Use rates listed in this label when using Princep 4L. Use equivalent rates when using Princep 80W or 90% dry flowable formulations. One quart of the 4L formulation equals 1.25 pounds of the 80W and 1.1 pounds of the 90% dry flowable formulations.

**APPLICATION RATES: Preemergence Surface**

SOIL TEXTURE	BROADCAST RATE PER ACRE			
	1% to 3% organic matter		3% or more organic matter	
	INTRRO (quarts)	+ Princep 4L* (quarts)	INTRRO (quarts)	+ Princep 4L* (quarts)
Coarse-sandy loam only	1.5 to 2	0.75 to 2	1.5 to 2	1 to 2
Medium	1.5 to 2	1 to 2	2 to 2.25	1.25 to 2
Fine	2 to 2.25	1.25 to 2	2 to 2.75	1.25 to 2

\* Use rates listed in this label when using Princep 4L. Use equivalent rates when using Princep 80W or 90% dry flowable formulations. One quart of the 4L formulation equals 1.25 pounds of the 80W formulation and 1.1 pounds of the 90% dry flowable formulations.

**INTRRO plus ATRAZINE plus PROWL® (TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of triazine-resistant lambsquarters and pigweed.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

**APPROVED APPLICATION SYSTEMS**

**Ground: Broadcast boom, banded; Aerial: Fixed wing, helicopter**

**APPROVED APPLICATION METHOD**

**Preemergence Surface**

DO NOT preplant incorporate this tank mixture as serious crop injury can result.

**NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS\*** can be planted the year following use of this mixture.

\* There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow-irrigated corn.

**APPLICATION RATES:**

SOIL TEXTURE	BROADCAST RATE PER ACRE		
	Less than 3% organic matter		
	INTRRO (quarts)	+ atrazine 4L* (quarts)	+ Prowl (quarts)
Coarse	1.5 to 2	0.75 to 2	0.75
Medium	1.5 to 2	1 to 2	0.75 to 1
Fine	2 to 2.25	1.25 to 2	1

**BROADCAST RATE PER ACRE**

SOIL TEXTURE	3% or more organic matter			
	INTRRO (quarts)	+	atrazine 4L* (quarts)	+ Prowl (quarts)
	Coarse	1.5 to 2		1 to 2
Medium	2 to 2.25		1.25 to 2	1
Fine	2 to 2.75		1.25 to 2	1

\* Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

**CORN-CONSERVATION OR MINIMUM TILLAGE (All Types)**

Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control, injury to crops, persons or animals, or other unintended consequences. Refer to specific product labels for crop rotation restrictions and cautionary statements of all products used in these tank mixtures, including precautions on soil pH, sensitive varieties, minimum re-cropping interval and rotational guidelines.

Use the higher rates in the recommended ranges in areas of heavy weed infestation or where otherwise specified. If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds. Do not apply when conditions favor drift.

Detailed information regarding "APPLICATION SYSTEMS" and "APPLICATION TIMINGS AND METHODS" appear in the "GENERAL INFORMATION" section of the INTRRO label booklet and should be carefully reviewed in conjunction with the information in this section. If the specific information in this section differs from the "GENERAL INFORMATION", this specific information should control.

This product and the following tank mixtures can be tank mixed with Roundup agricultural herbicides, Gramoxone Extra and/or 2,4-D.

**APPROVED APPLICATION SYSTEMS**

**Ground: Broadcast boom**

**Note:** For application into CONSERVATION OR MINIMUM TILLAGE SYSTEMS, apply in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre when adding a Roundup agricultural herbicide or 2,4-D to the mix. Apply in 20 to 60 gallons of water or nitrogen solution when adding Gramoxone herbicide to the mix. Do not apply during periods of gusty winds, when winds are in excess of 15 mph or when other conditions favoring drift exist.

**ROUNDUP UltraMAX**

**Annual Weeds**—Apply 13 to 20 ounces of Roundup UltraMAX per acre, or equivalent rates of other Roundup agricultural herbicides, in these tank mixtures when weeds are less than 6 inches tall. Use 26 ounces of Roundup UltraMAX per acre, or equivalent rates of other Roundup agricultural herbicides, for control of fall panicum, barnyardgrass, crabgrass and shattercane up to 2 inches tall; Pennsylvania smartweed up to 6 inches tall; other annual weeds 6 to 12 inches tall.

**Perennial Weeds**—At normal application dates in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. Use of 1.6 to 3.25 quarts of Roundup UltraMAX per acre, or equivalent rates of other Roundup agricultural herbicides, in the above mixtures under these conditions provides top kill and reduces competition from many emerged perennial grasses and broadleaf weeds.

**NOTE:** When using these tank mixtures, do not exceed 3.25 quarts of Roundup UltraMAX per acre or equivalent rates of other Roundup agricultural herbicides.

**GRAMOXONE® EXTRA**

**Broadcast Treatment**—Apply 1.5 to 3 pints of Gramoxone Extra in these tank mixtures immediately before, during or after planting. Use 1.5 to 2 pints when weeds are 1 to 3 inches tall, 2 to 2.5 pints when weeds are 3 to 6 inches tall and 2.5 to 3 pints when weeds are 6 inches tall. This mixture may not control weeds taller than 6 inches. Add a nonionic spreader surfactant (approved for use on crops) containing at least 75 percent surfactant active agent at 8 ounces per 100 gallons of diluted spray.

**2,4-D**

**Broadcast Treatment**—Apply 1 to 2 pints of 2,4-D (amine or low-volatile ester) in these tank mixtures. DO NOT use 2,4-D on light, sandy soils, or where soil moisture is inadequate for normal weed growth.

**PREEMERGENCE WEED CONTROL  
INTRRO**

**APPLICATION RATES:**

SOIL TEXTURE	BROADCAST RATE PER ACRE	
	INTRRO (quarts)	
Coarse	2.5	
Medium	3	
Fine	3 to 3.5	

Use the higher rate in the recommended ranges in areas where organic matter content is 3 percent or more.

**INTRRO plus ATRAZINE**

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

**APPLICATION RATES:**

SOIL TEXTURE	BROADCAST RATE PER ACRE		
	INTRRO (quarts)	+	atrazine 4L* (quarts)
Coarse-sandy loam only	1.75 to 2		1 to 2
Medium	2 to 2.5		1.2 to 2
Fine	2.5 to 3		1.6 to 2

\* Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

Use the higher rate of this product in the recommended ranges when fall panicum or crabgrass will be present. Use the higher rate of atrazine in the recommended ranges on soils with greater than 3 percent organic matter.

**NOTE:** CORN, PEANUTS, SORGHUM OR SOYBEANS\* can be planted the year following use of this mixture.

\* There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow-irrigated corn.

**INTRRO plus PRINCEP**

**APPLICATION RATES:**

SOIL TEXTURE	BROADCAST RATE PER ACRE		
	INTRRO* (quarts)	+	Princep 4L** (quarts)
Coarse-sandy loam only	1.75 to 2		1 to 1.2
Medium	2 to 2.5		1.2 to 1.6
Fine	2.5 to 3		1.6 to 2.2

\* Use the higher rate of this product in the recommended ranges when fall panicum or crabgrass will be present.

\*\* When using Princep 80W use equivalent rates. One quart of Princep 4L equals 1.25 pounds of Princep 80W. Use the higher rate of Princep in the recommended ranges on soils with greater than 3 percent organic matter.

**NOTE:** LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP, PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS.

**INTRRO plus ATRAZINE plus PRINCEP**

For additional preemergence control of heavy weed infestation and/or hard-to-control weeds, a three-way tank mixture of this product herbicide plus atrazine plus Princep may be applied.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

**APPLICATION RATES:**

SOIL TEXTURE *	BROADCAST RATE PER ACRE		
	INTRRO** (quarts)	+ atrazine 4L** (quarts)	+ Princep 4L*** (quarts)
Coarse	1.75 to 2	0.75 to 1	0.75 to 1
Medium	2 to 2.5	1 to 1.25	1 to 1.5
Fine	2.5 to 3	1 to 1.25	1 to 1.5

\* Use the higher rates of atrazine and Princep in the recommended ranges on soils with greater than 3 percent organic matter.

\*\* Use the higher rate of this product in the recommended ranges when fall panicum, crabgrass, or soils with greater than 3 percent organic matter will be present.

\*\*\* When using 4L formulations use the rates listed. Use equivalent rates when using 80W or 90% dry flowable formulations. One quart of the 4L formulation of these products equals 1.25 pounds of the 80W formulation. One quart of atrazine 4L or Princep 4L equals 1.1 pounds of the 90% dry flowable formulations of these products.

**NOTE:** LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP, PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS. CORN, PEANUTS, SORGHUM OR SOYBEANS\* CAN BE PLANTED THE YEAR FOLLOWING USE OF THIS MIXTURE.

\* There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow-irrigated corn.

Roundup Ready and Roundup UltraMAX are registered trademarks of Monsanto Technology LLC. All other trademarks are the property of their respective owners.

©2004 Monsanto Company  
St. Louis, Missouri 63167 USA

11/19/2003

12069C1-7




MONSANTO COMPANY  
 600 13TH STREET, N.W.  
 SUITE 660  
 WASHINGTON, D.C. 20005  
<http://www.monsanto.com>

4 May 2004

Document Processing Desk (NOTIF)  
 Registration Division – 7504C  
 U.S. Environmental Protection Agency  
 1921 Jefferson Davis Highway  
 Crystal Mall #2, Room 266A  
 Arlington, Virginia 22202

Attention: Mr. James A. Tompkins  
 Product Manager 25

Subject: **Intrro Preemergent Herbicide (EPA Reg. No. 524-314)  
 Final Printed Supplemental Labeling  
 For Use in Field Corn and Silage Corn in Hawaii Only and  
 For Use in Field Corn and Silage Corn in California Only  
 19-Nov-2003 Approval**



Dear Mr. Tompkins:

On 19 November 2003, EPA approved a Lasso herbicide label on 524-314 with full crop uses. On 17 December 2003, a subset of uses that did not include corn was approved as an alternate brand, Intrro herbicide, on 524-314. Monsanto is now notifying the Agency of 2 supplemental labeling for use in California and Hawaii to support the corn business need. These supplemental labels are identical to the approved Lasso label sections for use on field and silage corn in these states.

Label Type	Print Plate No.	Comparison of stamped v printed
Supplemental for HA	12069C1-7	Lasso Crops section for corn, stamped pgs 15 thru 28 of 19-Nov-03 stamp with tank mix deletions
Supplemental for CA	12069C1-8	Lasso Crops section for corn, stamped pgs 15 thru 28 of 19-Nov-03 stamp with tank mix deletions

Differences between 19-November-2003 Approved Label and Final Printed Supplemental labels

1. ~~Changed brand name to Intrro Preemergent Herbicide by Monsanto~~
2. Added standard supplemental label language and reference to read the product label for restrictions and precautionary language
3. Deletion of all language, weed list or other uses and tank mixes not applicable to field and silage corn
4. Deletion of tank-mix products not registered in Hawaii and California
  - a. Delete tank mixes with Balance Pro, Banvel, Marksman, linuron
5. Other minor language deletions referring to specific sections in the Lasso book that would not apply to Intrro supplemental labels.

*This notification of minor label revisions during preparation of the Final Printed Label is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR§ 152.46, and no other changes (beside those outlined herein) have been made to the labeling or the*

*confidential statement of formula of this product. I understand that it is a violation of 18 USC Sec. 1001 to willfully make any false statements to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement actions and penalties under sections 12 and 14 of FIFRA.*

The submitted Final Printed Supplemental Labels contain minor corrections of grammar and trademark reference changes. These changes do not expand the allowed scope of use sites, pests, or application rates beyond those already approved.

If you should have any questions regarding this submission, please call Dr. Russell P. Schneider at (202) 383-2866 or call me at (314) 694-6576.

Sincerely,



Dawn Fee-White  
Registration Manager

Cc: Russ Schneider