

READ AND FOLLOW THE ENTIRE LABEL FOR ACCORD™ HERBICIDE AND GARLON™ 4 BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the labeling for Accord and this supplement.

ACCORD LOGO  
Herbicide by Monsanto  
(EPA Reg. No. 524-326)

Accord plus Garlon 4 tank mixture for aid to burning treatment in forestry site preparation.


Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

\*Accord is a trademark of Monsanto Company.

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

FORESTRY SITE PREPARATION

TANK MIXTURE  
of  
ACCORD PLUS GARLON

When applied as directed under the conditions described, this product plus Garlon 4 provides burndown, partial control or suppression of woody brush, trees and herbaceous weeds labeled for this product. The tank mixture is recommended as an aid to burning in forestry site preparation.

Aid to burning - When applied prior to burning, this tank mixture may provide more rapid browning and desiccation of vegetation as part of a site preparation operation.

NOTE: THIS PRODUCT REQUIRES USE WITH A NONIONIC SURFACTANT. When using this product, mix 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution. Always read and follow the surfactant manufacturer's label recommendations.

Apply 3 to 5 quarts of this product plus 1 to 2 quarts of Garlon 4 per acre. Use the higher recommended rates where vegetation is heavy or dense, or where hard-to-control species are present. Apply ground applications in 10 to 60 gallons of total spray solution per acre. Apply aerial applications (helicopter only) in 5 to 30 gallons of total spray solution per acre. For best results, treatments should be made from late spring through early summer.

Refer to the Garlon 4 label for plantback restrictions.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of woody brush and trees, use the higher recommended rates.

Read and carefully observe the label claims, cautionary statements and all information on the labels of both products used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.


When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Accord before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

<sup>TM</sup>Garlon is a trademark of Dow Chemical Company.

MAP-2936  
©MONSANTO COMPANY 1988

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167



READ AND FOLLOW THE ENTIRE LABEL FOR ACCORD™ HERBICIDE AND ARSENAL™  
APPLICATORS CONCENTRATE BEFORE PROCEEDING WITH THE USE DIRECTIONS  
CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the labeling  
for Accord and this supplement.

---

ACCORD LOGO  
Herbicide by Monsanto  
(EPA Reg. No. 524-326)

Tank mixtures of Accord herbicide plus Arsenal Applicators Concentrate for  
forestry site preparation prior to planting loblolly pine.

---


Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or  
night, (314) 694-4000.

™Accord is a trademark of Monsanto Company.

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.



---

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner  
inconsistent with the labeling.

This label should be in the possession of the user at the time of  
the herbicide application.

---

FORESTRY SITE PREPARATION

---

TANK MIXTURE  
of  
ACCORD PLUS ARSENAL

Use as a site preparation treatment for forestry sites being planted to  
loblolly pine.

NOTE: THIS PRODUCT REQUIRES USE WITH A NONIONIC SURFACTANT. When using this product, mix 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution. Always read and follow the surfactant manufacturer's label recommendations.

Apply 3 to 5 quarts of this product with up to 1 pint of Arsenal Applicators Concentrate in 10 to 60 gallons of water per acre as a broadcast spray to woody brush, trees and herbaceous weeds after full leaf expansion until initiation of fall leaf color.

The lower tank mixture rates will provide acceptable herbaceous and woody brush and tree control for most forestry sites. Use the higher recommended rates for dense, tough-to-control woody brush and trees.

The tank mixture may be applied by ground or aerial application. When applied by ground, use the recommended rate in 10 to 60 gallons of spray solution per acre. When applied by air (helicopter only), use the recommended rates in 5 to 15 gallons of spray solution per acre.

Read and carefully observe the label claims, cautionary statements and all information on the labels of both products used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.

When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Accord before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

<sup>TM</sup>Arsenal is a trademark of American Cyanamid Company.

MAP-2934  
©MONSANTO COMPANY 1988

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167

536

READ AND FOLLOW THE ENTIRE LABEL FOR ACCORD™ HERBICIDE AND OUST™ BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the labeling for Accord and this supplement.

---

ACCORD LOGO  
Herbicide by Monsanto  
(EPA Reg. No. 524-326)

Tank mixture of Accord herbicide plus Oust for forestry site preparation prior to planting listed conifer species.

---


Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

™Accord is a trademark of Monsanto Company.

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.



---

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

---

FORESTRY SITE PREPARATION

---

TANK MIXTURE  
of  
ACCORD PLUS OUST

Use as a site preparation treatment for forestry sites being planted to jack pine, loblolly pine, red pine, slash pine and Virginia pine.

NOTE: THIS PRODUCT REQUIRES USE WITH A NONIONIC SURFACTANT. When using this product, mix 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution. Always read and follow the surfactant manufacturer's label recommendations.

Apply 2 to 5 quarts of this product with 2 to 4 ounces of Oust in 10 to 60 gallons of water per acre as a broadcast spray to woody brush, trees and herbaceous weeds.

For control of herbaceous weeds listed on the label for this product, use the lower recommended tank mixture rates. For control or partial control of labeled woody brush and trees, use the higher recommended tank mixture rates. Apply after full leaf expansion until initiation of fall leaf color.

The tank mixture may be applied by ground or aerial application. When applied by ground, use the recommended rate in 10 to 60 gallons of spray solution per acre. When applied by air (helicopter only), use the recommended rates in 5 to 15 gallons of spray solution per acre.

Read and carefully observe the label claims, cautionary statements and all information on the labels of both products used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.


When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Accord before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

"Oust is a trademark of E.I. du Pont de Nemours and Company.

MAP-2933  
©MONSANTO COMPANY 1988

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167



READ AND FOLLOW THE ENTIRE LABEL FOR ACCORD™ and ARSENAL™ HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the labeling for Accord and this supplement.

ACCORD LOGO  
Herbicide by Monsanto  
(EPA Reg. No. 524-326)

Accord plus Arsenal tank mixtures for control, partial control or suppression of woody brush, trees and herbaceous weeds in utility rights-of-way.


Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

™Accord is a trademark of Monsanto Company.

~~MONSANTO COMPANY~~  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.



DIRECTIONS FOR USE


It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

UTILITY RIGHTS-OF-WAY

TANK MIXTURE  
of  
ACCORD PLUS ARSENAL

When applied as directed under the conditions described, this product plus Arsenal provides control, partial control or suppression of woody brush, trees and herbaceous weeds labeled for this product. The tank mixture is recommended for use on utility rights-of-way.



NOTE: THIS PRODUCT REQUIRES USE WITH A NONIONIC SURFACTANT. When using this product, mix 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution. Always read and follow the surfactant manufacturer's label recommendations.

Use 3 to 5 quarts of this product with up to 1 quart of Arsenal per acre as a broadcast application. Use 10 to 60 gallons of water per acre for ground applications and 5 to 30 gallons per acre for aerial applications (helicopter only). Apply after full leaf expansion until initiation of fall leaf color.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

Read and carefully observe the label claims, cautionary statements and all information on the labels of both products used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.


When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Accord before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

<sup>TM</sup>Arsenal is a trademark of American Cyanamid Company.

MAP-2937  
©MONSANTO COMPANY 1988

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167



READ AND FOLLOW THE ENTIRE LABEL FOR ACCORD™ HERBICIDE AND GARLON™ 4 BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the labeling for Accord and this supplement.

---

ACCORD LOGO  
Herbicide by Monsanto  
(EPA Reg. No. 524-326)

Accord plus Garlon 4 tank mixture for burndown and partial control or suppression of woody brush, trees and herbaceous weeds in utility rights-of-way.

---


Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

™Accord is a trademark of Monsanto Company.

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.



---

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

---

UTILITY RIGHTS-OF-WAY

---

TANK MIXTURE  
of  
ACCORD PLUS GARLON

When applied as directed under the conditions described, this product plus Garlon 4 provides burndown and partial control or suppression of woody brush, trees and herbaceous weeds labeled for this product. The tank mixture is recommended for use on utility rights-of-way.

NOTE: THIS PRODUCT REQUIRES USE WITH A NONIONIC SURFACTANT. When using this product, mix 2 or more quarts of a nonionic surfactant per 100 gallons of spray solution. Always read and follow the surfactant manufacturer's label recommendations.

Hand-held and High-Volume Applications: Use 2 to 4 quarts of this product plus  $\frac{1}{2}$  to 2 quarts of Garlon 4 per 100 gallons of spray solution. Apply to foliage as a spray-to-wet application. Spray coverage should be uniform and complete for best results. Do not spray to the point of runoff.

Broadcast Applications with Ground Equipment: Use 2 to 5 quarts of this product plus  $\frac{1}{2}$  to 2 quarts of Garlon 4 in 10 to 60 gallons of water per acre.

Aerial Applications (helicopter only): Use 2 to 5 quarts of this product plus  $\frac{1}{2}$  to 2 quarts of Garlon 4 and apply in 5 to 15 gallons of water per acre.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of woody brush and trees, use the higher recommended rates.

Read and carefully observe the label claims, cautionary statements and all information on the labels of both products used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.

When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Accord before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

<sup>TM</sup>Garlon is a trademark of Dow Chemical Company.

MAP-2935  
©MONSANTO COMPANY 1988

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167

# DEPLOY™

Monsanto

FOR USE IN ILLINOIS, INDIANA, IOWA,  
KANSAS, MISSOURI, AND NEBRASKA.

## Complete Directions for Use

**AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.**

DEPLOY is a trademark of Monsanto Company  
EPA Reg. No. 524-325

1988-1 892 58-000 01 53

Read each of these sections of this label for essential product performance information.

USE THESE CONVENIENT SECTION MARKERS.



**ACCEPTED**

JUN - 1 1988

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-325

Read the entire label before using this product

Use only according to label instructions

Read **LIMIT OF WARRANTY AND LIABILITY** before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED

### LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet (Directions) when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This Warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including but not limited to incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Direc-

tions application in any manner not explicit by set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions on or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this **LIMIT OF WARRANTY AND LIABILITY** which may not be varied by any verbal or written agreement.

### PRECAUTIONARY STATEMENTS

Hazards to  
Humans and Domestic Animals

Keep out of reach of children

### CAUTION!

MAY CAUSE EYE IRRITATION.

Avoid contact with eyes, skin or clothing

Wash thoroughly with soap and water after handling

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Call a physician.

IF ON SKIN, immediately flush with plenty of water.

Wash clothing before reuse.

In case of an emergency involving this product

Call Collect, day or night, (814) 694-4000

### Environmental Hazards

Do not apply directly to water, wetland (swamps, bogs, marshes or potholes). Do not contaminate water or equipment or disposal of wastes.

### Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray

solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

#### ACTIVE INGREDIENT

*Isopropylamine salt of glyphosate	41.5%
INERT INGREDIENTS	58.5%
	100.0%

\*Contains 480 grams per litre or 4 pounds of the active ingredient isopropylamine salt of N-(phosphonomethyl) glycine per U.S. gallon. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid glyphosate.

Product is protected by  
U.S. Pat. No. 3,799,728 and  
U.S. Pat. No. 4,405,531  
Other patents pending

MONSANTO COMPANY 1988

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI, 63167 U.S.A.



### GENERAL INFORMATION

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

This product is a water soluble liquid which mixes readily with water to be applied as a foliar spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field-type sprayers after dilution and thorough mixing with water in accordance with label instructions.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, but on most perennial weeds may not occur for 7 days or more. Extremely cool or cloudy weather following treatment may slow down activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of aboveground growth and deterioration of underground plant parts.

Unless otherwise specified on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "Weeds Controlled" section of this label. Unemerged plants arising from unattached underground rhizomes or root stocks of perennials will not be affected by the spray and will continue to grow.

Always use the higher rate of this product per acre when: (1) weed growth is heavy or dense or (2) weeds are growing in an undisturbed (noncultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust. Reduced control may result when applications are made to weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of Daploy with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance of Daploy.

### MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. DO NOT APPLY UNDER WIND OR OTHER CONDITIONS WHICH ALLOW DRIFT TO OCCUR. HAND GUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS. NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

#### MIXING

This product requires use with a nonionic surfactant. When using this product, mix 4 to 8 quarts (1 to 2 percent solution) of a nonionic surfactant per 100 gallons of spray solution. Use the 8-quart rate for any surfactant containing less than 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water. Add the proper amount of this product (see "Directions for Use" and "Weeds Controlled" sections of this label) and nonionic surfactant near the end of the filling process and mix well. Remove hose from tank immediately after filling to avoid siphoning back into the corner source. During mixing and application, foaming of the spray solution may

occur. To prevent or minimize foam, avoid the use of mechanical agitators, place the filling hose below the surface of the spray solution, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved anti-foam or defoaming agent.

CLEAN SPRAYER AND PARTS IMMEDIATELY AFTER USING THIS PRODUCT BY THOROUGHLY FLUSHING WITH WATER AND DISPOSE OF RINSATE ACCORDING TO LABELED USE OR DISPOSAL INSTRUCTIONS.

#### TANK MIXTURES

Always predetermine the compatibility of labeled tank mixtures of this herbicide with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of Daploy herbicide with water as follows:

1. Place a 20 ft 35 mesh screen or netting basket over filling top.
2. Through the screen, fill the sprayer tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.

For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

Keep people and pets off treated areas until spray solution has dried to prevent transfer of this product onto desirable vegetation.

#### ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation, since minute quantities of this herbicide can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

BEST AVAILABLE COPY

- If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
- Continue filling the sprayer tank with water and add the required amount of **Deploy** herbicide near the end of the filling.
- Add nonionic surfactant to the spray tank before completing the filling process.
- Add individual formulations to the spray tank as follows: wettable powder, flowable emulsifiable concentrate, drift control additive, water-soluble liquid followed by surfactant.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water.

## ADDITIVES

### AMMONIUM SULFATE

The addition of 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water may increase the performance of **Deploy** and **Deploy** plus 2,4-D amine or Banvel<sup>®</sup> tank mixtures on annual and perennial weeds. The improvement in performance may be apparent where environmental stress is a concern. Low-quality ammonium sulfate may contain material that will not readily dissolve which could result in nozzle tip plugging. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, pre-dissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet lines. Ensure that ammonium sulfate is completely dissolved in the spray tank before adding herbicides or surfactant. Thoroughly rinse the spray system with clean water after use to reduce corrosion.

**NOTE:** The use of ammonium sulfate as an additive does not preclude the need for surfactant. Do not use herbicide rates lower than recommended in this label.

<sup>®</sup>Banvel is a trademark of Sandez, Inc.

## APPLICATION EQUIPMENT AND TECHNIQUES

Do not apply this product through any type of irrigation system.

This product may be applied with the following application equipment:

**Aerial** — Fixed-wing and Helicopter

**Boom** — Conventional Broadcast Spray

**Controlled Droplet Applicator (CDA)** — Hand-held or boom-mounted applicators which produce a spray consisting of a narrow range of droplet sizes.

**Hand-held and High-Volume Spray Equipment** — Knapsack and backpack sprayers, pump-up pressure sprayers, hand-pumps, handwands, lances and other hand-held spray equipment used to direct the spray onto weed foliage and vertically-mounted high-volume spray equipment for spray-to-wet applications.

### AERIAL EQUIPMENT

Use the recommended rates of this herbicide in 3 to 15 gallons of water per acre unless otherwise specified on this label. See the "Weeds Controlled" section of this label for specific rates. (Aerial applications of this product may only be made in fallow and reduced tillage systems, and preharvest. Refer to the individual use area sections of this label for recommended volumes and application rates.)

### BOOM EQUIPMENT

For control of annual or perennial weeds listed on this label using conventional boom equipment — Use the recommended rates of this product in 10 to 40 gallons of water per acre as a broadcast spray unless otherwise specified on this label. See the "Weeds Controlled" section of this label for specific rates. As density of weeds increases, spray gallonage should be increased within the recommended range to ensure complete coverage. Carefully select proper nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

### CONTROLLED DROPLET APPLICATION (CDA)

The rate of this product applied per acre by vehicle-mounted CDA equipment must not be less than the amount recommended in this label when applied by conventional broadcast equipment. For vehicle-mounted CDA equipment, apply 3 to 15 gallons of water per acre.

For the control of labeled annual weeds with hand-held CDA units, apply a 20 percent solution of this product plus a 5 percent solution of nonionic surfactant at a flow rate of 2 fluid ounces per minute and a walking speed of 1.5 MPH (1 quart per acre). For the control of labeled perennial weeds, apply a 20 to 40 percent solution of this product plus a 5 to 10 percent solution of nonionic surfactant at a flow rate of 2 fluid ounces per minute and a walking speed of 0.75 MPH (2 to 4 quarts per acre).

Avoid direct application to any body of water.

**AVOID DRIFT — DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION. APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.**

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which disperse spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application — To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

Controlled droplet application equipment produces a spray pattern which is not easily visible. Extreme care must be exercised to avoid spray or drift contacting the foliage or any other green tissue of desirable vegetation as damage or destruction may result.

### HAND-HELD and HIGH-VOLUME EQUIPMENT

Use coarse sprays only.

Mix this product in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff.

Use a 1 percent solution, unless otherwise specified. For best results, use a 5 percent solution on harder-to-control perennials, such as dock, field bindweed, hemp dogbane, milkweed and Canada thistle.

Less than complete coverage of weeds may result from the use of slugy equipment designed for motorized spot treatments. Where less than complete coverage of annual weeds occurs, use a 5 percent solution.

BEST APPLICATION TECHNIQUES

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the following table:

Spray Solution				
DESIRED VOLUME	AMOUNT OF DEPLET <sup>1</sup>			
	1%	1 1/2%	2%	5%
1 gallon	1-1/8 oz. 2 oz.	2-1/8 oz. 4 oz.	3-1/8 oz. 6 oz.	1-1/2 gal.
25 gallons	1 qt. 1-1/2 qt. 2 qt.	2 qt. 3 qt.	3 qt. 5 qt.	1-1/2 gal.
100 gallons	1 gal. 1-1/2 gal. 2 gal.	2 gal. 3 gal.	3 gal. 5 gal.	1-1/2 gal.

2 tablespoons = 1 ounce  
Add surfactant at a rate of 1 to 2 oz. per gallon of spray mix

For use in knapsack sprayers it is suggested that the proper amount of this product be mixed with water in a larger container. Fill sprayer with the mixed solution and add the correct amount of surfactant.

For hand held CONTROLLED DROPLET APPLICATORS see the "Controlled Droplet Applicator (CDA)" section of this label.

### WEEDS CONTROLLED

This herbicide controls many annual and perennial grasses and broadleaf weeds.

#### ANNUAL WEEDS

- Apply to actively growing grass and broadleaf weeds.
- Allow at least 3 days after treatment before tillage.
- For maximum agronomic benefit, apply when weeds are 6 inches or less in height.
- To prevent seed production, applications should be made prior to seedhead formation.
- This product does not provide residual control; therefore, delay application until maximum weed emergence. Repeat treatments may be necessary to control later germinating weeds.

#### LOW-VOLUME BROADCAST APPLICATION (LOW-RATE TECHNOLOGY)

When applied as directed under the conditions described, this product will control the weeds listed below when:

1. Water carrier volumes are 3 to 10 gallons per acre for ground applications and 3 to 5 gallons per acre for aerial applications. (See the "Aerial Equipment" section of this label for approved sites.)

#### NOTE

- The addition of 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water may increase the performance of this herbicide on annual weeds. The improvement in performance may be apparent where environmental stress is a concern. Refer to the "Mixing, Additives and Application Instructions" section of this label.
- Do not tank mix with soil residual herbicides when using these rates unless otherwise specified.
- Do not use these rates for weeds that have been mowed, grazed, or cut. Refer to the "High Volume Broadcast Application" portion of this section for recommended rates.
- Refer to the "Tank Mixtures" portion of this section for control of additional broadleaf weeds.

WEED SPECIES	MAXIMUM HEIGHT	RATE PER ACRE**
Foxtail <i>Setaria spp.</i>	12"	8 oz.
Barnyardgrass <i>Echinochloa crus-galli</i>	6"	12 oz.
Bluegrass, annual <i>Poa annua</i>		
Brome, downy <sup>1</sup> <i>Bromus tectorum</i>		
Mustard, blue <i>Chorispora tenella</i>		
Mustard, tansy <i>Descurainia pinnata</i>		
Mustard, tumble <i>Sisymbrium altissimum</i>		
Mustard, wild <i>Sinapis arvensis</i>		
Spry, umbrella <i>Holosteum umbellatum</i>		
Barley <i>Hordeum vulgare</i>	12"	
Rye <i>Secale cereale</i>		
Sandbar, field <i>Cenchrus spp.</i>		
Shattercane <i>Sorghum bicolor</i>		

WEED SPECIES	MAXIMUM HEIGHT	RATE PER ACRE**
Stinkgrass <i>Eragrostis ciliaris</i>	12"	12 oz.
Wheat <i>Triticum aestivum</i>	18"	
Morningglory <i>Ipomea spp.</i>	3"	16 oz.
Bluegrass, bulbous <i>Poa bulbosa</i>	6"	
Cheat <i>Bromus secalinus</i>		
Chickweed, mouseear <i>Cerastium vulgatum</i>		
Corn <i>Zea mays</i>		
Groundail, common <i>Senecio vulgaris</i>		
Horseweed/Marestail <i>Conyza canadensis</i>		
Pennycress, field Fanweed <i>Thlaspi arvense</i>		

WEED SPECIES	MAXIMUM HEIGHT	RATE PER ACRE**
Rocket, London <i>Sisymbrium irio</i>	3"	16 oz.
Ryegrass, Italian <i>Lolium multiflorum</i>		
Shepherdspurge <i>Capsella bursa-pastoris</i>		
Bittercap <i>Ranunculus spp.</i>	12"	
Cocklebur <i>Xanthium strumarium</i>		
Crabgrass <i>Digitaria spp.</i>		
Oxardandelion <i>Krigia cespitosa</i>		
Fabertail, trifoliate <i>Camelina microcarpa</i>		
Foxtail, Cocksfoot <i>Alopecurus carolinense</i>		
Johnsongrass, seedling <i>Sorghum halepense</i>		
Lambquarters, common <i>Chenopodium album</i>		
Oats, wild <i>Avena fatua</i>		

**BEST AVAILABLE COPY**

WEED SPECIES	MAXIMUM HEIGHT	RATE PER ACRE**
<b>Panicum, fall</b> <i>Panicum dichotomiflorum</i>	2"	32 oz
<b>Panicum, Texas</b> <i>Panicum terans</i>		
<b>Pigweed, redroot</b> <i>Amaranthus retroflexus</i>		
<b>Pigweed, smooth</b> <i>Amaranthus hybridus</i>		
<b>Witchgrass</b> <i>Panicum capillare</i>		
<b>Signalgrass, broadleaf</b> <i>Bracharia platyphylla</i>	6"	24 oz
<b>Tea weed</b> <i>Sida spinosa</i>	3"	32 oz
<b>Geranium, Carolina</b> <i>Geranium carolinianum</i>	12"	
<b>Goosegrass</b> <i>Eleusine indica</i>		

WEED SPECIES	MAXIMUM HEIGHT	RATE PER ACRE**
<b>Prunella, cutleaf evening</b> <i>Desmodium illinoense</i>	12"	32 oz
<b>Pursley, Florida</b> <i>Richardia scabra</i>		
<b>Sicklepod</b> <i>Cassia obtusifolia</i>		
<b>Spanishneedles</b> <i>Bidens bipinnata</i>		
<b>Filaree</b> <i>Erodium spp</i>	12"	48 oz

\*For control in no-till systems use 16 fluid ounces per acre  
 \*\*For those rates less than 32 ounces per acre. Deploy at rates up to 32 ounces per acre may be used where heavy weed densities exist

**TANK MIXTURES**  
 DEPLOY™ plus BANVEL  
 plus NONIONIC SURFACTANT  
 DEPLOY™ plus 2,4-D AMINE  
 plus NONIONIC SURFACTANT

These tank mixtures are recommended for use in fallow and reduced tillage areas only. Follow use directions as given in this Low Volume Broadcast Application section.

Deploy plus Banvel or 2,4-D amine will control the annual grasses and broadleaf weeds previously listed for Deploy alone at the indicated heights (except 8 ounces per acre applications) plus the following broadleaf weeds. For those weeds previously listed at 8 ounces of Deploy herbicide alone per acre use the 12-ounce rate in these tank mixtures.

NOTE: Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products used in tank mixtures. Some crop injury may occur if Banvel is applied within 45 days of planting. The addition of Banvel in a mixture with this product may provide short-term residual control of selected weed species.

Apply 12 to 16 ounces of this product plus 0.25 lb a.e. of Banvel or 0.5 pound a.e. of 2,4-D amine per acre to control dense populations of the following annual broadleaf weeds when less than 6 inches in height.

<b>Rochia*</b> <i>Kochia scoparia</i>	<b>Pigweed, redroot</b> <i>Amaranthus retroflexus</i>
<b>Lambs' ears</b> <i>Chenopodium album</i>	<b>Thistle, Russian</b> <i>Salsola kali</i>

**Lettuce, prickly**  
*Lactuca serriola*

\*Controlled with Banvel tank mixture only

**HIGH-VOLUME BROADCAST APPLICATION**

When applied as directed under the conditions described, this product will control the weeds listed below when water carrier volumes are 10 to 40 gallons per acre for ground applications.

Use 1 quart of this product per acre if weeds are less than 6 inches tall. If weeds are over 6 inches tall, use 1.5 quarts of this product per acre. If weeds have been mowed, grazed, or cut, allow adequate time for new growth to recommended stages prior to treatment. These rates will also provide control of weeds listed in the Low-volume Broadcast Application section.

**WEED SPECIES**

<b>'alsamapple*</b> <i>Momordica charantia</i>	<b>Panicum</b> <i>Panicum spp</i>
<b>Bassia, treehoop</b> <i>Bassia hyssopifolia</i>	<b>Ragweed, common</b> <i>Ambrosia artemisiifolia</i>
<b>Brome</b> <i>Bromus spp</i>	<b>Ragweed, giant</b> <i>Ambrosia trifida</i>

<b>Fiddleneck</b> <i>Amsinckia spp</i>	<b>Smartweed, Pennsylvania</b> <i>Polygonum pennsylvanicum</i>
<b>Flaxleaf, hairy</b> <i>Corylis bonariensis</i>	<b>Sowthistle, annual</b> <i>Scorpus oleraceus</i>
<b>Flanbane</b> <i>Erigeron spp</i>	<b>Sunflower</b> <i>Helianthus annuus</i>
<b>Kochia</b> <i>Kochia scoparia</i>	<b>Thistle, Russian</b> <i>Salsola kali</i>
<b>Lettuce, prickly</b> <i>Lactuca serriola</i>	<b>Valerian</b> <i>Abutilon theophrasti</i>

\*Apply with hand-held equipment only.

**PERENNIAL WEEDS**

**TANK MIXTURES**

When applied as directed, under the conditions described, Banvel plus Banvel or 2,4-D amine will suppress or control the following perennial broadleaf weed:

**Birdweed, field**  
*Convolvulus arvensis*

For suppression, apply 16 ounces of this product plus 0.5 lb a.e. of 2,4-D amine or 0.25 lb a.e. of Banvel in 5 to 10 gallons of water per acre for ground applications and 3 to 5 gallons of water per acre for aerial applications. Applications should be delayed until maximum emergence has occurred and when weeds are between 6 to 18 inches in length. These tank mixtures may be applied using aerial application equipment (except in California) in fallow and reduced tillage systems only.

Refer to the specific product labels for crop rotation restrictions and cautionary statements of all products in tank mixture.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

**Storage and Disposal**

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**STORAGE**  
STORE ABOVE 10°F (-12°C) TO KEEP PRODUCT FROM CRYSTALLIZING.

Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°-72°F for several days to redissolve and shake or roll to mix well before using.

**DISPOSAL**  
Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or buried on site in a safe place away from water supplies. All disposal should be in accordance with applicable Federal state or local procedures.

Emptied container retains approx. 10% product residue. Observe all label safeguards until container is destroyed.

Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**BEST AVAILABLE COPY**

See "General Information" and "Mixing Additives and Application Instructions" sections of this label for essential product performance information and the following sections for specific recommended uses.

**EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURF GRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.**

**NOTE:** If spraying areas adjacent to desirable plants, use a shield made of cardboard, sheet metal, or plywood while spraying to help prevent spray from contacting foliage of desirable plants.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

This product does not provide residual weed control. For subsequent weed control, follow a label approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

## CROPPING SYSTEMS

See "General Information" and "Mixing Additives and Application Instructions" sections of this label for essential product performance information.

See the following Cropping Systems sections for specific recommended uses.

**EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE, GREEN STEMS OR FRUIT OF DESIRABLE CROPS, PLANTS, TREES OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.**

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Except as otherwise specified on this label, repeat treatments must be made before the crop emerges in accordance with the instructions of this label.

Except as otherwise specified in a crop section of this label, the combined total of all treatments must not exceed 3 quarts per acre of this product per year.

Do not plant subsequent crops other than those on the label for one year following application.

For other cropping systems, except following spot treatment in forage grasses and legumes, do not harvest or feed treated crops for 8 weeks after application. Allow 14 days following spot treatment or selective equipment use before grazing domestic livestock or harvesting forage grasses and legumes.

When applied as directed for Cropping Systems under the conditions described, this product controls annual and perennial weeds listed on this label prior to the emergence of these crops.

ALFALFA

BARLEY

FORAGE GRASSES

FORAGE LEGUMES

OATS

WHEAT

## FALLOW AND REDUCED TILLAGE SYSTEMS

Use this product in fallow and reduced tillage systems for control of annual weeds prior to emergence of crops listed in this label. Refer to the "Weeds Controlled" section of this label for specific rates and instructions. This product may be applied using ground or aerial spray equipment. See the "Application Equipment and Techniques" section of this label for instructions.

### TANK MIXTURES

DEPLOY™ plus **BANVEL** plus  
NONIONIC SURFACTANT

DEPLOY plus **2,4-D AMINE** plus  
NONIONIC SURFACTANT

The addition of Banvel in a mixture with this product may provide short-term residual control of selected weed species. Some crop injury may occur if Banvel is applied within 45 days of planting. Refer to the Banvel and 2,4-D labels for cropping restrictions and other use instructions.

These recommended tank mixtures may be applied using ground or aerial spray equipment. Refer to the "Weeds Controlled" section of this label for specific rates and instructions.

## PASTURES

Apply this product prior to planting forage grasses and legumes.

**Pasture or Hay Crop Renovation** — When applied as a broadcast spray, this product controls the annual and perennial weeds listed in this label prior to planting forage grasses or legumes. Remove domestic livestock before application and wait 8 weeks after application before grazing or harvesting.

**Spot Treatment** — When applied as a spot treatment as recommended, this product controls annual and perennial weeds listed in this label which are growing in pastures, forage grasses and forage legumes composed of bluegrass, brome, fescue, orchardgrass, ryegrass, timothy, wheatgrass, alfalfa or clover.

For spot treatment, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Further applications may be made in the same area at 30 day intervals. Remove domestic livestock before application and wait 14 days after application before grazing livestock or harvesting.

See the "General Information," "Mixing Additives and Application Instructions" and the "Weeds Controlled" sections of this label.

EPA Reg. No. 524-326

892 68 000 01/53

©MONSANTO COMPANY 1988

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.



**BEST AVAILABLE COPY**

# DEPLOY

HERBICIDE BY **Monsanto**

For use in Illinois, Indiana, Iowa, Kansas, Missouri and Nebraska

Carefully follow detailed instructions in label booklet.  
**AVOID CONTACT WITH FOLIAGE, GREEN STEMS OR FRUIT OF CROPS, DESIRABLE  
 PLANTS AND TREES SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT!**

- Read the entire label before using this product
- Use only according to label instructions
- Read **LIMIT OF WARRANTY AND LIABILITY** before buying or using. If terms are not acceptable, return at once, unopened
- REFUNDATION OR REFRESHING IS PROHIBITED

<b>ACTIVE INGREDIENT</b>	
* Isopropylamine salt of glyphosate	41.5%
<b>INERT INGREDIENTS</b>	58.5%
	100.0%

\* Contains 480 grams per litre or 4 pounds of the active ingredient isopropylamine salt of N (phosphonomethyl) glycine per U.S. gallon. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid glyphosate

**TWO 2.5 GAL UNITS**

**ACCEPTED**  
 - 1 1988  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-326

Keep out of reach of children  
**CAUTION!** Read precautions on side panel

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

Keep out of reach of children

**CAUTION!**

**MAY CAUSE EYE IRRITATION**

Avoid contact with eyes, skin or clothing

Wash thoroughly with soap and water after handling

**FIRST AID: IF IN EYES**, immediately flush with plenty of water for at least 15 minutes. Call a physician

**IF ON SKIN**, immediately flush with plenty of water. Wash clothing before reuse

In case of an emergency involving this product  
 Call Collect day or night (314) 694 4000

**Environmental Hazards**

Do not apply directly to water or wetland (swamps, bogs, marshes or potholes). Do not contaminate water by cleaning of equipment or disposal of wastes

**Physical or Chemical Hazards**

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic lined steel containers

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product reacts with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling

**Storage and Disposal**

Do not contaminate water, landfills, seed or feed by storage or disposal

**STORAGE**  
 STORE ABOVE 10°F (-12°C) TO KEEP PRODUCT FROM CRYSTALLIZING

Crystals will settle to the bottom. If allowed to crystallize, place in warm area 80°F (27°C) for several days to redissolve and mix well before using

**DISPOSAL**  
 Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or buried on site in a safe place away from water supplies. All disposal should be in accordance with applicable Federal, State or local procedures

© 1987 Monsanto Company  
Registered under  
EPA Reg. No. 101-103-01

# DEPLOY™

HERBICIDE BY **Monsanto**

FOR USE IN ILLINOIS, INDIANA, IOWA, KANSAS, MISSOURI AND NEBRASKA

Carefully follow detailed instructions in attached label booklet.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS OR FRUIT OF CROPS DESIRABLE  
PLANTS AND TREES SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT

Keep out of reach of children

**CAUTION!**

Read the entire label before using this product.

Use only according to label instructions.

Read **LIMIT OF WARRANTY AND LIABILITY** before buying or  
using. If terms are not acceptable, return at once, unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED

 **Monsanto**  
Monsanto Company  
P.O. Box 131  
St. Louis, MO 63168

**ACTIVE INGREDIENT**

*Isopropylamine salt of glyphosate	4.5%
<b>INERT INGREDIENTS</b>	<u>55.5%</u>
	60.0%

\*Contains 480 grams per litre or 4 pounds of the active ingredient isopropylamine salt of N-phosphonomethyl glycine per U.S. gallon. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid glyphosate.

## NET 2.5 GAL

100 Page No. 524 326  
1000 1000 1000

1000 1000 1000  
1000 1000 1000

*1000 1000 1000*

# DEPLOY™

HERBICIDE BY **Monsanto**

## PRECAUTIONARY STATEMENTS

Hazards to  
Humans and Domestic Animals

Keep out of reach of children

### CAUTION!

MAY CAUSE EYE IRRITATION

Avoid contact with eyes, skin or clothing

Wash thoroughly with soap and water after handling

**FIRST AID IF IN EYES:** Immediately flush with plenty of water for at least 15 minutes. Call a physician.

**IF ON SKIN:** Immediately flush with plenty of water. Wash clothing before reuse.

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000

## Environmental Hazards

Do not apply directly to water or wetland (swamps, bogs, marshes or borofores). Do not contaminate water by cleaning of equipment or disposal of wastes.

## Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic-lined steel containers.

**DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

## Storage and Disposal

Do not contaminate water, foodstuffs, seed or feed by storage or disposal.

### STORAGE

STORE ABOVE 10°F (12°C) TO KEEP PRODUCT FROM CRYSTALLIZING.

Crystals will settle to the bottom. If allowed to crystallize, place in a warm room (68° F/20° C) for several days to redissolve and shake or roll to mix well before using.

### DISPOSAL

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or buried on site in a safe place away from water supplies. All disposal should be in accordance with applicable Federal, state or local procedures.

Empty container retains vapor and product residue. Observe all label safeguards until container is destroyed.

Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

See label booklet for Complete Directions for Use.

## LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims, whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

Product protected by U.S. Pat. Nos. 2,759,759  
and 4,405,531. Other patents pending.

MONTANTO COMPANY, 1996

® Trademarks of Monsanto Company

Page 2 of 2

892 69-090 03-531