

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ACCORD (R) HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Accord herbicide and this supplement.

ACCORD

HERBICIDE BY MONSANTO

EPA Reg. No. 524-326

ACCORD HERBICIDE FOR NONCROP AREAS SUCH AS AIRPORTS, DITCH BANKS, DRY DITCHES, DRY CANALS, FENCEROWS, HIGHWAYS, INDUSTRIAL SITES, PIPELINES, RAILROADS, ROADSIDES, STORAGE AREAS, AND SIMILAR INDUSTRIAL OR NONCROP AREAS.

Keep out of reach of children.

CAUTION!


In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

Accord (R) is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.



Industrial Uses

See "General Information" and "Mixing and Application Instructions" sections of the label booklet for Accord for essential product performance information and the following sections for specific recommended uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds.

This product does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

When applied as directed under conditions described, this product controls annual weeds and perennial weeds listed in this supplemental label and woody brush listed on the label booklet for Accord in areas such as airports, ditch banks, dry ditches, dry canals, fencerows, golf courses, highways, industrial plant sites, lumber yards, parking areas, parks, petroleum tank farms and pumping installations, pipelines, power and telephone rights-of-way, railroads, schools, storage areas, other public areas and similar industrial or noncrop areas.

WEEDS CONTROLLED

Annual weeds

When applied as directed under the conditions described, this product will control the weeds listed below when water carrier volumes are 10 to 40 gallons per acre for ground applications.

Broadcast Application-Use 1 quart of this product per acre if weeds are less than 6 inches tall. If weeds are over 6 inches tall, use 1.5 quarts of this product per acre. If weeds have been mowed, grazed, or cut, allow adequate time for new growth to recommended stages prior to treatment.

BEST AVAILABLE COPY

High-volume Application-For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff.

For control of annual weeds listed on this label, apply a 0.5 percent solution of this product to weeds less than 6 inches in height or runner length. Apply prior to seedhead formation in grass or bud formation in broadleaves. Allow three or more days before tillage or mowing.

For annual weeds over 6 inches tall, or unless otherwise specified, use a 1 percent solution.

ANNUAL WEED SPECIES

Balsamapple <i>Monardella charantia</i>	Falsedoe, smallseed <i>Camelina microcarpa</i>	Mustard, tansy <i>Descurainia pinnata</i>	Shattercan <i>Sorghum bicolor</i>
Barley <i>Hordeum vulgare</i>	Fiddleneck <i>Amsinckia spp</i>	Mustard, tumble <i>Sisymbrium altissimum</i>	Shepherdspurse <i>Capsella bursa-pastoris</i>
Barnyardgrass <i>Echinochloa crus-galli</i>	Flaxleaf Fleawane <i>Conyza bonariensis</i>	Mustard, wild <i>Sinapis arvensis</i>	Signalgrass, broadleaf <i>Brachiaria platyphylla</i>
Bassia, livehoak <i>Bassia hyssopifolia</i>	Florbane <i>Erigeron spp</i>	Oats, wild <i>Avena fatua</i>	Smartweed, Pennsylvania <i>Polygonum pennsylvanicum</i>
Bluegrass, annual <i>Poa annua</i>	Fentail <i>Setaria spp.</i>	Panicum <i>Panicum spp</i>	Southwale, annual <i>Sonchus oleraceus</i>
Bluegrass, bulbous <i>Poa bulbosa</i>	Fentail, Caroline <i>Alopecurus carolinianus</i>	Pennycress, field <i>Thlaspi arvense</i>	Spanishneedles <i>Bidens bipinnata</i>
Brome <i>Bromus spp</i>	Groundsel, common <i>Senecio vulgaris</i>	Pigweed, redroot <i>Amaranthus retroflexus</i>	Stinkgrass <i>Eragrostis cilianensis</i>
Buttercup <i>Ranunculus spp</i>	Horseweed/Marestail <i>Conyza canadensis</i>	Pigweed, smooth <i>Amaranthus hybridus</i>	Sunflower <i>Helianthus annuus</i>
Cheat <i>Bromus secalinus</i>	Kochia <i>Kochia scoparia</i>	Ragweed, common <i>Ambrosia artemisiifolia</i>	Thistle, Russian <i>Salsola kali</i>
Chickweed, mouseear <i>Cerastium vulgatum</i>	Lambquarters, common <i>Chenopodium album</i>	Ragweed, giant <i>Ambrosia trifida</i>	Spurry, umbrella <i>Holosteum umbellatum</i>
Cocklebur <i>Xanthum strumarium</i>	Lettuce, prickly <i>Lactuca scariola</i>	Rocket, London <i>Sisymbrium irio</i>	Vetchling <i>Abutilon theophrasti</i>
Corn, volunteer <i>Zea mays</i>	Morningglory <i>Ipomoea spp</i>	Rye <i>Secale cereale</i>	Wheat <i>Triticum aestivum</i>
Crabgrass <i>Digitaria spp</i>	Mustard, blue <i>Chorispora tenella</i>	Ryegrass, Italian <i>Lolium multiflorum</i>	Witchgrass <i>Panicum capillare</i>
Dwarf Mandarin <i>Argemone cespitosa</i>		Sandbar, field <i>Cenchrus spp</i>	

PERENNIAL WEEDS

Apply this product as follows to control or destroy most perennial weeds:

NOTE. If weeds have been mowed or tilled, do not treat until plants have resumed active growth and are at the recommended stages.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed. Repeat treatments must be made prior to crop emergence.

The addition of 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water may increase the performance of this product on perennial weeds. The improvement in performance may be apparent where environmental stress is a concern.

When applied as recommended under the conditions described, this product WILL CONTROL the following PERENNIAL WEEDS:

Alfalfa <i>Medicago sativa</i>	Horse radish <i>Armoracia rusticana</i>
Alligatorweed* <i>Alternanthera philoxeroides</i>	Johnsongrass <i>Sorghum halepense</i>
Artichoke, Jerusalem <i>Helianthus tuberosus</i>	Kilburygrass <i>Pennisetum clandestinum</i>
Bahiagrass <i>Paspalum notatum</i>	Knapweed <i>Centaurea repens</i>
Bentgrass <i>Agrostis spp.</i>	Lantana <i>Lantana camara</i>
Bermudagrass <i>Cynodon dactylon</i>	Milkweed <i>Asclepias spp.</i>
Bindweed, field <i>Convolvulus arvensis</i>	Muhly, wirestem <i>Muhlenbergia frondosa</i>
Bluegrass, Kentucky <i>Poa spp.</i>	Mullein, common <i>Verbascum thapsus</i>
Blueweed, Texas <i>Helianthus ciliaris</i>	Napiergrass <i>Pennisetum purpureum</i>
Brackenfern <i>Pteridium squilinum</i>	Nightshade, silverleaf <i>Solanum elaeagnifolium</i>
Bromegrass, smooth <i>Bromus inermis</i>	Notweed, purple, yellow <i>Cyperus rotundus</i> <i>Cyperus esculentus</i>
Bursage, woollyleaf <i>Franseria tomentosa</i>	Orchardgrass <i>Dactylis glomerata</i>

Canarygrass, reed <i>Phalaris arundinacea</i>	Pampas <i>Cortaderia jubata</i>
Cattail <i>Typha</i> spp	Paragrass <i>Brachiaria mutica</i>
Clover, red <i>Trifolium pratense</i>	Phragmites* <i>Phragmites</i> spp
Clover, white <i>Trifolium repens</i>	Quackgrass <i>Agropyron repens</i>
Cogongrass <i>Imperata cylindrica</i>	Reed, giant <i>Arundo donax</i>
Dallisgrass <i>Paspalum dilatatum</i>	Ryegrass, perennial <i>Lolium perenne</i>
Dandelion <i>Taraxacum officinale</i>	Smartweed, swamp <i>Polygonum coccineum</i>
Dock, curly <i>Rumex crispus</i>	Sweet potato, wild* <i>Ipomoea pandurata</i>
Dogbane, hemp <i>Apocynum cannabinum</i>	Thistle, Canada <i>Cirsium arvense</i>
Fescues <i>Festuca</i> spp	Timothy <i>Phleum pratense</i>
Fescue, tall <i>Festuca arundinacea</i>	Torpedograss* <i>Panicum repens</i>
Guineagrass <i>Panicum maximum</i>	Vaseygrass <i>Paspalum urvillei</i>
Hornnettle <i>Solanum carolinense</i>	Wheatgrass, western <i>Agropyron smithii</i>

*Partial Control

Alfalfa—Apply 1 quart of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. ~~Make application after the last hay cutting in the fall. Allow alfalfa to regrow to a height of 6 to 8 inches or more prior to treatment. Applications should be followed with deep tillage at least 7 days after treatment, but be sure seed is covered up.~~

Alligatorweed—Apply 4 quarts of this product per acre or apply a 1½ percent solution with hand-held equipment to provide partial control. Apply when most of the plants are in bloom. Repeat applications will be required to maintain such control.

~~**Bentgrass**—For suppression in grass seed production areas. For ground applications only, apply 1.5 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 20 gallons of water per acre. Ensure entire crown area has resumed growth prior to a fall application. Bentgrass should be actively growing and have at least 3 inches of growth. Tillage prior to treatment should be avoided. Tillage 7 to 10 days after application is recommended for best results. Failure to use tillage after treatment may result in unacceptable control.~~

Bermudagrass—For control, apply 5 quarts of this product per acre and, for partial control, apply 3 quarts per acre. Treat when bermudagrass is actively growing and seedheads are present. Retreatment may be neces-

sary to maintain control. Allow 7 or more days after application before tillage

Bindweed, field / Bindweed, Texas—Apply 4 to 5 quarts of this product per acre west of the Mississippi River and 3 to 4 quarts per acre east of the Mississippi River. Apply when weed is actively growing and is at or beyond full bloom. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. New leaf development indicates active growth. For best results, apply in late summer or fall. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage

~~**Bindweed, field (suppression-treated lands where annual tillage is performed)**—Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing bindweed that has reached a length of 12 inches or greater. Allow maximum weed emergence and runner growth. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth. Allow 3 or more days after application before tillage~~

Bluegrass, Kentucky / Bromegrass, smooth / Orchardgrass—Apply 2 quarts of this product in 10 to 40 gallons of water per acre when the grasses are actively growing and most plants have reached boot-to-early seedhead stage of development. ~~For partial control in pasture hay and renovation, apply 1 to 1½ quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to actively growing plants when most have reached 4 to 12~~

~~inches in height. Allow 7 or more days after application before tillage~~

~~**Orchardgrass (sods going in no-till corn)**—Apply 1 to 1.5 quarts of this product per acre plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to orchardgrass that is a minimum of 12 inches tall for spring applications and 6 inches tall for fall applications. Allow at least three days following application before planting. A sequential application of 3.75 to 4.5 quarts of Lariat[®] herbicide or equivalent, will be necessary for optimum results. Lariat should be applied within 3 to 10 days following spring applications to prevent annual weed growth.~~

~~Least-toxic registered formulation of Monocrotophos~~

Brackenfern—Apply 3 to 4 quarts of this product per acre as a broadcast spray or as a 1 to 1½ percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 1½ inches long

~~**Burrago, woollyleaf**—For control, apply 2 quarts of this product plus 1 pint of Banvel per acre. For partial control, apply 1 quart of this product plus 1 pint of Banvel per acre. Add 0.5 to 1 percent nonionic surfactant by total spray volume and apply in 20 gallons of water per acre. Apply when plants are producing new active growth which has been initiated by moisture for at least 2 weeks and when plants are at or beyond flowering~~

Canarygrass, reed / Timothy / Wheatgrass, western—Apply 2 to 3 quarts of this product per acre. For best results, apply to actively growing plants when most have

reached the boot-to-head stage of growth. Allow 7 or more days after application before tillage.

Cogongrass—Apply 3 to 5 quarts of this product plus 0.5 to 1 percent nonionic surfactant in 10 to 40 gallons of water per acre. Apply when Cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.

Dogbane, hemp/Knapweed/Horseradish—Apply 4 quarts of this product per acre. Apply when actively growing and when most weeds have reached the late bud to flower stage of growth. Following crop harvest or mowing, allow weeds to regrow to a mature stage prior to treatment. For best results, apply in late summer or fall. Allow 7 or more days after application before tillage.

Fescue, tall—Apply 3 quarts of this product in 10 to 40 gallons of water per acre to actively growing plants when most have reached boot-to-early seedhead stage of development.

~~Fall application only—Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Apply to fescue in the fall when actively growing and plants have 6 to 12 inches of new growth. Allow 7 or more days after application before tillage. A sequential application of 1 pint per acre of this product plus nonionic surfactant will improve long-term control and control seedlings germinating after fall treatments or the following spring.~~

Guineagrass—Apply 3 quarts of this product per acre or use a 1 percent solution with hand-held equipment. Apply to actively growing guineagrass when most has reached at least the 7-leaf stage of growth. Ensure thorough coverage when using hand-held equipment. Allow 7 or more days after application before tillage.

Johnsongrass/Ryegrass, perennial—Apply 1 to 3 quarts of this product per acre in annual cropping systems; apply 1 to 2 quarts of this product per acre. Apply 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre. In noncrop areas where annual tillage and tillage is not performed, apply 2 to 3 quarts of this product in 10 to 40 gallons of water per acre. For best results, apply to actively growing plants when most have reached the boot-to-head stage of growth or in the fall prior to frost. Allow 7 or more days after application before tillage. Do not tank mix with residual herbicides when using the 1 quart per acre rate.

~~For knockdown of Johnsongrass, apply 1 pint per acre plus 0.5 to 1 percent nonionic surfactant in 3 to 10 gallons of water per acre before the plants reach a height of 12 inches. For this use, allow at least 3 days after treatment before tillage.~~

Spot Treatment (partial control or suppression)—Apply a 1 percent solution of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume when Johnson grass is 12 to 18 inches in height. Coverage should be uniform and complete.

Kikuyugrass—Apply 2 to 3 quarts of this product per acre. Spray when most kikuyugrass is at least 8 inches in height (3 or 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.

Lantana—Apply this product as a 1 to 1½ percent solution using hand-held equipment only. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth. Allow 7 or more days after application before tillage.

Milkweed, common—Apply 3 quarts of this product per acre. Apply when actively growing and most of the milkweed has reached the late bud to flower stage of growth. Following small grain harvest or mowing, allow milkweed to regrow to a mature stage prior to treatment. Allow 7 or more days after application before tillage.

~~**Muhly, wirestem**—Apply 1 to 2 quarts of this product per acre. Use 1 quart of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 40 gallons of water per acre. Use 2 quarts of this product when applying 10 to 40 gallons of water per acre or in pasture sod or noncrop areas. Spray when the wirestem muhly is 8 inches or more in height and actively growing. Do not till between harvest and fall applications or in the fall or spring prior to spring applications. Allow 3 or more days after application before tillage. This product will not provide residual control of wirestem muhly from seeds which germinate after application of this product. Do not tank mix with residual herbicides when using the 1 quart per acre rate.~~

~~**Nutsedge, white**—For control, apply 3 quarts of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. Applications should be made when at least 60 percent of the plants have berries. Fall treatments must be applied before a killing frost. Allow 7 or more days after application before tillage. Do not treat when weed is under drought stress as good soil moisture is necessary for active growth.~~

Nutsedge, purple, yellow—Apply 3 quarts of this product per acre as a broadcast spray or apply a 1 percent solution from hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Treat when plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control. Wait 7 days after treatment before tillage. Tillage will stimulate nutlet germination.

~~Apply 1 quart to 1 quart of this product per acre plus 0.5 to 1 percent nonionic surfactant in 3 to 40 gallons of water per acre to provide suppression or partial control of existing nutsedge plants. Treat when plants have 3 to 5 leaves (less than 6 inches). Repeat treatments will be required to control subsequent emerging plants. Wait 7 days after treatment before tillage or mowing.~~

Pampas—Apply this product as a 1½ to 2 percent solution using hand-held equipment. Apply to plants that are actively growing at or beyond the boot stage of growth. Thorough coverage is necessary for best control.

Phragmites—For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 5 quarts per acre as a broadcast spray or apply a 2 percent solution from hand-held equipment. In other areas of the U.S., apply 3 quarts per acre as a broadcast spray or apply a 1 percent solution from hand-held equipment for partial control. For best results, treat during late summer or fall months or when plants are actively growing and in full bloom. Treatment before or after this stage may lead to reduced control. Due to the dense nature of the vegetation, which may prevent good spray coverage or uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.

~~**Quackgrass**—In Annual Cropping Systems or in Pastures and Soils Followed by Deep Tillage: Apply 1 to 2 quarts of this product per acre. For the one quart rate, apply 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre. For the 2 quart rate, apply in 10 to 40 gallons of water per acre. Do not tank mix with residual herbicides when using the 1 quart rate. Spray when quackgrass is 8 to 12 inches in height and actively growing. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage. In pastures or soils, for best results use a moldboard plow.~~

Quackgrass—Pasture or Soil or Other Noncrop Areas Where Deep Tillage is Not Planned Following Application: Apply 2 to 3 quarts in 10 to 40 gallons of water per acre. Spray when the quackgrass is greater than 8 inches tall

and actively growing. Do not till between harvest and fall applications or in fall or spring prior to spring application. Allow 3 or more days after application before tillage.

Reed, giant—For control of giant reed, apply a 2 percent solution of this product when plants are actively growing. Best results are obtained when applications are made in late summer to fall.

Sweet Potato, wild—Apply this product as a 2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment. Allow 7 or more days before tillage.

Thistle, Canada—Apply 2 to 3 quarts of this product per acre. Apply to actively growing thistles when most are at or beyond the bud stage of growth. After harvest, mowing or tillage in the late summer or fall allow at least 4 weeks for initiation of active growth and rosette development prior to the application of this product. Fall treatments must be applied before a killing frost. Allow 3 or more days after application before tillage.

~~For suppression of Canada thistle, apply 1 quart per acre of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 3 to 10 gallons of water per acre in the late summer or fall after harvest, mowing or tillage. Allow rosette regrowth to a minimum of 6 inches in diameter before treating. Applications can be made as long as leaves are still green and plants are actively growing at the time of application. Allow 3 or more days after application before tillage.~~

10

Torpedograss—Apply 4 to 5 quarts of this product per acre to provide partial control of torpedograss. Apply to actively growing torpedograss when most plants are at or beyond the seedhead stage of growth. Repeat applications will be required to maintain control. Fall treatments must be applied before frost. Allow 7 or more days after application before tillage.

Other perennials listed on this label—Apply 3 to 5 quarts of this product per acre. Apply when actively growing and most have reached early head or early bud stage of growth. Allow 7 or more days after application before tillage.

Accord (R) plus Oust (TM)
For Industrial Sites

Use on industrial sites including airports, industrial plants, lumberyards, petroleum tank farms, pumping stations, pipelines, railroads, roadsides, storage areas or other similar sites where bare ground is desired.

When applied as directed under the conditions described, this product plus Oust provides control of annual weeds listed in the "Weeds Controlled" section of the label for this product and Oust, and control or partial control of the perennial weeds listed below.

Apply 1 to 2 quarts of this product with 2 to 4 ounces of Oust in 10 to 40 gallons of spray solution per acre as a broadcast spray to actively growing weeds.

For control of annual weeds, use the lower rates of these products.

For control of the listed perennial weeds, use the higher rates of both products. For partial control, use the lower rates.

Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Bahugrass <i>Paspalum notatum</i>	Johnsongrass** <i>Sorghum halepense</i>
Bermudagrass* <i>Cynodon dactylon</i>	Peonies** <i>Diodia teres</i>
Broomsedge <i>Andropogon virginicus</i>	Quackgrass <i>Agropyron repens</i>
Dock, curly <i>Rumex crispus</i>	Trumpetcrasper* <i>Campsis radicans</i>
Dogfennel <i>Eupatorium capilliflorum</i>	Vaseygrass <i>Paspalum urvillei</i>
Fescue, tall <i>Festuca arundinacea</i>	Vervain, blue <i>Verbena hastata</i>

*Suppression at the higher rates only

**Control at the lower rates

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used

*Oust is a trademark of E. I. du Pont de Nemours and Company

Tank Mixtures For Industrial Sites

When applied as a tank mixture, this product provides control of the emerged annual weeds and partial control of the emerged perennial weeds listed in this label. When applied as a tank mixture, the following residual herbicides will provide preemergence control of the weeds listed in the individual product labels.

Accord plus Diuron

Accord plus Krovar (TM) II

Accord plus Krovar I

Accord plus Simazine, Princep Caliber (TM) 90

Accord plus Simazine 4L

Accord plus Simazine 80 W

Accord plus Surflan (TM) 75 W

Accord plus Surflan AS

Read and carefully observe the label claims, cautionary statements, recommended use rates and all other information on the labels of all products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Control of Emerged Weeds

Annual Weeds-Apply 1 quart per acre of this product in these tank mixtures when weeds are less than 6 inches tall and 1 1/2 quarts per acre when weeds are more than 6 inches tall.

Perennial Weeds-For partial control of perennial weeds using these tank mixtures, apply 2 to 5 quarts per acre of this product. Follow the recommendations in the "Weeds Controlled" section of this label for the stage of growth and rate of application for specific perennial weeds.

Preemergence Weed Control

The following are the labeled rates for the recommended residual herbicides for noncrop uses:

PRODUCT	RATE/ACRE
Diuron, Karmex (TM) 80DF,	2 to 8 lbs
Krovar I	4 to 6 lbs
Krovar II	2 to 5 lbs
Simazine, Princep Caliber 90	5.3 to 11.1 lbs
Simazine 80 W	6 to 12 1/2 lbs
Simazine 4L	4.8 to 10 qts
Surflan 75 W	2 2/3 to 5 1/3 lbs
Surflan AS	2 to 4 qts

Refer to the individual product labels for specific noncrop sites, rates, carrier volumes, and precautionary statements.

Mix only the quantity of spray solution which can be used during the same day. Do not allow these tank mixtures to stand overnight as this may result in reduced weed control.

APPLY THESE TANK MIXTURES THROUGH CONVENTIONAL BROADCAST EQUIPMENT ONLY.

TM Caliber is a trademark of Ciba-Geigy Corporation.

TM Karmex and TM Krovar are trademarks of E.I. du Pont de Nemours and Company.

TM Surflan is a trademark of Elanco Products Company

RELEASE OF BERMUDAGRASS OR BAHIA GRASS

NOTE: Use only in areas where bermudagrass or bahia grass are desirable ground covers and where some temporary injury or discoloration can be tolerated. Use Roundup plus Oust tank mixtures only on railroads, highways, utility plant sites, or other right-of-way areas.

When applied as directed for ~~Nonemop~~ uses under the conditions described, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahia grass. This product may be tank-mixed with Oust as recommended for residual control. Make applications to dormant bermudagrass or bahia grass. Roundup plus Oust tank mixture may delay greenup. To avoid delays in greenup and minimize injury, do not add more than 1 ounce per acre of Oust on bermudagrass or more than 1/2 ounce per acre on bahia grass, or treat when these grasses are in a semi-dormant condition.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 - 6 leaf stage.

79

80

WEEDS CONTROLLED

Rate recommendations for control or suppression of winter annuals and tall fescue are listed below.

Apply the recommended rates of this product alone or as a tank mixture in 10 to 25 gallons of water plus 0.5 to 1 percent nonionic surfactant by total spray volume per acre.

For the best recommendation for the mixture of weeds within your geographic areas, contact your Monsanto sales representative.

WEEDS CONTROLLED OR SUPPRESSED WITH ROUNDUP® ALONE*

NOTE: C = Control
S = Suppression

WEED SPECIES	ROUNDUP® OZ/ACRE					
	8	12	16	24	32	64
Barley, little <i>Hordeum pusillum</i>	S	C	C	C	C	C
Bodotrow, catchweed <i>Galium aparine</i>	S	C	C	C	C	C
Bluegrass, annual <i>Poa annua</i>	S	C	C	C	C	C
Chenop <i>Chaerophyllum tainturieri</i>	S	C	C	C	C	C

← Accord

← Delete

← Accord

← Accord

← Accord

Chickweed, common <i>Stellaria media</i>	S	C	C	C	C	C
Clover, crimson <i>Trifolium incarnatum</i>	•	S	S	C	C	C
Clover, large leaf <i>Trifolium campestre</i>	•	S	S	C	C	C
Fescue, tall <i>Festuca arundinaceae</i>	•	•	•	•	S	S
Geranium, Carolina <i>Geranium carolinianum</i>	•	•	S	S	C	C
Hemlock <i>Lamium amplexicaule</i>	•	S	C	C	C	C
Ryegrass Italian <i>Lolium multiflorum</i>	•	•	S	C	C	C
Speedwell, corn <i>Veronica arvensis</i>	S	C	C	C	C	C
Vetch, common <i>Vicia sativa</i>	•	•	S	C	C	C

*These rates apply only to sites where an established competitive turf is present.

81
82

**WEEDS CONTROLLED OR SUPPRESSED
WITH ROUNDUP® PLUS OUST***

NOTE C = Control
S = Suppression

WEED SPECIES	ROUNDUP® + OUST (OZ/ACRE)								
	ROUNDUP	8	12	12	16	12	16		
	+	+	+	+	+	+	+		
	OUST	1/4	1/4	1/4	1/4	1/4	1	1	
Barley, little <i>Hordeum pusillum</i>		C	C	C	C	C	C	C	
Bedstraw, catchweed <i>Galium aparine</i>		C	C	C	C	C	C	C	
Bleeding heart <i>Poa annua</i>		S	C	C	C	C	C	C	
Chenopod <i>Chaerophyllum tarnifolium</i>		C	C	C	C	C	C	C	
Chickweed, common <i>Stellaria media</i>		S	C	C	C	C	C	C	
Clover, crimson <i>Trifolium incarnatum</i>		S	S	S	S	C	C	C	
Clover, large leaf <i>Trifolium campestre</i>		•	•	S	S	S	C	C	
Fescue, tall <i>Festuca arundinaceae</i>		•	•	•	•	•	S	S	
Geranium, Carolina <i>Geranium carolinianum</i>		•	S	S	C	C	C	C	

← Accord

← Accord

Hemlock	• S C C C C C
<i>Larix laricina</i>	
Ryegrass, Italian	• S S C C C C
<i>Lolium multiflorum</i>	
Speedwell, corn	S C C C C C C
<i>Veronica arvensis</i>	
Vetch, common	C C C C C C C
<i>Vicia sativa</i>	

*These rates or mixtures of rates apply only to sites where an established competitive turf is present.

RELEASE OF ACTIVELY GROWING BERMUDAGRASS

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "Weeds Controlled" section of this and the Oust label, and suppression or partial control of certain perennial weeds.

For control or suppression of those annual species listed on this label, use 1 to 3 pints of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use higher rate as size of plants increases or as they approach flower or seedhead formation.

Use the higher rate of this product for partial control of the following perennial species. Use the lower rates for suppression of growth. For best results, see the "Weeds Controlled" section of this label for proper stage of growth.

83



84

Bahiagrass	Johnsongrass**
<i>Paspalum notatum</i>	<i>Sorghum halepense</i>
Bloodroot, silver	Trumpetcrimper*
<i>Andropogon saccharoides</i>	<i>Campsis radicans</i>
Fescue, tall	Vaseygrass
<i>Festuca arundinacea</i>	<i>Paspalum urvillei</i>

*Suppression at higher rates only

**Control at the higher rates

This product may be tank-mixed with Oust. If tank-mixed, use no more than 1 to 2 pints per acre of Roundup herbicide with 1 to 2 ounces of Oust per acre.

← Accord

Use the lower rates of both mixtures to control annual weeds below 6 inches in height (or runner length in annual vines) that are listed in the "Weeds Controlled" section of this booklet and the Oust label. Use the higher rates as annual weeds increase in size and approach the flower or seedhead stages.

Use the higher rates of this product to provide partial control of the following perennial weeds. Use the lower rates for suppression of growth.

Bahiagrass	Johnsongrass*
<i>Paspalum notatum</i>	<i>Sorghum halepense</i>
Bloodroot, silver	Peonies**
<i>Andropogon saccharoides</i>	<i>Diodia teres</i>
Broomrape	Trumpetcrimper*
<i>Andropogon virginicus</i>	<i>Campsis radicans</i>
Oach, curly	Vaseygrass
<i>Rumex crispus</i>	<i>Paspalum urvillei</i>

Dogfennel **Vervain, blue**
Eupatorium capilliflorum *Verbena hastata*

Fescue, tall
Festuca arundinacea

*Suppression at higher rates only

**Control at the higher rates

Use only on well-established bermudagrass. Bermuda-grass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may result.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

COOL SEASON TURF GROWTH REGULATION

When applied as directed, this product will suppress growth and seedhead development of listed turf species in industrial sites.

This product is recommended for management of coarse turfs on roadside rights-of-way or other industrial areas. Do not use on high-quality turfs or other areas where some turf color changes cannot be tolerated. Slight turf discoloration may occur but turf will regreen and regrow under moist conditions as effects of this product wear off.

85

86

Apply 4 to 6 fluid ounces of this product per acre alone or in a recommended tank mixture. Spray volumes of 10 to 40 gallons per acre are recommended.

When using this product, mix 2 quarts of a nonionic surfactant per 100 gallons of spray solution.

This product can be used for growth and seedhead suppression of

Tall Fescue
Smooth Brome

For best results, apply this product in a recommended tank mixture to actively growing turfgrasses after greenup in the spring of the year. For suppression of seedheads, applications must be made before boot-to-seedhead stage of development. Applications made from seedhead emergence until maturity may result in turf discoloration or injury.

After mowing or removal of seedheads, this product in a recommended tank mixture may also be used to suppress the growth of certain turfgrasses. Allow turf to recover from stress caused by heat, drought, or mowing before making applications. Applications made to turf under stress may increase the potential for discoloration or injury.

TANK MIXTURES

For the following tank mixtures, consult each product label for weeds controlled and the correct stage of application. Do not treat turf under stress.

Tank mixtures plus 2,4-D Amine

For additional weed control benefits, up to 1 quart per acre of 2,4-D amine may be added to the following tank mixtures. Consult the label for 2,4-D amine for weeds controlled.

TALL FESCUE

~~Roundup plus Telar™~~

Accord

For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 1/2 ounce of Telar per acre.

This tank mixture can also be applied after mowing or removal of tall fescue seedheads for turf growth suppression. Make only one of the above applications per growing season.

~~Roundup plus Oust™~~

Accord

For suppression of tall fescue growth and seedheads, and control or partial control of some annual weeds, apply

87

88

this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 1/4 ounce of Oust per acre.

SMOOTH BROME

~~Roundup plus Oust~~

Accord

For suppression of smooth brome growth and seedheads and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 1/4 ounce of Oust per acre.

Insert:

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

BAHIA GRASS SEEDHEAD AND VEGETATIVE SUPPRESSION

When applied as directed in the indicated noncrop areas (roadsides, airports, golf course roughs, and plant sites), this product will provide significant inhibition of seed-head emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications.

(TM) Oust and Telar are trademarks of E.I. du Pont de Nemours and Company

Apply this product 1 to 2 weeks after full greenup of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 6 fluid ounces per acre of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume in 10 to 25 gallons of water per acre.

Sequential applications of this product plus 0.5 to 1 percent nonionic surfactant by total spray volume may be made at approximately 45 day intervals to extend the period of seedhead and vegetative growth suppression. For continued seedhead suppression, sequential applications must be made prior to seedhead emergence. Apply no more than 2 sequential applications per year. As a first sequential application, apply 4 fluid ounces of this product per acre plus nonionic surfactant. A second sequential application of 2 to 4 fluid ounces per acre plus nonionic surfactant may be made approximately 45 days after the last application.

A tank mixture of Roundup plus Oust may be applied only on roadsides for seedhead inhibition and vegetative suppression. Apply 6 fluid ounces per acre of this product plus 0.25 ounce per acre of Oust, plus 0.5 to 1 percent nonionic surfactant by total spray volume 1 to 2 weeks following an initial spring mowing. When using Roundup plus Oust for suppression of bahiagrass, make only one application per year.

← Accord

← Accord

Read and carefully observe the label claims, cautionary statements and all information on the labels of each product used. Use according to the most restrictive label directions for each product used in recommended tank mixtures.

When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" i.e. the label booklet for Accord before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

(C) MONSANTO COMPANY 1990

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167