

PM 25

524-326

1 of 37

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ACCORD® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Accord herbicide and this supplement.

ACCORD HERBICIDE

BY MONSANTO

EPA Reg. No. 524-326

BASAL AND DORMANT STEM APPLICATIONS OF ACCORD HERBICIDE FOR CONTROL OR PARTIAL CONTROL OF WOODY TREES AND BRUSH

REC'D EPA/PP/DPDI

'97 JAN 30 1 13 PM

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, call Collect, day or night, 314-694-4000.

Accord® is a registered trademark of Monsanto Company.

Directions for Use

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

When applied as directed, this product will control or partially control woody brush and trees in areas such as utility rights-of-way, forestry sites, roadsides, railroads, drainage ditches and canals, fence rows, and similar non-crop areas.

This application technique is recommended for directed spot treatment of trees and brush. Apply this product using backpack, hand-held, handgun or similar equipment. Use flat fan, cone, or similar nozzles that will provide effective spray coverage of target vegetation. These applications are more effective on small brush less than 15 feet in height or 3-inch stem diameter. Use the higher rates for difficult to control species, trees with bigger stems, brush growing in clusters, or species growing under stressful environmental conditions.

If needed, use propylene glycol to prevent spray solution from freezing. Refer to the product label for proper application rate and use instructions.

BEST AVAILABLE COPY

BASAL APPLICATION

Apply a solution consisting of 25-50% v/v of this product and up to 50% v/v of MON 59120 spray adjuvant. Fill spray tank or container with clean water to the desired volume if necessary. Completely cover the lower 18-24 inches of the brush stems or trunks. For larger stems over 3 inches in diameter, use the higher rates and treat up to 48 inches or higher from the ground level. Treat every stem of brush existing in clusters. Spray-to-wet to the point of run-off.


DORMANT STEM APPLICATION

Apply a solution consisting of 10-25% v/v of this product and up to 50% v/v of MON 59120 spray adjuvant. Fill spray tank or container with clean water to the desired volume if necessary. Apply from fall to spring. Spray coverage should be uniform, covering at least 75% of the trunks, branches, and stems of the target brush or trees.

Read and carefully observe the label claims, cautionary statements, and all information on the labels of all products used in this tank mixture.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Accord before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

©MONSANTO COMPANY 1997
MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167



BEST AVAILABLE COPY

ACCORD®
Herbicide by Monsanto

Complete Directions for Use
in Forestry and Utility Rights-of-Way

EPA Reg. No. 524-326

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NONWOODY ROOTS,
OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY
OR DESTRUCTION WILL RESULT.

® Accord is a registered trademark of Monsanto Company.

1997-1

892.65-000.37/53

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or
using. If terms are not acceptable, return at once unopened.

REFORMULATION IS PROHIBITED. SEE CONTAINER LABEL FOR REPACKAGING
LIMITATIONS.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical
description on the label and is reasonably fit for the purposes
set forth in the Complete Directions for Use label booklet
("Directions") when used in accordance with those Directions
under the conditions described therein. NO OTHER EXPRESS WARRANTY
OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR
MERCHANTABILITY IS MADE. This warranty is also subject to the
conditions and limitations stated herein.

(ACCORD 96-1)

ACCEPTED
MAR - 4 1997
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 524-326

(1/16/97-19/96)

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

Hazards to
Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

HARMFUL IF INHALED.

Avoid breathing vapors or spray mist.

FIRST AID: IF INHALED, remove individual to fresh air. Seek medical attention if breathing difficulty develops.

Personal Protective Equipment

Applicators and other handlers must wear long-sleeved shirt and long pants and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendation

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not contaminate water when disposing of equipment washwaters. Treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss can cause fish suffocation.

In case of:

SPILL or LEAK, soak up and remove to a landfill.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt

41.5%

INERT INGREDIENTS:

58.5%

100.0%

*Contains 480 grams per litre or 4 pounds per U.S. gallon of glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt. Equivalent to 356 grams per litre or 3 pounds per U.S. gallon of the acid, glyphosate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers

on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are: coveralls, waterproof gloves and shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Uses of this product in utility rights-of-way and all other utility sites are not within the scope of the Worker Protection Standard requirements. Requirements in the Agricultural Use Requirement box do not apply to utility uses. Follow all other label requirements for applications to utility sites.

For more product information, call toll free 1-800-332-3111.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: STORE ABOVE 10°F (-12°C) TO KEEP PRODUCT FROM CRYSTALLIZING.

Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F (20°C) for several days to redissolve and roll or shake container or recirculate in mini-bulk containers to mix well before using. For bulk containers, see container label.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

(See the individual container labels for disposal information)

GENERAL INFORMATION

This product, a water soluble liquid, mixes readily with water and surfactant to be applied as a foliar spray for the control or destruction of most herbaceous and woody plants.

This product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most herbaceous weeds occur within 7 days but on most woody plants may not occur for 30 days or more.

After any site disturbance, such as logging, mechanical brush removal or mowing, allow stump sprouts, resprouts and foliar regrowth from woody brush and perennial herbaceous weeds sufficient time to regrow before treatment.

Always use the higher recommended rates of this product and surfactant when treating dense, multicanopied sites of woody vegetation or difficult-to-control woody and herbaceous plants.

Reduced control may result when woody brush, trees and herbaceous weeds are treated under poor growing conditions caused by drought, disease or insect damage. Reduced control may result if the foliage of undesirable vegetation is covered with dust at the time of treatment.

Rainfall occurring within 6 hours after application may reduce effectiveness. Heavy rainfall within 2 hours after application may wash the chemical off the foliage and a repeat treatment may be required.

When this product comes in contact with soil (on the soil surface or as suspended soil or sediment in water) it is bound to soil particles. When used in accordance with label directions, once this product is bound it is not available for plant uptake and will not harm off-site vegetation where roots grow into the treatment area or if the soil is transported off-site. When used in accordance with label directions, the strong affinity of this product to soil particles prevents this product from leaching out of the soil profile and entering ground water. The affinity between this product and soil particles remains until this product is degraded, which is primarily a biological degradation process carried out under both aerobic and anaerobic conditions by soil microflora.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this label. Mixing this product with herbicides or other materials not recommended in this label may result in reduced performance.

**FORESTRY SITE PREPARATION
AND UTILITY RIGHTS-OF-WAY**

This product is recommended for the control or partial control of woody brush, trees and herbaceous weeds. This product is labeled for use in forestry and utility sites. This product is also recommended for use in preparing or establishing wildlife openings

10/2/37

within these sites and maintaining logging roads, and for side trimming along utility rights-of-way.

In forestry, this product is recommended for use in site preparation prior to planting any tree species, including Christmas trees and silvicultural nursery sites.

In utilities, this product is recommended for use along electrical power, pipeline and telephone rights-of-way, and in other utility sites associated with these rights-of-way, such as substations.

APPLICATION RATES AND TIMING

APPLICATION	ACCORD®	SPRAY VOLUME GAL/A
<u>BROADCAST</u>		
Aerial	2 to 10 qts/a	5 to 30
Ground	2 to 10 qts/a	10 to 60
<u>SPRAY-TO-WET</u>		
Handgun, Backpack, Mistblower	3/4 % to 2 % by volume	spray-to-wet
<u>LOW VOLUME DIRECTED SPRAY</u>		
Handgun, Backpack, Mistblower	5% to 10% by volume	partial coverage*

*For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results.

In forestry site preparation and utility rights-of-way applications, this product requires use with a nonionic surfactant. Use a nonionic surfactant with greater than 80 percent active ingredient and labeled for use with herbicides. Use of this product without surfactant will result in reduced

performance. See the "MIXING AND APPLICATION INSTRUCTION" section of this label for more information.

Mix 2 or more quarts of the nonionic surfactant per 100 gallons of spray solution (0.5 percent or more by spray volume). Use of surfactant concentrations greater than 1.5 percent by spray volume with handgun applications or 2.5 percent by spray volume with broadcast applications is not recommended.

Use higher rates of this product within the recommended range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the recommended range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the recommended range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

This product has no herbicidal or residual activity in the soil. Where repeat applications are necessary, do not exceed 10.6 quarts of this product per acre per year.

TANK MIXTURES

Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of both products used. Use according to the most restrictive precautionary statements for each product in the mixture. Any recommended rate of this product may be used in a tank mix.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions. For side trimming

treatments in utility rights-of-way, tank mixtures with Arsenal™ 2WSL are not recommended. For side trimming treatments, it is recommended that this product be used alone as recommended, or as a tank mixture with Garlon™ 4.

PRODUCT	BROADCAST RATE	USE SITE
Arsenal Applicators Concentrate	2 to 16 fl oz /a	Forestry site preparation
Oust™	1 to 4 oz/a	Forestry site preparation, Utility sites
Garlon 3A*, Garlon 4	1 to 4 qts/a	Forestry site preparation, Utility sites
Arsenal 2WSL	2 to 32 fl oz/a	Utility sites

PRODUCT	SPRAY-TO-WET RATES	USE SITE
Arsenal Applicators Concentrate	1/32 % to 1/2 % by volume	Forestry site preparation
Arsenal 2WSL	1/32 % to 1/2 % by volume	Utility sites

PRODUCT	LOW VOLUME DIRECTED SPRAY RATES	USE SITES
Arsenal Applicators Concentrate	1/8% to 1/2% by volume	Forestry site preparation
Arsenal 2WSL	1/8% to 1/2% by volume	Utility sites

* Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray compatibility problems.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

*Arsenal is a trademark of American Cyanamid Company.

*Oust is a trademark of E. I. du Pont de Nemours and Company.

*Garlon is a trademark of DowElanco Products Company.

FORESTRY CONIFER AND HARDWOOD RELEASE

DIRECTED SPRAY AND SELECTIVE EQUIPMENT

This product may be applied as a directed spray or by using selective equipment in forestry conifer and hardwood sites, including Christmas tree plantations and silvicultural nurseries. Mix 2 to 6 quarts of a nonionic surfactant per 100 gallons of spray solution (0.5 to 1.5 percent by spray volume) for all spray applications. Use a surfactant with greater than 80 percent active ingredient.

In hardwood plantations, tank mixtures with Oust may be used. In pine plantations, tank mixtures with Garlon 4 or Arsenal AC may be used. Comply with all site restrictions, forestry species limitations and precautions on the tank mix product label.

Avoid contact of spray, drift, mist or drips with foliage, green bark or non-woody surface roots of desirable species.

See all sections in the APPLICATION EQUIPMENT AND TECHNIQUES portion of this label for specific equipment recommendations and precautions.

For spray-to-wet applications, use a 2 percent spray solution for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent solution.

For low volume directed spray applications, use a 5 to 10 percent spray solution. Coverage should be uniform with at least 50

percent of the foliage contacted. Coverage of the top one-half of the unwanted vegetation is important.

For equipment calibrated for broadcast applications, use 2 to 10 quarts of this product per acre. Apply in 10 to 60 gallons of clean water per acre. Sheilded application equipment may be used to avoid contact of the spray solution with desirable plants. Sheilds should be adjusted to prevent spray contact with the foliage or green bark of desirable vegetation.

Wiper application equipment may be used. See the SELECTIVE EQUIPMENT portion of this label for equipment and rate recommendations

BROADCAST SPRAY

Except where specifically recommended below, use only where conifers have been established for more than one year.

APPLICATION MUST BE MADE AFTER FORMATION OF FINAL CONIFER RESTING BUDS IN THE FALL OR PRIOR TO INITIAL BUD SWELLING IN THE SPRING.

Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied. Damage can be accentuated if applications are made when conifers are actively growing, or are under stress from drought, flood water, improper planting, insects, animal damage or diseases.

This product may require use with a surfactant. Unless otherwise recommended in this section of this label, use Entry[™] II surfactant at 10 to 30 fluid ounces per acre. Follow the instructions under the "Mixing" portion of the "MIXING AND APPLICATION INSTRUCTIONS" section of this label.

For release of the following conifer species outside the Southeastern United States:

- Douglas fir
- Pseudotsuga menziesii*

15 of 31

Fir

Abies spp.

Hemlock**

Tsuga spp.

Pines*

Pinus spp.

Redwood, California**

Sequoia spp.

Spruce

Picea spp.

* Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.

**Use of a surfactant is not recommended for release of hemlock species or California redwood. In mixed conifer stands, injury to these species may result if a surfactant is used.

Apply 1 to 2 quarts of this product per acre as a broadcast spray.

NOTE: For release of Douglas fir with this product or recommended tank mixtures of this product, Entry II or a nonionic surfactant recommended for over-the-top foliar sprays may be used. To avoid possible conifer injury, Entry II rates should not exceed 20 fluid ounces per acre at elevations above 1500 feet, or 10 fluid ounces per acre in the coastal range or at elevations below 1500 feet in Washington and Oregon. Nonionic surfactants may be used at 2 fluid ounces per acre at elevations above 1500 feet, or 1 fluid ounce per acre in the coastal range or at elevations below 1500 feet. Use of surfactant rates exceeding those listed above may result in unacceptable conifer injury and are not recommended. Ensure that the nonionic surfactant has been adequately tested for Douglas fir safety before use.

In Maine, up to 3 quarts per acre of this product may be used for the control of difficult species.

160/31

To release Douglas fir, and pine and spruce species at the end of the first growing season(except in California), apply 1 to 1.5 quarts of this product per acre. Ensure that the conifers are well hardened off.

OUST TANK MIXTURES - To release jack pine, white pine and white spruce, apply 1 to 2 quarts of this product with 1 to 3 ounces(1 to 1.5 for white pine) of Oust per acre. Make applications to actively growing weeds as a broadcast spray over the top of established conifers. Applications at these rates should be made after formation of conifer resting buds in the late summer or fall.

ARSENAL APPLICATORS CONCENTRATE TANK MIXTURES - This product may be tank mixed with Arsenal Applicators Concentrate for release of Douglas fir. Use 1 to 1 1/2 quarts of this product tank mixed with 2 to 6 fluid ounces of Arsenal per acre. For release of balsam fir and red spruce, apply a mixture of 2 quarts of this product and 1 to 2 1/2 fluid ounces of Arsenal Applicators Concentrate per acre.

For release of the following conifer species in the Southeastern United States:

Loblolly pine
Pinus taeda

Eastern white pine
Pinus strobus

Shortleaf pine
Pinus echinata

Slash pine
Pinus elliottii

Virginia pine
Pinus virginiana

Longleaf pine

Pinus palustris

Apply 1 1/2 to 2 1/2 quarts of this product per acre as a broadcast spray during late summer or early fall after the conifers have hardened off. For applications at the end of the first growing season, use 1 quart per acre of this product alone or in a recommended tank mixture.

ARSENAL APPLICATORS CONCENTRATE TANK MIXTURES - Apply 1 to 2 quarts of this product with 2 to 16 fluid ounces of Arsenal Applicators Concentrate per acre as a broadcast spray for conifer release. Use only on conifer species that are labeled for over-the-top sprays for both products. Use the higher recommended rates for dense, tough-to-control woody brush and trees.

Read and carefully observe the label claims, cautionary statements and all information on the labels of each product used in these tank mixtures. Use according to the most restrictive precautionary statements for each product in the mixture.

*Entry is a trademark of Monsanto Company.

HERBACEOUS RELEASE

When applied as directed, this product plus listed residual herbicides provides postemergence control of the annual weeds and control or suppression of the perennial weeds listed in this label, and residual control of the weeds listed in the residual herbicide label. Make applications to actively growing weeds as a broadcast spray over the top of labeled conifers.

Oust tank mixtures - To release loblolly pines, apply 16 to 24 fluid ounces of this product, plus 2 to 4 ounces of Oust per acre. To release slash pines, apply 12 to 16 fluid ounces of this product, plus 2 to 4 ounces of Oust per acre.

Mix up to 3.2 fluid ounces per acre of Entry II with the recommended rate of this product plus Oust. Applications can be made over newly planted pines after the emergence of herbaceous weeds in the spring or early summer. Best results are obtained from applications made in May and June.

18 of 31

Weed control may be reduced if water volumes exceed 25 gallons per acre for these treatments.

Atrazine tank mixtures - To release Douglas fir, apply 1 quart of this product, plus 4 pounds a.i. of atrazine per acre. Apply only over Douglas fir that has been established for at least one full growing season. Apply in the early Spring, usually mid-March through early April. Injury will occur if applications are made after bud swell in the Spring. Do not add surfactant to this mix for this use.

Always read and follow the manufacturer's label recommendations for all herbicides and surfactants used.

WETLAND SITES

This product may be used in and around water (aquatic areas) and wetlands found in forestry and in power, telephone and pipeline rights-of-way sites, including where these sites are adjacent to or surrounding domestic water supply reservoirs, supply streams, lakes and ponds. Read and observe the following before making applications in and around water.

Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat in such areas.

There is no restriction on the use of treated water for irrigation, recreation or domestic purposes.

NOTE: Do not apply this product directly to water within 1/2 mile up-stream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. These aquatic applications may be made **ONLY** in those cases where there are alternative water sources

or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications.

This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.

Do not spray open bodies of water where woody brush, trees and herbaceous weeds do not exist. The maximum application rate of 5 quarts per acre must not be exceeded in a single over-water broadcast application except as follows, where any recommended rate may be applied:

- Stream crossings in utility rights-of-way
- Where applications will result in less than 20 percent of the total water area being treated.

**MIXING AND APPLICATION
INSTRUCTIONS**

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES. HANDGUN APPLICATIONS SHOULD BE PROPERLY DIRECTED TO AVOID SPRAYING DESIRABLE PLANTS. NOTE: REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

MIXING

This product mixes readily with water. Mix spray solutions of this product as follows: Fill the mixing or spray tank with the required amount of water while adding the required amount of this product (see the "DIRECTIONS FOR USE" and "WEEDS CONTROLLED" sections of this label). For tank mixtures, add the tank-mix product before adding this product. If tank mixing with Garlon 3A, ensure that the Garlon 3A is well mixed with at least 75 percent of the total spray volume before adding this product to avoid incompatibility. Near the end of the filling process, add the required surfactant and mix well. Maintain an air break between the filling hose and the spray solution and remove the

hose from the tank immediately after filling to avoid siphoning back into the water source. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, terminate by-pass and return lines at the bottom of the tank and, if needed, use an approved antifoam or defoaming agent.

**APPLICATION EQUIPMENT
AND TECHNIQUES**

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of plant or crop injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

■ **AERIAL EQUIPMENT** ■

This product is recommended for application by helicopter only in forestry sites and utility rights-of-way. Use the recommended rates of this product and surfactant in 5 to 30 gallons of spray

solution per acre as a broadcast spray. (THIS PRODUCT PLUS OUST TANK MIXTURES MAY NOT BE APPLIED BY AIR IN CALIFORNIA.)

IN CALIFORNIA, AERIAL APPLICATION MAY ONLY BE MADE IN NONRESIDENTIAL, FORESTRY SITES AND CHAPARRAL AREAS.

AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITION, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT; DRIFT WILL CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT IS NOT INTENDED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine droplets.

Drift control additives may be used for forestry site preparation and utility rights-of-way applications. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label. The use of a drift control agent for conifer and herbaceous release applications may result in conifer injury and is not recommended.

PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion. To prevent corrosion of exposed parts, thoroughly wash aircraft after each day of spraying to remove residues of this product accumulated during spraying or from spills. Landing gear are most susceptible.

■ GROUND BROADCAST EQUIPMENT ■

This product is recommended for broadcast applications using suitable ground equipment in forestry sites, utility sites and utility rights-of-way. Use the recommended rates of this product plus surfactant in 10 to 60 gallons of clean water per acre as a broadcast spray. Check for even spray distribution throughout the spray pattern.

BACKPACK, HANDGUN OR
MISTBLOWER EQUIPMENT

This product is recommended for application through backpack, handgun or hand-held mistblower* equipment. For spray-to-wet applications, coverage should be uniform and complete, but not to the point of runoff.

This product can be used for low volume directed sprays for spot treatment of trees and brush. It is most effective in areas where there is a low density of undesirable trees or brush. If a straight stream nozzle is used, start the application at the top of the targeted vegetation and spray from top to bottom in a lateral zigzag motion. For flat fan and cone nozzles and with mistblowers, mist the application over the foliage of the targeted vegetation. Small, open branched trees need only be treated from one side. If the foliage is thick or there are multiple root sprouts, applications must be made from several sides to ensure adequate spray coverage.

It is suggested that the recommended amount of this product and surfactant be mixed in a larger container and then added to the sprayer.

*This product is not registered in California or Arizona for use in mistblowers.

SELECTIVE EQUIPMENT

This product may be applied through shielded sprayers or wiper application equipment. This equipment may be used to selectively control undesirable vegetation without harming desirable vegetation.

Shielded sprayers direct the herbicide solution onto weeds while shielding desirable vegetation from the spray solution. Any recommended rate or tank mixture of this product may be used employing this equipment.

23 of 31

Wiper applicators physically wipe product directly onto undesirable vegetation. Care should be taken to avoid wiping desirable vegetation. Use a 33 to 100 percent solution of this product, diluted in water for wiper applications. Use a 33 percent solution for wick or gravity feed systems. Higher concentrations may be used in pressurized systems that are capable of handling thicker solutions. Addition of a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution is recommended.

WEEDS CONTROLLED

When applied as recommended under the conditions described, this product CONTROLS, PARTIALLY CONTROLS or SUPPRESSES most woody brush, trees and herbaceous weeds, some of which are listed below.

WOODY BRUSH AND TREES

Alder

Alnus spp.

Ash

Fraxinus spp.

Aspen, quaking

Populus tremuloides

Bearmat (Bearclover)

Chamaebatia foliolosa

Beech

Fagus grandifolia

Birch

Betula spp.

Blackberry

Rubus spp.

Blackgum
Nyssa spp.

Bracken
Pteridium spp.

Broom:
French
Cytisus monspessulanus
Scotch
Cytisus scoparius

Buckwheat, California
Eriogonum fasciculatum

Cascara
Rhamnus purshiana

Catsclaw
Acacia greggi

Ceanothus
Ceanothus spp.

Chamise
Adenostoma fasciculatum

Cherry:
Bitter
Prunus emarginata

Black
Prunus serotina

Pin
Prunus pensylvanica

Coyote brush
Baccharis consanguinea

Creeper, Virginia

Parthenocissus quinquefolia

Dewberry

Rubus trivialis

Dogwood

Cornus spp.

Elderberry

Sambucus spp.

Elm

Ulmus spp.

Eucalyptus, bluegum

Eucalyptus globulus

Hasardia

Haplopappus squamosus

Hawthorn

Crataegus spp.

Hazel

Corylus spp.

Hickory

Carya spp.

Holly, Florida; Brazilian Peppertree

Schinus terebinthifolius

Honeysuckle

Lonicera spp.

Hornbeam, American

Carpinus caroliniana

Kudzu

Pueraria lobata

Locust, black
Robina pseudoacacia

Madrone
Arbutus menziesii

Manzanita
Arctostaphylos spp.

Maple
Acer spp.

Monkey Flower
Mimulus guttatus

Oak
Quercus spp.

Persimmon
Diospyros spp.

Pine
Pinus spp.

Poison Ivy
Rhus radicans

Poison Oak
Rhus toxicodendron

Poplar, yellow
Liriodendron tulipifera

Prunus
Prunus spp.

Raspberry
Rubus spp.

Redbud, eastern
Cercis canadensis

Rose, multiflora
Rosa multiflora

Sage, black
Salvia mellifera

Sagebrush, California
Artemisia californica

Salmonberry
Rubus spectabilis

Saltbush; Sea myrtle
Baccharis halimifolia

Sassafras
Sassafras albidum

Sourwood
Oxydendrum arboreum

Sumac
Rhus vernix

Sweetgum
Liquidambar styraciflua

Swordfern
Polystichum munitum

Tallowtree, Chinese
Sapium sebiferum

Tan Oak
Lithocarpus densiflorus

Thimbleberry
Rubus parviflorus

Tobacco, tree

- Nicotiana glauca*
- Trumpet creeper**
Campsis radicans
- Waxmyrtle, southern**
Myrica cerifera
- Willow**
Salix spp.

HERBACEOUS WEEDS

- Bahiagrass**
Paspalum notatum
- Balsamapple**
Momordica charantia
- Barnyardgrass**
Echinochloa crus-galli
- Bassia, fivehook**
Bassia hysopifolia
- Bermudagrass**
Cynodon dactylon
- Bindweed, field**
Convolvulus arvensis
- Bluegrass, Kentucky**
Poa pratensis
- Brackenfern**
Pteridium aquilinum
- Brome**
Brome spp.

Bromegrass, smooth
Bromus inermis

Broomsedge
Andropogon spp.

Buttercup
Ranunculus spp.

Cheat
Bromus secalinus

Chickweed, mouseear
Cerastium vulgatum

Clover, red
Trifolium pratense

Clover, white
Trifolium repens

Cocklebur
Xanthium strumarium

Crabgrass
Digitaria spp.

Dallasgrass
Paspalum dilatatum

Dock, curly
Rumex crispus

Dwarf dandelion
Krigia cespitosa

Falseflax, smallseed
Camelina microcarpa

Fescue

30 of 37

Festuca spp.

Fiddleneck
Amsinckia spp.

Flaxleaf fleabane
Conyza bonariensis

Fleabane
Erigeron spp.

Foxtail
Setaria spp.

Groundsel, common
Senecio vulgaris

Guineagrass
Panicum maximum

Horsenettle
Solanum carolinense

Horseweed/Marestail
Conyza canadensis

Johnsongrass
Sorghum halepense

Kikuyugrass
Pennisetum clandestinum

Knapweed
Centaurea repens

Kochia
Kochia scoparia

Lambsquarters, common
Chenopodium album

Lespedeza: common, sericea

Lespedeza striata

Lespedeza cuneata

Lettuce, prickly

Lactuca serriola

Morningglory

Ipomoea spp.

Muhly, wirestem

Muhlenbergia frondosa

Mullein, common

Verbascum thapsus

Mustard, blue

Chorispora tenella

Mustard, tansy

Descurainia pinnata

Mustard, tumble

Sisymbrium altissimum

Mustard, wild

Sinapis arvensis

Napiersgrass

Pennisetum purpureum

Nightshade, silverleaf

Solanum elaeagnifolium

Nutsedge: purple, yellow

Cyperus rotundus

Cyperus esculentus

Oats, wild

Avena fatua

Orchardgrass

Dactylis glomerata

Panicum

Panicum spp.

Pampasgrass

Cortaderia jubata

Pennycress, field

Thlaspi arvense

Pigweed, redroot

Amaranthus retroflexus

Pigweed, smooth

Amaranthus hybridus

Quackgrass

Agropyron repens

Ragweed, common

Ambrosia artemisiifolia

Ragweed, giant

Ambrosia trifida

Reed, giant

Arundo donax

Ryegrass, perennial

Lolium perenne

Saltcedar

Tamarix spp.

Sandbur, field

Cenchrus spp.

Shepherdspurse

Capsella bursa-pastoris

Signalgrass, broadleaf
Brachiaria platyphylla

Smartweed, Pennsylvania
Polygonum pennsylvanicum

Sowthistle, annual
Sonchus oleraceus

Spanishneedles
Bidens bipinnata

Spurry, umbrella
Holosteum umbellatum

Starthistle, yellow
Centaurea solstitialis

Stinkgrass
Eragrostis cilianensis

Thistle, Canada
Cirsium arvense

Thistle, Russian
Salsola kali

Vaseygrass
Paaspalum urvillei

Velvetgrass
Holcus spp.

Witchgrass
Panicum capillare

INJECTION AND CUT STUMP APPLICATIONS

Woody brush and trees may be controlled using injection or cut stump applications of this product in forestry and utility right-of-way sites.

INJECTION APPLICATIONS

Apply the equivalent of 1 ml of this product per each 2 inches of trunk diameter. This is best achieved by applying a 25 to 100 percent concentration of this material either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying diluted material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In these species, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, avoid applications during peak sap flow in the Spring.

CUT STUMP TREATMENTS

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. Apply a 50 to 100 percent solution of this product to the freshly cut surface immediately after cutting. Delays in application may result in reduced performance. For best results, avoid applications during peak sap flow in the Spring.

When used according to directions for cut stump application, this product will CONTROL, PARTIALLY CONTROL or SUPPRESS most woody brush and tree species, some of which are listed below:

Alder

Alnus spp.

Coyotebrush

Baccharis consanguinea

Dogwood

BEST AVAILABLE COPY

Cornus spp.

Eucalyptus

Eucalyptus spp.

Hickory

Carya spp.

Madrone

Arbutus menziesii

Maple

Acer spp.

Oak

Quercus spp.

Poplar

Populus spp.

Saltcedar

Tamarisk spp.

Sweetgum

Liquidambar styraciflua

Sycamore

Platanus occidentalis

Tan Oak

Lithocarpus densiflorus

Willow

Salix spp.

Product protected by
U.S. Patent No. 4,405,531.
Other patents pending.

36 of 37

©MONSANTO COMPANY 19976

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

(ACCORD 96-1)

3433

(1/16/971/19/96)

CALIFORNIA

Accord® herbicide has been approved by the U.S. Environmental Protection Agency for the uses, crops and sites listed on this label and by California under label designation 19976-1. Approval of the items listed below is pending under the State of California registration requirements. With the exception of these items, this booklet contains the material approved by California in label 19976-1.

These use conditions, crops and sites may not be treated with this product in California until approval is received.

- Use of this product with Entry II surfactant, Arsenal Applicators Concentrate or Arsenal 2WSL.

- Use of this product for injection treatments on the following species:

- Alder
- Coyotebrush
- Eucalyptus
- Madrone
- Saltcedar
- Tan Oak
- Willow

- Use of this product for cut stump treatments on the following species:

- Coyotebrush
- Dogwood
- Hickory
- Maple
- Poplar
- Sycamore