

PM

25

524-326

Page 1 of 3

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ACCORD® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Accord herbicide and this supplement.

ACCORD  
HERBICIDE BY MONSANTO

EPA Reg. No. 524-326

ACCORD HERBICIDE FOR CONTROL OR SUPPRESSION OF BROMUS SPECIES AND MEDUSAHEAD RYE FOUND IN INDUSTRIAL, RANGELAND AND PASTURE SITES IN COLORADO, IDAHO, IOWA, KANSAS, MONTANA, NEBRASKA, NORTH DAKOTA, OREGON, SOUTH DAKOTA, UTAH, WASHINGTON AND WYOMING.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, call Collect, day or night, 314-694-4000.

Accord® is a registered trademark of Monsanto Company.

Directions for Use

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

When applied as directed in this label, there are no grazing restrictions.

APPLICATION EQUIPMENT AND TECHNIQUES

ACCEPTED

APR 21 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-326

BEST AVAILABLE COPY

Applications may be made using ground or aerial equipment. Aerial applications for these uses may be made using fixed wing or helicopter equipment. For aerial applications, apply in 2 to 10 gallons of water per acre. For applications using ground equipment, apply in 10 to 20 gallons of water per acre.

Mix 0.5 to 1.0 percent nonionic surfactant (2 to 4 quarts per 100 gallons of spray solution) with the spray solution. The surfactant should be at least 80 percent active ingredient.

---

BROMUS SPECIES

---

This product may be used to treat downy brome (*Bromus tectorum*), Japanese brome (*Bromus japonicus*), soft chess (*Bromus mollis*) and cheatgrass (*Bromus secalinus*) found in industrial, rangeland and pasture sites. Apply 8 to 16 fluid ounces of product per acre on a broadcast basis.

For best results, treatment should coincide with early seedhead emergence of the most mature plants. Delaying the application until this growth stage will maximize the emergence of other weedy grass flushes. Applications should be made to the same site each year until seed banks are depleted and the desirable perennial grasses are able to become reestablished on the site.

---

MEDUSAHEAD RYE

---

To treat medusahead rye, apply 16 fluid ounces of this product per acre as soon as plants are actively growing, and prior to seedhead emergence the 4 leaf stage. Applications may be made in the fall or spring.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Accord before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167

