

PROPOSED BRONCO® LABEL CHANGES

- ① "applied alone or in tank mixtures"
- ② Substitute for this section the following:

ATTENTION

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this herbicide can cause severe damage or destruction to the crop, plants, or other areas on which treatment was not intended. The likelihood of injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

- ③ Substitute for BroncoTM the phrase "this product"
- ④ "alone or"
- ⑤ Add to the list "Johnsongrass (*Sorghum halepensis*)"
- ⑥ Substitute for this section the following:

When applied as directed under conditions described, tank mixtures of Bronco with products listed in the following table will control or reduce competition from the additional grass and broadleaf weeds as indicated. Refer to the crop on which treatment is intended in the "Directions for Use" section of this label to determine the tank mixtures and rates recommended for use in each crop.

NOTE: C = Control R = Reduced Competition

	Bronco	Atrazine	Bladex	Lorox	Prinsep	Sencor/ Laxzone
Carpetweed <i>Mollugo verticillata</i>	C	C	C	C	C	C
Cocklebur <i>Xanthium pensylvanicum</i>	*	R	R	R	R	R
Jimsonweed <i>Datura stramonium</i>	*	C	R	R	R	R
Kochia <i>Kochia scoparia</i>	*	C	*	*	*	*
Lambsquarters <i>Chenopodium album</i>	R	C	C	C	C	C
Morningglory (annual) <i>Ipomoea purpurea</i>	*	R	R	R	R	R
Mustard <i>Brassica spp.</i>	*	C	C	C	C	C
Nightshade (black) <i>Solanum nigrum</i>	C	C	C	C	C	C
Nightshade (hairy) <i>Solanum sarachoides</i>	C	C	C	C	C	C
Pigweed, Carelessweed (redroot) <i>Amaranthus retroflexus</i>	C	C	C	C	C	C
Pigweed, (smooth) <i>Amaranthus hybridus</i>	C	C	C	C	C	C

100% 100%
100% 100%

NOTE: C = Control R = Competition

	Bronco	Atrazine	Bladex	Lorox	Princetown	Sencor/ Lexone
Purslane <i>Portulaca oleracea</i>	C	C	C	C	C	C
Pusley (Florida) <i>Richardia scabra</i>	C	C	C	C	C	C
Ragweed (common) <i>Ambrosia artemisiifolia</i>	R	C	C	C	C	C
Ragweed (giant) <i>Ambrosia trifida</i>	*	C	R	R	R	R
Sandbur (field) <i>Cenchrus spp.</i>	*	C	*	*	*	*
Sida (spiny), teaweed <i>Sida spinosa</i>	R	C	C	C	C	C
Smartweed <i>Polygonum pensylvanicum</i> <i>Polygonum persicaria</i>	R	C	C	C	C	C
Velvetleaf, Buttonweed <i>Abutilon theophrasti</i>	*	R	R	R	R	R

- (7) Substitute for 20 to 30, "10 to 30"
- (8) Insert the following sentence:
For best results with ground application equipment, use flat fan or whirl-chamber nozzle.
- (9) Omit the subheading "Tank Mixtures"
- (10) Substitute for "These tank mixtures" the following phrase:
"This product or labelled tank mixtures"
- (11) Insert the following section:

Tank Mixtures with Roundup and/or Lasso

For additional postemergent or preemergent weed control, Roundup or Lasso, respectively, may be added to the recommended treatment rates for this product. Consult the Roundup® and Lasso® labels for specific recommendations. The following table shows the amounts of Roundup and/or Lasso that can be added to specific treatment rates of this product.

<u>Recommended Rate (Quarts)</u>	<u>Additional Product (Quarts)</u>	
<u>BRONCO</u>	<u>ROUNDUP</u>	<u>LASSO</u>
3.25	1.9	1.9
3.5	1.8	1.7
4.0	1.6	1.4
4.5	1.4	1.1
5.0	1.25	0.75

DRAFT
DO NOT USE

