

READ THE ENTIRE LABEL FOR BRONCO® AND LASSO® HERBICIDES BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED ON THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the Label Pamphlet for Bronco, the Label Booklet for Lasso and this supplement.



For weed control in Corn and Soybeans

EPA Reg. No. 524-314-AA

ACCEPTED

JUN - 3 1987

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-341

FOR DISTRIBUTION AND USE ONLY IN KANSAS AND NEBRASKA.

For postemergence control of volunteer wheat and downy brome and preemergence control of annual weeds listed in the label for Bronco in ecofallow grain sorghum systems.

Keep out of reach of children.

**DANGER!**

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

• Bronco and Lasso® are registered trademarks of Monsanto Company.

MONSANTO COMPANY • ST. LOUIS, MISSOURI, 63167 U.S.A.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of herbicide application.

Observe the protective clothing requirements as listed in the Precautionary Statements of the labels for Bronco and Lasso.

Observe all mixing, spraying and handling requirements on the labels for Bronco and Lasso.

When Bronco herbicide is applied in a tank mixture with Lasso plus atrazine, this product can be applied up to 30 days prior to planting of grain sorghum to control emerged volunteer wheat and downy brome in reduced tillage and ecofallow systems. This tank mixture will also provide preemergence control or reduced competition of annual grasses and broadleaf weeds listed on the label for Bronco. Observe all precautions and limitations on the labels for Bronco, Lasso and atrazine before use of this tank mixture.

APPLY BRONCO PLUS LASSO PLUS ATRAZINE ONLY ON FIELDS WHERE THE GRAIN SORGHUM SEED HAS BEEN PROPERLY TREATED WITH A SEED PROTECTANT CONTAINING THE ACTIVE INGREDIENT, FLURAZOLE.

## METHOD OF APPLICATION

Broadcast Treatment — Apply this tank mixture in 10 to 30 gallons of water, or 10 to 50 gallons of liquid nitrogen (28 or 32 percent only) per acre.

## RECOMMENDED RATE AND TIMING OF APPLICATION

These recommendations are for ecofallow systems in which atrazine was applied the previous season (July to November) after wheat harvest. For fields not treated with atrazine the previous season, refer to the label booklet for Lasso (the section regarding Lasso plus atrazine for grain sorghum) for the recommended rates.

Apply this tank mixture when volunteer wheat and downy brome are actively growing, less than 24 inches tall and prior to the heading stage of development.

## Medium and Fine Textured Soils

Single Application — Apply 2.5 quarts of Bronco plus 1.5 quarts of Lasso plus 1 quart of atrazine 4L per acre (1.25 pounds 80W or 1.1 pounds 90 percent dry flowable formulation) in tank mixture up to 30 days prior to planting.

Split Application — Apply 2.5 quarts of Bronco plus 0.5 quarts of Lasso plus 1 quart of atrazine 4L per acre (1.25 pounds 80W or 1.1 pounds 90 percent dry flowable formulation) in tank mixture up to 30 days prior to planting. At planting, apply 1 quart of Lasso per acre.

## Coarse Textured Soils (except sands):

Single Application — Apply 2.5 quarts of Bronco plus 1.5 quarts of Lasso plus 0.5 quarts of atrazine 4L per acre (0.625 pound 80W or 0.55 pound 90 percent dry flowable formulation) in tank mixture up to 14 days prior to planting.

## ADDITIVES

Ammonium Sulfate — The addition of "fine or feed grade" ammonium sulfate to the spray solution may increase the performance of Bronco and tank mixtures containing Bronco applied in water on emerged annual weeds. When using this additive, add 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water. Refer to the label for Bronco for additional important information on the use of ammonium sulfate with Bronco.

Surfactants — Nonionic surfactants which are labeled for use with herbicides may be used with Bronco. DO NOT reduce rates of Bronco when adding surfactant. Use 0.5 percent surfactant concentration (2 quarts per hundred gallons of spray solution) when using surfactants which contain at least 50 percent active ingredient, or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 50 percent active ingredient. Read and observe cautionary statements and information on the surfactant label.

## MIXING INSTRUCTIONS

Fill the sprayer tank one-half full with water or a labeled fluid fertilizer before adding the tank mix components. If ammonium sulfate is used, add it slowly to the tank

with agitation in the following order: 1. fluid fertilizer, 2. surfactant, 3. water.

• Avoid driving this 1 plants, to speed or

• DO NOT (ers, herb listed on

• DO NOT sorghum

• DO NOT in 8 week Flush spray

Read the "Label pamphlets apply terms are at once.

©MONSANTO

MONSANTO AGRICULTURAL ST. LOUIS

BEST AVAILABLE COPY

121