


- 7
- U
- 7
- (A) any weed or brush species
- (B) THE USE OF A NONIONIC SURFACTANT APPROVED FOR THE SITE OF THE DESIRED APPLICATION IS REQUIRED FOR USE WITH THIS PRODUCT. Use 1/4 to 1/2 percent surfactant by volume. Carefully observe all cautionary statements and other information appearing on the surfactant label.
- (C) 3 to 20
- (D) weed or brush species
- (E) 3 to 20
- (F) Apply to actively growing annual grasses and broadleaf weeds. Use 1-1/2 pints of this product plus 1/4 to 1/2 % nonionic surfactant per acre if weeds are less than 6 inches tall. If weeds are over 6 inches tall, apply 2-1/2 pints of this product plus 1/4 to 1/2 % surfactant per acre. Allow at least 3 days after application before disturbing treated vegetation. After that period the weeds may be mowed, tilled or burned. See "Application Equipment and Techniques" for specific volumes of water.
- (G) When applied as directed under the conditions described in this label, this product plus nonionic surfactant will control the following annual weeds:
- (H) actively growing
- (I) Unless otherwise specified, allow at least 7 days after application before disturbing vegetation.
- (J) Add 1/4 to 1/2 % nonionic surfactant by volume to the rates of this product given in this list. See the "General Information" and "Directions for Use" sections of this label for additional information.
- (K) Perennial weeds. See attachment A.
- (L) Add 1/4 to 1/2 % nonionic surfactant by volume to the rates of this product given in this list. See the "General Information" and "Directions for Use" sections of this label for additional information.
- (M) Ensure thorough coverage when using hand-held equipment.
- (N) AQUATIC SITES

When applied as directed under conditions described, this product plus nonionic surfactant will control or partially control emerged annual and perennial weeds and woody brush and trees listed in this label. This product does not control plants which are either completely submerged or have a majority of the foliage under water. See the "Weeds Controlled" section of this label for rates and degree of control provided.

U.S. DEPARTMENT OF AGRICULTURE
 OFFICE OF PESTICIDE REGISTRATION
 WASHINGTON, D.C. 20250

7



This product may be used in and around aquatic sites, including all bodies of fresh and brackish water, which may be flowing, non-flowing, or transient. This includes lakes, rivers, streams, ponds, seeps, irrigation and drainage ditches, canals, reservoirs, and similar sites. There is no restriction on the use of water for irrigation, recreation, or domestic purposes.

For treatments after drawdown of water or in dry ditches, allow 7 or more days after treatment before reintroduction of water. Apply the product within one day after drawdown to ensure application to actively growing weeds.

When using this product in aquatic sites where water is present, add 1 to 2 quarts of Ortho X-77 surfactant per 100 gallons of spray solution (1/4 to 1/2 % surfactant by volume).

When using this product in sites where water is not present (dry ditches, ditchbanks, dry canals), use 1 to 2 quarts of nonionic surfactant per 100 gallons of spray solution (1/4 to 1/2 % surfactant by volume).

Consult local state fish and game agency and water control authorities before applying this product to public water. Permits may be required to treat such water.

NOTE: Do not apply this product within 0.5 miles upstream of potable water intakes.

Do not apply this product on rice levees when flood water is present.

Do not apply in tidewater areas.

Floating mats of vegetation may require retreatment. Avoid washoff of sprayed foliage by spray boat or recreational boat backwash or by rainfall within six hours of application. Do not retreat within 24 hours following the initial treatment.

Applications made to moving bodies of water must be made while traveling upstream to prevent concentration of the product in water. When making any bankside applications, do not overlap more than 1 foot into open water. The maximum application rate of 7-1/2 pints per acre must not be exceeded in any single application. Do not spray across open moving bodies of water.

When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in fish kill.

Ortho

ATTACHMENT A

The following perennial weeds will be added to the perennial weed list, in alphabetical order:

Alligatorweed*
Alternanthera philoxeroides

Maidencane
Panicum hematomon

Cattail
Typha Spp.

Paragrass
Brachiaria mutica

Giant Cutgrass*
Zizaniopsis miliacea

Phragmites**
Phragmites Spp.

Guineagrass
Panicum maximum

Spatterdock
Nuphar luteum

* Partial control.

** Partial control in Southeastern states. See description below.

Alligatorweed:

Broadcast 6 pints of this product per acre or apply a 1½ percent solution with hand-held equipment, to provide partial control of alligatorweed. Apply when most of the plants are in bloom. Repeat applications will be required to maintain such control.

Cattail:

Broadcast 4½ to 7½ pints of this product per acre or apply a ¾ percent solution with hand-held equipment, providing thorough coverage. Apply when most of the plants are in bloom. For best results, apply during the summer or fall months.

Giant Cutgrass:

Broadcast 6 pints of this product per acre or apply a 1 percent solution with hand-held equipment to provide control of giant cutgrass. Repeat applications will be required to maintain such control, especially of vegetation partially submerged in water. Allow for substantial regrowth to the seven to ten leaf stage prior to retreatment

0000

4

Maidencane/Paragrass:

Broadcast 6 pints of this product per acre or apply a 3/4 percent solution with hand-held equipment. Repeat treatments will be required, especially to vegetation partially submerged in water. Under these conditions, allow for regrowth to the seven to ten leaf stage prior to retreatment.

Phragmites (Southeastern States; SC, GA, AL, FL, MS, LA, TX):

Broadcast 7½ pints of this product per acre or apply a 1½ percent solution with hand-held equipment to provide partial control of Phragmites. Apply when most of the plants are in full bloom, or during the fall months. Repeat treatments will be required to maintain such control.

Phragmites (all other states):

Broadcast 6 pints of this product per acre or apply a 3/4 percent solution with hand-held equipment. Repeat treatments may be required to maintain control, due to the dense growth of these species preventing thorough spray coverage.

Spatterdock:

Broadcast 6 pints of this product per acre or apply a 3/4 percent solution with hand-held equipment. Apply when most plants are in full bloom. For best results, apply during the summer or fall months.