


193
124-343

READ THE ENTIRE LABEL FOR RODEO® BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED ON THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the Label Booklet for Rodeo and this supplement



EPA Reg No 524-343

WIPER APPLICATIONS FOR PARTIAL CONTROL OR SUPPRESSION OF SELECTED DITCHBANK AND AQUATIC WEEDS IN NONCROP AREAS.

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-343

Keep out of reach of children

CAUTION!



In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

• RODEO is a registered trademark of Monsanto Company.

MONSANTO COMPANY • AGRICULTURAL PRODUCTS • ST LOUIS, MISSOURI 63167 U.S.A.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in possession of the user at time of herbicide application.

GENERAL INFORMATION

Rodeo herbicide plus an approved surfactant, when applied by wiper application to actively growing plants, is effective for partial control or suppression of selected ditchbank and aquatic weeds.

AQUATIC SITES USAGE

See the "GENERAL INFORMATION," "MIXING AND APPLICATION INSTRUCTIONS" AND "AQUATIC SITES" sections of the label booklet for Rodeo for essential product performance information. See the "WEEDS CONTROLLED" section of the label booklet for Rodeo for optimum stage of treatment of weed for best results.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF THIS HERBICIDE WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS OR OTHER DESIRABLE VEGETATION SINCE DAMAGE OR DESTRUCTION MAY RESULT.

APPLICATION RECOMMENDATIONS WITH WIPER EQUIPMENT

This product may be applied through a wiper applicator, after dilution with water and thorough mixing, to weeds listed on this label growing in or along aquatic sites.

Wiper applicators, including wick devices, apply the herbicide solution by rubbing the weed with an absorbent material containing the herbicide solution.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desired vegetation should be adjusted so that the lowest wiper contact point is at least two (2) inches above this vegetation. Application made above desirable vegetation should be made when the weeds are a minimum of six (6) inches above this vegetation.

Best results may be attained when more of the weed is exposed to the herbicide solution. Weeds not contacted (wiped) with the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when the height of the weed varies so that not all weeds are contacted.

In severe infestations, reduce equipment ground speed to ensure that adequate amounts of this herbicide solution are wiped onto the weeds. When wiping moderate weed infestations an adequate flow rate should be 3 to 4 quarts of herbicide solution per mile of canal (wiping 4 foot band) for best results, do not allow wiper applicator to contact water.

NOTE:

- Maintain wiper equipment in good operating condition.

- Adjust height of wiper applicator to ensure adequate contact with weeds.
- Keep wiping surfaces clean.
- Keep wiper material at proper degree of saturation with herbicide solution.
- DO NOT use wiper equipment when weeds are wet, or under conditions where wave action or other water immersions will wash the solution off the weed.
- DO NOT operate equipment at ground speeds of greater than 5 MPH. As weed density increases, reduce equipment ground speed to ensure good coverage of weeds.
- Be aware that on sloping ground, the herbicide solution may migrate, causing dripping on the lower end and drying on the upper end of the wiper applicator.
- Variation in equipment design may affect weed control. With wiper applicators, the wiping material and its orientation must allow delivery of sufficient quantities of the recommended herbicide solution directly to the weeds.
- Mix only the amount of solution to be used during a one day period, as reduced activity may result from use of leftover solutions.

MIXING INSTRUCTIONS

Mix 2½ gallons of Rodeo herbicide with 7½ gallons of water to prepare a 25 percent solution. Add 1 quart of an approved surfactant per 10 gallons of herbicide solution (2½ percent surfactant by total volume). Apply this solution to weeds listed below.

WEEDS PARTIALLY CONTROLLED OR SUPPRESSED

When applied as recommended under the conditions described for "Wiper Equipment," this product will partially control or suppress the following weeds:

Bromegrass (smooth)	Mullein (common)
Bromus inermis	Verbascum thapsus
Canarygrass (reed)	Quackgrass
Phalaris arundinacea	Agropyron repens
Dock (curly)	Thistle (Canada)
Rumex crispus	Cirsium arvense

Read "LIMIT OF WARRANTY AND LIABILITY" in the Rodeo booklet before buying or using this product. If the terms are not acceptable, return the product unopened at once.

*Ortho II is a trademark of The Ortho Chemical Company.

21061T-1-8

3/30/95

© MONSANTO COMPANY 1995


MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST LOUIS, MISSOURI 63167 U.S.A.




243

READ THE ENTIRE LABEL FOR RODEO® AND FOR GARLON™ 4 BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the Label Booklet for Rodeo and this supplement.



EPA Reg. No. 524-343



Rodeo herbicide plus Garlon 4 tank mixture for burndown and partial control or suppression of woody brush and weeds in industrial sites.

Keep out of reach of children.

CAUTION!



In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4700

® RODEO is a registered trademark of Monsanto Company.

MONSANTO COMPANY • AGRICULTURAL PRODUCTS • ST. LOUIS, MISSOURI 63167 U.S.A.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

This label should be in the possession of the user at the time of herbicide application.

NONCROP USE

See the "GENERAL INFORMATION" and "MIXING AND APPLICATION INSTRUCTIONS" sections of this label for essential product performance information and the following NONCROP SECTIONS for specific recommended uses.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURFGRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT

Repeat treatments may be necessary to control weeds regenerating from underground parts or seeds or regrowth from areas not adequately contacted by the spray mixture.

Rodeo herbicide does not provide residual weed control. For subsequent weed control, follow a label-approved herbicide program.

Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

INDUSTRIAL SITES

When applied as directed for "Noncrop Uses" under the conditions described, this product, and an approved surfactant plus Garlon 4, provides burndown and partial control or suppression of woody brush and vegetation labeled for this product. This tank mixture is recommended for use on rights-of-way (utility, railroad, highway, pipeline), fence rows, roadsides, nonirrigation ditchbanks, wasteland and similar noncrop or industrial sites.

Hand-Held and High-Volume Applications.

Use 3 to 6 pints of Rodeo herbicide and 2 or more quarts of an approved surfactant, plus 1 to 2 quarts of Garlon 4 per 100 gallons of spray solution and apply to foliage of actively growing woody brush and weeds. Applications should be made on a spray to wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff.

Broadcast Applications with Ground Equipment:

Use 3 to 6 pints of Rodeo plus 1/4 to 2 quarts of Garlon 4 in sufficient water and make 20 to 100 gallons of total spray per acre. Use 2 to 4 quarts of an approved surfactant per 100 gallons of spray solution with this product.

Aerial Applications (Helicopter Only):

Use 3 to 6 pints of Rodeo plus surfactant plus 1 to 2 quarts of Garlon 4 and apply in a total spray volume of 10 to 20 gallons per acre. Aerial sprays should be applied using suitable drift control. Use 2 to 4 quarts of an approved surfactant per 100 gallons of spray solution with this product.

Apply when plants are actively growing and after full leaf expansion of woody brush. Use the higher rates of these products where vegetation is heavy or dense, or where hard-to-control brush species are prevalent. Repeat applications may be necessary to maintain control and to suppress areas where canopying of vegetation prevents good spray coverage and penetration.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Read and carefully observe the label claims, cautionary statements and all information on the labels of both products used in this tank mixture. Use according to the most restrictive label directions for each product in the mixture.

When used in combination as recommended by Monsanto Company, the liability of Monsanto shall in no manner extend to any damage, loss or injury not directly caused by the inclusion of the Monsanto product in such combination use.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Rodeo before buying or using this product. If the terms are not acceptable, return the product unopened at once.

*Garlon is a trademark of Dow Chemical Company.

21061T1-10

3/30/95

© MONSANTO COMPANY 1995

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.



NOV - 7 1995

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 524-343



(EPA Reg. No. 524-343)

RODEO HERBICIDE FOR FOLIAR APPLICATION TO SUPPRESS GROWTH AND SEEDHEAD DEVELOPMENT OF LISTED TURF SPECIES IN INDUSTRIAL SITES.

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

Apply 3 to 5 fluid ounces of this product per acre alone or in a recommended tank mixture. Spray volumes of 10 to 40 gallons per acre are recommended.

When using this product, mix 2 quarts of a non-ionic surfactant per 100 gallons of spray solution.

This product can be used for growth and seed-head suppression of:

Tall Fescue
Smooth Brome

For best results, apply this product in a recommended tank mixture to actively growing turf-grasses after greenup in the spring of the year. For suppression of seedheads, applications must be made before boot-to-seedhead stage of development. Applications made from seedhead emergence until maturity may result in turf discoloration or injury.

After mowing or removal of seedheads, this product in a recommended tank mixture may also be used to suppress the growth of certain turfgrasses. Allow turf to recover from stress caused by heat, drought or mowing before making applications. Applications made to turf under stress may increase the potential for discoloration or injury.

ANNUAL GRASSES

For growth suppression of some annual grasses such as annual ryegrass, wild barley and wild oats, apply 3 to 4 ounces of this product in 10 to 40 gallons of spray solution per acre. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

TANK MIXTURES OR INDUSTRIAL TURF GRASSES

For the following tank mixtures, consult each product label for weeds controlled and the proper stage of application. Do not treat turf under stress.

Tank Mixtures plus 2,4-D Amine

For additional weed control benefits, up to 1 quart per acre of 2,4-D amine may be added to the following tank mixtures.

TALL FESCUE Rodeo plus Telar™

For suppression of tall fescue growth and seed-heads, and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 1/2 ounce of Telar per acre.

This tank mixture can also be applied after mowing or removal of tall fescue seedheads for turf growth suppression and control of partial control of some annual weeds. Make only one of the above applications per growing season.

Rodeo plus Oust™

For suppression of tall fescue growth and seed-heads, and control or partial control of some

annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 1/4 ounce of Oust per acre.

Rodeo plus Escort™

This tank mixture can be applied after mowing or removal of tall fescue seedheads for turf growth suppression and control of partial control of some annual weeds. Use up to 1/3 ounce of Escort per acre.

SMOOTH BROME Rodeo plus Oust

For suppression of smooth brome growth and seedheads and control or partial control of some annual weeds, apply this tank mixture after greenup and prior to boot-to-seedhead stage of development. Use up to 1/4 ounce of Oust per acre.

Read and carefully observe the label claims, cautionary statements, and all information on the label for Rodeo herbicide.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Rodeo before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

Rodeo® is a registered trademark of Monsanto Company.

™Escort, Oust and Telar are trademarks of E. I. du Pont de Nemours and Company.

21061T1-9

3/30/95

© MONSANTO COMPANY 1995

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

