

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR ROUNDUP CUSTOM™ HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Custom herbicide and this supplement.

Roundup Custom Logo

EPA Reg. No. 524-343

Roundup Custom is a trademark of Monsanto Company.

WIPER, PREEMERGENCE, SPOT TREATMENT AND CREATING ROW APPLICATIONS IN GRASS SEED PRODUCTION

Keep out of reach of children.

CAUTION!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Custom herbicide for essential product performance information.

Do not feed or graze treated areas for 8 weeks following application.


RECOMMENDATIONS

1.0 Wiper Applications

When applied as directed in the "SELECTIVE EQUIPMENT" section of the Round Custom herbicide label, this product may be used for weed control in grasses grown for seed production.

Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators should be adjusted so that the wiper contact point is at least two (2) inches above the desirable vegetation.

Weeds should be a minimum of six (6) inches above the desirable vegetation. Better results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected.



This may occur in dense clumps, severe infestations, or when height of weeds varies so that not all weeds are contacted. In these instances, repeat treatments may be necessary. Better results may be obtained if 2 applications are made in opposite directions.

2.0 Preemergence applications

This product may be applied postplant and preemergence to grasses grown for seed production. Applications **MUST** be made prior to the emergence of the crop to avoid crop injury.

See the "WEEDS CONTROLLED" section of the Roundup Custom herbicide label for specific rate recommendations and the "APPLICATION EQUIPMENT AND TECHNIQUES" section for application recommendations.

3.0 Spot Treatments

When applied as directed under the conditions described, Roundup Custom herbicide will control grass and broadleaf weeds listed in the Roundup Custom herbicide label booklet.

Use a 1 to 5 percent solution of Roundup Custom herbicide on a volume to volume basis with water. See the "SELECTIVE EQUIPMENT" section for additional application recommendations.

Apply this product prior to heading of grasses. Do not treat more than 10 percent of the total field to be harvested. The crop receiving the spray in the treated area will be killed. Take care to avoid drift or spray outside the target area for the same reason.

4.0 Creating Rows in Annual Ryegrass

Roundup Custom herbicide may be used to create rows in annual ryegrass stands.

Use 12 to 24 fluid ounces of this product per acre mixed with water. Use the higher rate when the ryegrass is greater than 6 inches in height.

Set nozzle heights to allow the establishment of the desired row spacing while preventing spray droplets, spray fines, or drift to contact the ryegrass plants not treated. Use of low pressure nozzles, or drop nozzles designed to target the application over a narrow band are recommended.

Grower assumes all responsibility for crop losses from misapplication.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Custom herbicide before using this product. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2000 MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR ROUNDUP® CUSTOM HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Roundup Custom herbicide and this supplement.

EPA Reg. No. 524-343

ROUNDUP CUSTOM
Herbicide by Monsanto

ACCEPTED
FEB 16 2000
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-343

ROUNDUP CUSTOM HERBICIDE FOR POSTEMERGENCE WEED CONTROL IN SILVICULTURAL SITES AND UTILITY RIGHTS OF WAY.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Roundup Custom herbicide for essential product performance information.

This product requires use of a nonionic surfactant. When using this product, mix 2-4 quarts of a nonionic surfactant per 100 gallons of spray solution. Examples of when to use the higher surfactant rate include, but are not limited to: high water volumes, adverse environmental conditions, tough to control weeds, weeds under stress, surfactants with less than 70% active ingredient, tank mixes, etc.

NOTE: Applications of this product should not contact leaves of desirable plants since foliar injury, discoloration or death may result.

FORESTRY SITE PREPARATION AND UTILITY RIGHTS-OF-WAY

This product is recommended for the control or partial control of woody brush, trees and herbaceous weeds. This product is labeled for use in forestry and utility sites. This product is also recommended for use in preparing or establishing wildlife openings within these sites and maintaining logging roads, and for side trimming along utility rights-of-way.

In forestry, this product is recommended for use in site preparation prior to planting any tree species, including Christmas trees, and silvicultural nursery sites.

In utilities, this product is recommended for use along electrical power, pipeline and telephone rights-of-way, and in other utility sites associated with these rights-of-way, such as substations.

APPLICATION RATES AND TIMING

APPLICATION	ROUNDUP CUSTOM®	SPRAY VOLUME GAL/A
<u>BROADCAST</u>		
Aerial	1.5 to 7.5 qts/a	5 to 30
Ground	1.5 to 7.5 qts/a	10 to 60
<u>SPRAY-TO-WET</u>		
Handgun, Backpack, Mistblower	0.6 % to 2 % by volume	spray-to-wet
<u>LOW VOLUME DIRECTED SPRAY</u>		
Handgun, Backpack, Mistblower	4% to 7.5% by volume	partial coverage*

*For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results.

In forestry site preparation and utility rights-of-way applications, this product requires use with a nonionic surfactant. Use a nonionic surfactant with greater than 80 percent active ingredient and labeled for use with herbicides. Use of this product without surfactant will result in reduced performance. See the "MIXING AND APPLICATION INSTRUCTION" section of this label for more information.

Mix 2 or more quarts of the nonionic surfactant per 100 gallons of spray solution (0.5 percent or more by spray volume). Use of surfactant concentrations greater than 1.5 percent by spray volume with handgun applications or 2.5 percent by spray volume with broadcast applications is not recommended.

Use higher rates of this product within the recommended range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the recommended range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the recommended range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

This product has no herbicidal or residual activity in the soil. Where repeat applications are necessary, do not exceed 8 quarts of this product per acre per year.

TANK MIXTURES

Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of both products used. Use according to the most restrictive precautionary statements for each product in the mixture. Any recommended rate of this product may be used in a tank mix.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions. For side trimming treatments in utility rights-of-way, tank mixtures with Arsenal™ 2WG are not recommended. For side trimming treatments, it is recommended that this product be used alone as recommended, or as a tank mixture with Garon™ 4.

PRODUCT	BROADCAST RATE	USE SITES
Arsenal Applicators Concentrate**	2 to 16 fl oz /a	Forestry site preparation
Chopper™	4 to 32 oz/a	Forestry site preparation
Escort™**	½ to 3 ½ oz/a	Forestry site preparation
Oust™	1 to 4 oz/a	Forestry site preparation, Utility sites
Garlon 3A*, Garlon 4	1 to 4 qts/a	Forestry site preparation, Utility sites
Arsenal 2WSL**	4 to 32 fl oz/a	Utility sites

PRODUCT	SPRAY-TO-WET RATES	USE SITES
Arsenal Applicators Concentrate	1/32 % to 1/2 % by volume	Forestry site preparation
Arsenal 2WSL	1/16 % to 1/2 % by volume	Utility sites

PRODUCT	LOW VOLUME DIRECTED SPRAY RATES	USE SITES
Arsenal Applicators Concentrate	1/8% to 1/2% by volume	Forestry site preparation
Arsenal 2WSL	1/8% to 1/2% by volume	Utility sites

* Ensure that Garlon 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray compatibility problems.

**Arsenal and Escort are not registered in the state of California.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

FORESTRY CONIFER AND HARDWOOD RELEASE

DIRECTED SPRAY AND SELECTIVE EQUIPMENT

This product may be applied as a directed spray or by using selective equipment in forestry conifer and hardwood sites, including Christmas tree plantations and silvicultural nurseries. Mix 2 to 6 quarts of a nonionic surfactant per 100 gallons of spray solution (0.5 to 1.5 percent by spray volume) for all spray applications. Use a surfactant with greater than 80 percent active ingredient.

In hardwood plantations, tank mixtures with Oust may be used. In pine plantations, tank mixtures with Garlon 4 or Arsenal AC may be used. Comply with all site restrictions, forestry species limitations and precautions on the tank mix product label.

Avoid contact of spray, drift, mist or drips with foliage, green bark or non-woody surface roots of desirable species.

See all sections in the "APPLICATION EQUIPMENT AND TECHNIQUES" portion of this label for specific equipment recommendations and precautions.

For spray-to-wet applications, use a 1.5 percent spray solution for the control of undesirable woody brush and trees. To control herbaceous weeds, use a 0.75 to 1.5 percent solution.

For low volume directed spray applications, use a 4 to 7.5 percent spray solution. Coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the unwanted vegetation is important.

For equipment calibrated for broadcast applications, use 1.5 to 7.5 quarts of this product per acre. Apply in 10 to 60 gallons of clean water per acre. Shielded application equipment may be used to avoid contact of the spray solution with desirable plants. Shields should be adjusted to prevent spray contact with the foliage or green bark of desirable vegetation.

Wiper application equipment may be used. See the SELECTIVE EQUIPMENT portion of this label for equipment and rate recommendations.

BROADCAST SPRAY

Except where specifically recommended below, use only where conifers have been established for more than one year.

APPLICATION MUST BE MADE AFTER FORMATION OF FINAL CONIFER RESTING BUDS IN THE FALL OR PRIOR TO INITIAL BUD SWELLING IN THE SPRING.

Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied. Damage can be accentuated if applications are made when conifers are actively growing, or are under stress from drought, flood water, improper planting, insects, animal damage or diseases.

This product may require use with a surfactant. Unless otherwise recommended in this section of this label, use Entry™ II surfactant at 10 to 30 fluid ounces per acre. Follow the instructions under the "MIXING" portion of the "MIXING AND APPLICATION INSTRUCTIONS" section of this label.

For release of the following conifer species outside the Southeastern United States:

Douglas fir
Pseudotsuga menziesii

Fir
Abies spp.

Hemlock**
Tsuga spp.

Pines*
Pinus spp.

Redwood, California**
Sequoia spp.

Spruce
Picea spp.

RoundupCustom
Forestry/UROW

* Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.

**Use of a surfactant is not recommended for release of hemlock species or California redwood. In mixed conifer stands, injury to these species may result if a surfactant is used.

Apply 0.75 to 1.5 quarts of this product per acre as a broadcast spray.

NOTE: For release of Douglas fir with this product or recommended tank mixtures of this product, Entry II or a nonionic surfactant recommended for over-the-top foliar sprays may be used. To avoid possible conifer injury, Entry II rates should not exceed 20 fluid ounces per acre at elevations above 1500 feet, or 10 fluid ounces per acre in the coastal range or at elevations below 1500 feet in Washington and Oregon. Nonionic surfactants may be used at 2 fluid ounces per acre at elevations above 1500 feet, or 1 fluid ounce per acre in the coastal range or at elevations below 1500 feet. Use of surfactant rates exceeding those listed above may result in unacceptable conifer injury and are not recommended. Ensure that the nonionic surfactant has been adequately tested for Douglas fir safety before use.

In Maine, up to 2.25 quarts per acre of this product or a tank mix with 1 oz/a of Arsenal Applicators Concentrate may be used for the control of difficult species.

To release Douglas fir, pine and spruce species at the end of the first growing season(except in California), apply 0.75 to 1.125 quarts of this product per acre. Ensure that the conifers are well hardened off.

OUST TANK MIXTURES - To release jack pine, white pine and white spruce, apply 0.75 to 1.5 quarts of this product with 1 to 3 ounces(1 to 1.5 for white pine) of Oust per acre. Make applications to actively growing weeds as a broadcast spray over the top of established conifers. Applications at these rates should be made after formation of conifer resting buds in the late summer or fall.

ARSENAL APPLICATORS CONCENTRATE TANK MIXTURES - This product may be tank mixed with Arsenal Applicators Concentrate for release of Douglas fir. Use 0.75 to 1.125 quarts of this product tank mixed with 2 to 6 fluid ounces of Arsenal per acre. For release of balsam fir and red spruce, apply a mixture of 1.5 quarts of this product and 1 to 2 1/2 fluid ounces of Arsenal Applicators Concentrate per acre.

For release of the following conifer species in the Southeastern United States:

Loblolly pine

Pinus taeda

Eastern white pine

Pinus strobus

Shortleaf pine

Pinus echinata

Slash pine


Pinus elliottii

Virginia pine

Pinus virginiana

Longleaf pine

Pinus palustris



Apply 1.125 to 1.875 quarts of this product per acre as a broadcast spray during late summer or early fall after the conifers have hardened off. For applications at the end of the first growing season, use 0.75 quart per acre of this product alone or in a recommended tank mixture.

ARSENAL APPLICATORS CONCENTRATE TANK MIXTURES - Apply 0.75 to 1.5 quarts of this product with 2 to 16 fluid ounces of Arsenal Applicators Concentrate per acre as a broadcast spray for conifer release. Use only on conifer species that are labeled for over-the-top sprays for both products. Use the higher recommended rates for dense, tough-to-control woody brush and trees.

Read and carefully observe the label claims, cautionary statements and all information on the labels of each product used in these tank mixtures. Use according to the most restrictive precautionary statements for each product in the mixture.

HERBACEOUS RELEASE

When applied as directed, this product plus listed residual herbicides provides postemergence control of the annual weeds and control or suppression of the perennial weeds listed in this label, and residual control of the weeds listed in the residual herbicide label. Make applications to actively growing weeds as a broadcast spray over the top of labeled conifers.

Oust tank mixtures - To release loblolly pines, apply 12 to 18 fluid ounces of this product, plus 2 to 4 ounces of Oust per acre. To release slash pines, apply 9 to 12 fluid ounces of this product, plus 2 to 4 ounces of Oust per acre.

Mix up to 3.2 fluid ounces per acre of Entry II with the recommended rate of this product plus Oust. Applications can be made over newly planted pines after the emergence of herbaceous weeds in the spring or early summer. Best results are obtained from applications made in May and June.

Weed control may be reduced if water volumes exceed 25 gallons per acre for these treatments.

Atrazine tank mixtures - To release Douglas fir, apply 0.75 quart of this product, plus 4 pounds a.i. of atrazine per acre. Apply only over Douglas fir that has been established for at least one full growing season. Apply in the early Spring, usually mid-March through early April. Injury will occur if applications are made after bud swell in the Spring. Do not add surfactant to this mix for this use.

Always read and follow the manufacturer's label recommendations for all herbicides and surfactants used.

WETLAND SITES

This product may be used in and around water(aquatic areas) and wetlands found in forestry and in power, telephone and pipeline rights-of-way sites, including where these sites are adjacent to or surrounding domestic water supply reservoirs, supply streams, lakes and ponds. Read and observe the following before making applications in and around water.

Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat in such areas.

There is no restriction on the use of treated water for irrigation, recreation or domestic purposes.

NOTE: Do not apply this product directly to water within 1/2 mile up-stream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.

Do not spray open bodies of water where woody brush, trees and herbaceous weeds do not exist. The maximum application rate of 3.75 quarts per acre must not be exceeded in a single over-water broadcast application except as follows, where any recommended rate may be applied:

- Stream crossings in utility rights-of-way
- Where applications will result in less than 20 percent of the total water area being treated.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Roundup Custom herbicide before using this product. Those terms apply to this supplemental labeling and, if those terms are not acceptable, return the product unopened at once.

Roundup is a registered trademark of Monsanto Company
 Entry is a trademark of Monsanto Company.
 Oust and Escort are trademarks of E.I. DuPont de Nemours and Company.
 Garlon is a trademark of Dow Agrosciences .
 Arsenal and Chopper are trademarks of American Cyanamid Company.

©2000 MONSANTO COMPANY
 ST. LOUIS, MISSOURI 63167

