

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Aquamaster® herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.



EPA Reg. No. 524-343


AquaMaster is a trademark of Monsanto Company, LLC.

For Control of Cordgrass (*Spartina* spp.)

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night 314-694-4000.



Directions for Use

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label must be in the possession of the user at the time of the herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for Aquamaster herbicide for essential product performance information.

BROADCAST APPLICATIONS

AIR: Apply 4 to 7.5 pints of this product in 5 to 20 gallons of spray solution per acre. Add 1 to 2 quarts of nonionic surfactant per 100 gallons of spray solution.

GROUND: Apply 4 to 7.5 pints of this product in 10 to 60 gallons of spray solution per acre. For best results, ensure that complete coverage of cordgrass clumps is achieved. Add 1 to 2 quarts of a nonionic surfactant per 100 gallons of spray solution.

HAND-HELD EQUIPMENT

Apply a 2 to 8 percent solution of this product. Ensure that complete coverage of cordgrass clumps is achieved. Do not spray to the point of run-off. Add 1 to 2 quarts of a nonionic surfactant per 100 gallons of spray solution.

WIPER APPLICATIONS

For wick or wiper applications, mix 1 gallon of this product with 2 gallons of clean water to make a 33 percent solution. Addition of a nonionic surfactant at a rate of 10 percent by volume of the total herbicide solution is recommended.

In heavy stands, a double application in opposite directions may improve results

APPLICATION CONDITIONS

NOTE: Schedule application in order to allow 6 hours before treated plants are covered by tidewater. Rainfall or immersion within 6 hours after application may reduce effectiveness.

The presence of debris or silt on the cordgrass plants will reduce performance of this product. It may be necessary to wash targeted plants prior to application to improve uptake of this product into the plant. Where cordgrass has been cut or mowed prior to application with AquaMaster, ensure adequate regrowth of cordgrass occurs to allow for interception or absorption of the herbicide solution.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for AquaMaster before using this product. Those terms apply to this supplemental label and, if those terms are not acceptable, return the product unopened at once.

© [DATE] MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

