5au - 3uu

ł


IODIDAS SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR MICRO-TECH® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using Micro-Tech herbicide as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.



Herbicide by Monsanto



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "MIXING INSTRUCTIONS" and "APPLICATION EQUIPMENT AND INSTRUCTIONS" sections of the label booklet for Micro-Tech herbicide for essential product performance information.

RECOMMENDATION

This product may be used preemergence and postemergence to Roundup Ready Corn 2 until the corn reaches 5 inches in height. This product may be tank mixed with a Roundup® WeatherMAX or other Roundup agricultural herbicide – refer to the Roundup agricultural herbicide label for specific weeds controlled postemergence.

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING ROUNDUP WEATHERMAX OR OTHER ROUNDUP AGRICULTURAL HERBICIDE TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

Approved Application Systems

Ground--Broadcast boom

Approved Application Methods

Preemergence Surface - Sequential Program - This product may be applied at the Roundup Ready Rate of 2 quarts per acre preemergence to Roundup Ready Corn 2 in a planned sequential program followed by a postemergence application of Roundup WeatherMAX or other Roundup agricultural herbicide.

Use the full labeled rate of this product when the application is 14 days or more prior to planting or when tough grasses are present, i.e., *barnyardgrass, shattercane, woolly cupgrass, wild proso millet, seedling johnsongrass, sandbur.*

Labeled use rates are listed in the table below.

Postemergence Tank Mix

This product may be applied postemergence to Roundup Ready Corn 2 in a tank-mix with Roundup WeatherMAX or other Roundup agricultural herbicide from emergence until the crop is 5 inches in height. The Roundup Ready Rate for this product is 2.0 quarts per acre. Labeled use rates for the tank mix are defined in the table below. Use the higher rate on larger weeds and where heavy weed infestations exist. This tank mix should be applied when weeds are 2 to 4 inches in height and before the weed height and/or density become competitive with the crop. For difficult to control weeds such as *fall panicum*, *barnyardgrass, crabgrass, shattercane, broadleaf signalgrass and Pennsylvania smartweed*, use the higher rate of Roundup WeatherMAX or other Roundup agricultural herbicide.

ROUNDUP READY RATE – Micro-Tech herbicide at 2.0 quarts per acre.

Application Rates (minimum and maximum range)

5

ť

	Broadcast Rate Per Acre		
	Micro-Tech Herbicide (quarts)		Roundup WeatherMAX (ounces)
Soil Texture	Less than 3% organic matter	3% or more organic matter	
Coarse	2.0 to 2.25	2.0 to 2.75	16 to 22
Medium	2.0 to 2.75	2.0 to 2.75	16 to 22
Fine	2.0 to 2.75	2.0 to 3.25	16 to 22

Tank mixtures may cause a temporary burn, discoloration and/or growth reduction under stress conditions.

Micro-Tech, Roundup and Roundup Ready are trademarks of Monsanto Technology LLC

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label booklet for Micro-Tech herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

©2005 MONSANTO COMPANY ST. LOUIS, MISSOURI 63167

[date]

÷

[print code]