



HERBICIDE by Monsanto

For control or suppression of emerged grass and broadleaf weeds, woody brush and vines in bermudagrass roadside and highway rights-of-way in Alabama, Arkansas, Florida, Georgia, Louisiana, New Mexico, Mississippi, Oklahoma, South Carolina, and Texas.

Complete Directions for Use


AVOID CONTACT WITH FOLIAGE OF CROP OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT

EPA Reg. No. 524-351

Campaign® is a registered trademark of Monsanto Company

Change to:

For control or suppression of emerged grasses, broadleaf weeds and vines in bermudagrass, bahiagrass, and tall fescue turf on roadsides, highway rights-of-way, ditchbanks and dry ditches.



1988-1


897.16-000.01.53

1989-1

897.16-000.03/53

MONSANTO COMPANY 1988

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.



Read the entire label before using product.
Use only according to label instructions.

Read **LIMIT OF WARRANTY AND LIABILITY** before buying.
If terms are not acceptable, return at once.
No refund.

REFORMULATION OR REPACKAGING PROHIBITED

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for the label booklet (Directions) when used in accordance with those Directions under the conditions described therein. **NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE.** This warranty is also subject to conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with undesirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OF, HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this Limit Of Warranty and Liability which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

**Hazards to
Humans and Domestic Animals**

Keep out of reach of children.

DANGER!

CAUSES EYE BURNS.

HARMFUL IF SWALLOWED.

MAY CAUSE ALLERGIC SKIN REACTION

Do not get in eyes, on skin, or on clothing.

Wear goggles or face shield and rubber gloves in transferring and mixing and when adjusting, repairing or cleaning equipment.

Wash thoroughly with soap and water after handling.

FIRST AID - IN EYES: Immediately flush with plenty of water for at least 15 minutes. Call a physician.

ON SKIN: Immediately flush with plenty of water for at least 15 minutes while removing contaminated clothing. As soon as soap is available, wash thoroughly with soap and water. Wash clothing before reuse. Sensitized persons should avoid further contact and reuse of contaminated clothing.

IF SWALLOWED: Induce vomiting immediately by either giving two glasses of water and sticking finger down throat or, when available, administering syrup of ipecac, as prescribed on its container. Call a physician. **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water or wetlands (swamps, bogs, marshes or potholes). Do not contaminate water by cleaning equipment or disposal of wastes.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENTS*

Glyphosate (N-phosphonomethyl glycine) as the isopropylamine salt 12.9%

2,4-D(2,4-dichlorophenoxyacetic acid) as the isopropylamine salt 20.6%

INERT INGREDIENTS: 66.5%

100.0%

*Contains 144 grams per litre or 1.2 lbs per U.S. gallon of the active ingredient, glyphosate, as its isopropylamine salt and 227 grams per litre or 1.9 lbs per U.S. gallon of the active ingredient, 2,4-D, as its isopropylamine salt. Equivalent to 108 grams per litre or 0.9 lb per U.S. gallon of the acid, glyphosate and 182 grams per litre or 1.5 lbs per U.S. gallon of the acid, 2,4-D.

Product is protected by U.S. Patent No. 3,799,758, and U.S. Patent No. 4,405,531. Other patents are pending.

GENERAL INFORMATION

Campaign® is a postemergence herbicide for control or suppression of emerged weeds. This product is recommended for use on highway and roadside rights-of-way.

This product enters the plant through the foliage and moves throughout the plant. Visual effects of control are a gradual wilting or yellowing of the plant, which advances to complete browning of above-ground growth and deterioration of affected underground plant parts.

Change to:

Do not contaminate water when disposing of equipment washwaters.

Add:

No license granted under any non-U.S. patent.

Visible symptoms will usually develop on labeled weeds within 2 to 4 days after application, but may not occur for 7 or more days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay the visual effects of control.

APPLICATION PRECAUTIONS AND WARNINGS

- DO NOT APPLY IN THE VICINITY OF 2,4-D SENSITIVE CROPS SUCH AS COTTON, GRAPES, TOMATOES AND OTHER DESIRABLE VEGETATION.
- Applications should be made only when there is no hazard from spray drift since very small quantities of spray, which may not be visible, may severely injure susceptible crops or desirable vegetation.
- The likelihood of injury occurring to adjacent crops from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions including lesser wind velocities will favor spray drift.
- Movement of this product on soil particles during windstorms may cause damage to susceptible plants that are contacted. This hazard is reduced if rainfall occurs shortly after application.
- Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this herbicide or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

← Delete

Campaign is subject to all state and county regulations for 2,4-D amine.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. Do not store near fertilizers, seeds, insecticides or fungicides.

STORAGE

STORE ABOVE 40°F to keep product in solution. If crystals form, place in a warm room (72°F), allow the product to reach room temperature and roll or shake periodically until crystals have dissolved. Keep container closed to prevent spills and contamination.

Change to:

← ... allow the product to reach room temperature and roll or shake the container periodically...

DISPOSAL

Wastes of this pesticide may cause eye burns and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

Do not reuse container. Triple rinse container, then puncture and dispose in a sanitary landfill, or by incineration, if it is allowed by state and local authorities. Do not burn. Do not burn. Do not burn.

TIMING OF APPLICATION

This product should be applied postemergence to vigorously growing weeds at a rate recommended in the **RECOMMENDED RATES AND WEEDS CONTROLLED** section of this label. Application should be delayed until maximum emergence of the target weeds but before mature seeds are produced.

Reduced control may result if treatments are made during poor growing conditions such as drought stress, disease or insect damage or if weeds have been mowed, grazed or cut. Heavy dust on foliage or an overstory canopy covering targeted weeds may also reduce control.

Rainfall occurring within 6 hours after application may reduce effectiveness.

RELEASE OF DORMANT BERMUDAGRASS

When applied as directed, this product will provide control or suppression of many winter annual weeds and tall rescue for effective release of dormant bermudagrass. To avoid delays in greenup and minimize injury, do not treat when bermudagrass is in a semi-dormant condition. For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall rescue, treat when rescue is in or beyond the 4 to 6-leaf stage. See the **RECOMMENDED RATES AND WEEDS CONTROLLED** section for proper rate selection.

RELEASE OF ACTIVELY GROWING BERMUDAGRASS

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual and biennial species listed in the **WEEDS CONTROLLED** section of this label, and control or suppression of certain perennial weeds.

Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as size of plants increases or as they approach flower or seedhead formation. Use the higher rate of this product for control or partial control of perennial species. Use the lower rates for suppression of growth. For best results, see the **RECOMMENDED RATES AND WEEDS CONTROLLED** section for the proper rate selection.

Use only in areas where bermudagrass is a desirable ground cover and where some temporary injury or discoloration can be tolerated.

RECOMMENDED RATES AND WEEDS CONTROLLED

Apply 62 to 64 fluid ounces of this product in 15 to 30 gallons of spray solution per acre. Rate recommendations for specific weed species are given below.

For the best rate recommendation for the mixture of weeds within your geographic area, contact your Monsanto sales representative.

Replace with the following pages:

WEEDS CONTROLLED OR SUPPRESSED			
Note: S = Control • = Suppression			
ANNUAL WEED SPECIES	CAMPAIGN# 02-ACRE		
	32	48	64
Barley, little <i>Hordeum pusillum</i>	S	•	•
Bedstraw, catchweed <i>Galium aparine</i>	•	•	•
Bluegrass, annual <i>Poa annua</i>	S	•	•
Chenel <i>Chaerophyllum tainturieri</i>	•	•	•
Chickweed, common <i>Stellaria media</i>	•	•	•
Clover, crimson <i>Trifolium incarnatum</i>	•	•	•
Clover, lesp <i>Trifolium spp</i>	S	•	•
Crabgrass <i>Digitaria spp</i>	•	•	•
Foxtail <i>Setaria spp.</i>	S	•	•
Geranium, Caroline <i>Geranium carolinianum</i>	•	•	•
Henbit <i>Lamium amplexicaule</i>	•	•	•
Partridgepea <i>Cassia fasciculata</i>	S	•	•
Ragweed, common <i>Ambrosia artemisiifolia</i>	S	S	•
Speedwell, corn <i>Veronica arvensis</i>	S	•	•
Sponge, spotted <i>Euphorbia maculata</i>	S	•	•
Verbena, blue <i>verbena hastata</i>	S	•	•
BIENNIAL WEED SPECIES			
Carrot, wild <i>Daucus carota</i>	•	S	•
PERENNIAL WEED SPECIES			
Fescue, tall <i>Festuca arundinacea</i>	S	S	S
Greenbrier <i>Smilax spp</i>	•	•	S
Honeysuckle <i>Lonicera spp.</i>	•	S	•
Horsebottle <i>Solanum carolinense</i>	•	S	S
Johnsongrass <i>Sorghum halepense</i>	S	S	•
Plantain, broadleaf <i>Plantago lanceolata</i>	•	S	S
Raspberry <i>Rubus spp.</i>	•	•	S

Replace with the following pages:

PERENNIAL WEED SPECIES			
Ryegrass, Italian <i>Lolium multiflorum</i>	•	•	S
Trumpetcrapper <i>Campsis radicans</i>	•	•	S
Waxygrass <i>Paspalum firmile</i>	S	C	C
Velvet, Common <i>Vicia sativa</i>	•	•	S

Replace with the following pages:

8618
8

TANK MIXTURES

CAMPAIGN ALONE
CAMPAIGN + OUST TM
CAMPAIGN + ATRAZINE

BERMUDAGRASS

Read and carefully observe the label claims, cautionary statements, and all information on the labels of both products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Spring applications:

When applied as directed, this product will provide control or suppression of vines and many winter annual weeds for release of bermudagrass. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated.

Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.25 to 1 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

Summer applications:

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines for release of actively growing bermudagrass.

(Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.25 to 2 ounces of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

For spring and summer applications to bermudagrass, use only in areas where bermudagrass is a desirable ground cover and where some temporary discoloration can be tolerated.

TM Oust is a TRADEMARK OF E.I. duPont de Nemours and Company.

BAHIAGRASS

Dormant applications:

When applied as directed, this product will provide control or suppression of vines and winter annual weeds for release of dormant bahiagrass. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated. To avoid delays in greenup and minimize injury, treat when bahiagrass is dormant.

Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture apply 2 to 4 pints of this product plus 0.25 to 0.5 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

Active applications:

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines for release of bahiagrass.

Apply 1.5 to 2 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 1.5 to 2 pints of this product plus 0.25 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

TALL FESCUE

Spring applications:

When applied as directed, this product will provide control or suppression of vines and many winter annual weeds in tall fescue. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated. To minimize tall fescue injury, apply this product when tall fescue is 4 to 6 inches tall but prior to seedhead emergence.

Apply 2 to 3 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control of certain weeds. For this tank mixture, apply 2 pints of this product plus 0.25 ounces of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

Summer applications:

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines in tall fescue.

Apply 2 to 3 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control of certain weeds. For this tank mixture, apply 2 to 3 pints of this product plus 0.25 to 0.5 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct selection.

(CAMPAIGN® plus ATRAZINE TANK MIXTURES

A tank mixture of Campaign plus atrazine may be used on bermudagrass or bahiagrass. For the proper application timing and Campaign rate selection see the directions for CAMPAIGN ALONE in the respective bermudagrass and bahiagrass sections above. For this tank mixture use 1.5 to 3 pounds of atrazine per acre. Refer to the atrazine label for weeds controlled by atrazine. Do not use this tank mixture on tall fescue.

RECOMMENDED RATES AND WEEDS CONTROLLED

Rate recommendations for specific weed species are given below

Use the lower rate when treating annual weeds below 6 inches in height. Use the higher rate on weeds taller than 6 inches or as they approach flower or seedhead formation. Use the higher rate of this product for control or partial control of perennial species. Use the lower rates for growth suppression of perennials.

For the best rate recommendation on the mixture of weeds within your geographic area, contact your local Monsanto representative.

120618

(12)

WEEDS CONTROLLED OR SUPPRESSED
 Note C = Control
 S = Suppression

ANNUAL WEED SPECIES	CAMPAIGN# 02-AGRE			
	1	2	3	4
Barley, little <i>Hordeum pusillum</i>	S	S	C	C
Bedstraw, catchweed <i>Galium aparine</i>	S	S	C	C
Bluegrass, annual <i>Poa annua</i>	S	S	C	C
Charvil <i>Chaerophyllum tainturieri</i>	S	S	C	C
Chickweed, common <i>Stellaria media</i>	S	S	C	C
Clover, crimson <i>Trifolium incarnatum</i>	•	S	S	SC
Clover, lesp <i>Trifolium spp</i>	•	S	SC	SC
Crabgrass <i>Digitaria spp</i>	•	S	C	C
Festail <i>Setaria spp.</i>	•	S	C	C
Geranium, Carolina <i>Geranium carolinianum</i>	S	SC	SC	SC
Heabit <i>Lamium amplexicaule</i>	S	SC	SC	C
Partridgepea <i>Cassia fasciculata</i>	•	S	C	C
Ragweed, common <i>Ambrosia artemisiifolia</i>	•	S	SC	C
Spewdwell, corn <i>Veronica arvensis</i>	S	S	C	C
Spurge, spotted <i>Euphorbia maculata</i>	•	S	C	C
Vervain, blue <i>Verbena hastata</i>	•	S	C	C
BIENNIAL WEED SPECIES				
Carrot, wild <i>Daucus carota</i>	•	•	S	C
PERENNIAL WEED SPECIES				
Fescue, tall <i>Festuca arundinacea</i>	S	S	S	S
Greenbrier <i>Smilax spp.</i>	•	•	•	S
Honeylocust <i>Lonicera spp.</i>	•	S	S	C
Horsebottle <i>Solanum carolinense</i>	•	•	S	S
Johnsongrass <i>Sorghum halepense</i>	•	S	S	C
Minktail, buckhorn <i>Plantago lanceolata</i>	S	VS	SC	SC
Raspberry <i>Rubus spp.</i>	•	•	S	

Add:
 CAMPAIGN# ALONE

Change to:
 Pints/Acre
 1.5 2 3 4

Add:
 Vetch • S C C
Vicia spp.

Bahiagrass S S S S
Paspalum notatum

Dock (early) • S C C
Rumex crispus

Peppervine • S C C
Ampelopsis arborea

PERENNIAL WEED SPECIES				
Ryegrass, Italian	•	•	S	← Replace with Ryegrass <u>Lolium spp.</u>
Lolium multiflorum				
Trumpetcrueper	•	•	S	
Campsis radicans				
Vasegrass	S	C	C	
Paspalum urvillei				
Vetch, common	•	•		← Delete
Vase-erue				

• • • S

WEEDS CONTROLLED OR SUPPRESSED
CAMPAIGN® plus OUST

Note: C = Control

S = Suppression

ANNUAL WEED SPECIES	CAMPAIGN (PINTS/ACRE)	1.5	2	3	3	4	4	4
		+	+	+	+	+	+	+
	OUST (OZ/ACRE)	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
Barley, little <i>Hordeum pusillum</i>		C	C	C	C	C	C	C
Bedstraw, catchweed <i>Galium aparine</i>		C	C	C	C	C	C	C
Bluegrass, annual <i>Poa annua</i>		S	S	C	C	C	C	C
Chervil <i>Chaerophyllum tainturieri</i>		C	C	C	C	C	C	C
Chickweed, common <i>Stellaria media</i>		S	S	C	C	C	C	C
Clover, crimson <i>Trifolium incarnatum</i>		S	S	S	C	C	C	C
Clover, hop <i>Trifolium spp.</i>		•	S	C	C	C	C	C
Crabgrass <i>Digitaria spp.</i>		•	S	C	C	C	C	C
Foxtail <i>Setaria spp.</i>		•	S	C	C	C	C	C
Geranium, Carolina <i>Geranium carolinianum</i>		S	C	C	C	C	C	C
Henbit <i>Lamium amplexicaule</i>		S	C	C	C	C	C	C
Partridgepea <i>Cassia fasciculata</i>		•	S	C	C	C	C	C

Ragweed, common <i>Ambrosia artemisiifolia</i>	•	S	C	C	C	C	C
Speedwell, corn <i>Veronica arvensis</i>	S	S	C	C	C	C	C
Spurge, spotted <i>Euphorbia maculata</i>	•	S	C	C	C	C	C
Vervain, blue <i>Verbena hastata</i>	•	S	C	C	C	C	C
Vetch <i>Vicia</i> spp.	•	S	C	C	C	C	C

BIENNIAL WEED
SPECIES

Carrot, wild <i>Daucus carota</i>	•	•	S	S	C	C	C
--------------------------------------	---	---	---	---	---	---	---

PERENNIAL WEED
SPECIES

Bahiagrass <i>Paspalum notatum</i>	S	S	S	S	S	S	S
Dock, curly <i>Rumex crispus</i>	•	S	C	C	C	C	C
Fescue, tall <i>Festuca arundinacea</i>	S	S	S	S	S	S	S
Greenbriar <i>Smilax</i> spp.	•	•	•	•	S	S	S
Honeysuckle <i>Lonicera</i> spp.	•	S	C	C	C	C	C
Horsenettle <i>Solanum carolinense</i>	•	•	S	S	S	S	S
Johnsongrass <i>Sorghum halepense</i>	•	S	S	S	C	C	C
Peppervine <i>Ampelopsis arborea</i>	•	S	C	C	C	C	C
Plantain, buckhorn <i>Plantago lanceolata</i>	S	S	C	C	C	C	C
Raspberry <i>Rubus</i> spp.	•	•	•	•	S	S	S

Ryegrass, Italian • • S S C C C
 Lolium spp.

Trumpet creeper • S C C C C C
 Campsis radicans

Vaseygrass • S C C C C C
 Paspalum urvillei

APPLICATION EQUIPMENT

This product may be applied using ground spray equipment. Use extreme care to avoid misting or drifting of herbicide solution onto foliage, green stems or fruit of desirable crops, trees, or plants during both growing and dormant periods since even very small quantities of spray can cause severe plant injury.

BROADCAST APPLICATION: Apply recommended rates of this product in 15 to 30 gallons of water per acre as a broadcast spray. For optimum spray distribution and coverage, use flat fan or low-volume flood nozzles. When using flood nozzles, space them no more than 40 inches apart and ensure double overlap of spray pattern. Refer to the manufacturer's recommendations for correct pressure and nozzle height above the target canopy. Avoid pressure and nozzles which produce fine droplets or mist.

MIXING INSTRUCTIONS

Fill the spray tank to about 3/4 of the desired volume with clean water. Add the recommended amount of this product, then complete the filling process while maintaining agitation. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source.

During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, terminate by-pass and return lines at the tank bottom and/or use an agriculturally approved anti-foam or defoaming agent.

Additional surfactant is not necessary for this formulation.

NOTE: Reduced results may occur if water containing silt is used, such as water from ponds and unlined ditches.

Add:

CAMPAIGN® ALONE

Add:

CAMPAIGN TANK MIXTURES WITH OUST OR ATRAZINE

Always predetermine the compatibility of labeled tank mixtures of this herbicide with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of Campaign herbicide with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with water and start agitation.
3. Add the required amount of Oust or atrazine slowly while maintaining agitation.
4. Continue filling the sprayer tank with water and add the required amount of Campaign herbicide near the end of the filling process.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near bottom of tank to minimize foaming.

SPRAYER CLEANUP

CLEAN THE ENTIRE SPRAYER AFTER APPLICATION OF THIS PRODUCT. Add clean water to the tank and thoroughly rinse the entire sprayer system, then fill the tank with water and ammonia. Add 1 quart of household ammonia per 25 gallons of water. Pump enough solution through the hoses, boom and nozzles to fill these parts completely. Then fill the tank, close and leave for 24 hours before draining and rinsing thoroughly with water.

897 16-000 01/53

MONSANTO COMPANY 1988

In case of emergency involving this product
Call Collect, day or night, (314) 694-4000

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.



Change Lot:

897 16-000.08/53

Reg no: 524-351

PM-25

1918
①



HERBICIDE by Monsanto

For control or suppression of emerged grass and broadleaf weeds, woody brush and vines in bermudagrass roadside and highway rights-of-way in Alabama, Arkansas, Florida, Georgia, Louisiana, New Mexico, Mississippi, Oklahoma, South Carolina, and Texas.

Change to:

For control or suppression of emerged grasses, broadleaf weeds and vines in bermudagrass, bahiagrass, and tall fescue turf on roadsides, highway rights-of-way, ditchbanks and dry ditches.

Complete Directions for Use

AVOID CONTACT WITH FOLIAGE OF CROP OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

EPA Reg. No. 524-351

Campaign® is a registered trademark of Monsanto Company.

ACCEPTED

23

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-351

1988-1

897.16-000.01/53

1989-1

897.16-000.02/53

BEST AVAILABLE COPY

©MONSANTO COMPANY 1988

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.



Read the entire label before using product.
Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING PROHIBITED.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this Limit Of Warranty and Liability which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

Hazards to
Humans and Domestic Animals

Keep out of reach of children.

DANGER!

CAUSES EYE BURNS.
HARMFUL IF SWALLOWED.
MAY CAUSE ALLERGIC SKIN REACTION.

Do not get in eyes, on skin, or on clothing.

Wear goggles or face shield and rubber gloves in transferring and mixing and when adjusting, repairing or cleaning equipment.

BEST AVAILABLE COPY

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Call a physician.

IF ON SKIN, immediately flush with plenty of water for at least 15 minutes while removing contaminated clothing. As soon as soap is available, wash thoroughly with soap and water. Wash clothing before reuse. Sanitized persons should avoid further contact and reuse of contaminated clothing.

IF SWALLOWED, induce vomiting immediately by either giving two glasses of water and sticking finger down throat or, when available, administering syrup of ipecac as prescribed on its container. Call a physician. **NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.**

In case of an emergency involving this product,
 Call Collect, day or night, (314) 694-4808.

Environmental Hazards

Do not apply directly to water or wetlands (swamps, bogs, marshes or ponds). Do not contaminate water by cleaning of equipment or disposal of wastes.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENTS*

Glyphosate [N-phosphonomethyl glycine] as the isopropylamine salt	12.9%
2,4-D(2,4-dichlorophenoxyacetic acid) as the isopropylamine salt	20.6%
INERT INGREDIENTS:	66.5%
	100.0%

*Contains 144 grams per litre or 1.2 lbs per U.S. gallon of the active ingredient, glyphosate, as its isopropylamine salt and 227 grams per litre or 1.9 lbs per U.S. gallon of the active ingredient, 2,4-D, as its isopropylamine salt. Equivalent to 108 grams per litre or 0.9 lb per U.S. gallon of the acid, glyphosate and 182 grams per litre or 1.5 lbs per U.S. gallon of the acid, 2,4-D.

Product is protected by U.S. Patent No. 3,799,758, and U.S. Patent No. 4,405,531. Other patents are pending.

GENERAL INFORMATION

Campana® is a postemergence herbicide for control or suppression of emerged weeds. This product is recommended for use on highway and roadside rights-of-way.

This product enters the plant through the foliage and moves throughout the plant. Visual effects of control are a gradual wilting or yellowing of the plant, which advances to complete browning of above-ground growth and deterioration of affected underground plant parts.

Change to:

Do not contaminate water when disposing of equipment washwaters.

BEST AVAILABLE COPY

Add:

No license granted under any non-U.S. patent.

Visible symptoms will usually develop on labeled weeds within 2 to 4 days after application, but may not occur for 7 or more days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay the visual effects of control.

APPLICATION PRECAUTIONS AND WARNINGS

- DO NOT APPLY IN THE VICINITY OF 2,4-D SENSITIVE CROPS SUCH AS COTTON, GRAPES, TOMATOES AND OTHER DESIRABLE VEGETATION.
- Applications should be made only when there is no hazard from spray drift since very small quantities of spray, which may not be visible, may severely injure susceptible crops or desirable vegetation.
- The likelihood of injury occurring to adjacent crops from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions including lesser wind velocities will favor spray drift.
- Movement of this product on soil particles during windstorms may cause damage to susceptible plants that are contacted. This hazard is reduced if rainfall occurs shortly after application.
- Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this herbicide or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

← Delete

Campaign is subject to all state and county regulations for 2,4-D amine.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. Do not store near fertilizers, seeds, insecticides or fungicides.

STORAGE:

STORE ABOVE 40°F to keep product in solution. If crystals form, place in a warm room (72°F), allow the product to reach room temperature and roll or shake periodically until crystals have dissolved. Keep container closed to prevent spills and contamination.

DISPOSAL:

Wastes of this pesticide may cause eye burns and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Empty container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

BEST AVAILABLE COPY

Change to:

← ... allow the product to reach room temperature and roll or shake the container periodically...

Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

TIMING OF APPLICATION

This product should be applied postemergence to vigorously growing weeds at a rate recommended in the "RECOMMENDED RATES AND WEEDS CONTROLLED" section of this label. Application should be delayed until maximum emergence of the target weeds but before mature seeds are produced.

Reduced control may result if treatments are made during poor growing conditions such as drought stress, disease or insect damage or if weeds have been mowed, grazed or cut. Heavy dust on foliage or an overstory canopy covering targeted weeds may also reduce control.

Rainfall occurring within 6 hours after application may reduce effectiveness.

RELEASE OF DORMANT BERMUDAGRASS

When applied as directed, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass. To avoid delays in greening and minimize injury, do not treat when bermudagrass is in a semi-dormant condition.

For best results on winter annuals, treat when plants are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for proper rate selection.

RELEASE OF ACTIVELY GROWING BERMUDAGRASS

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual and biennial species listed in the "WEEDS CONTROLLED" section of this label, and control or suppression of certain perennial weeds.

Use the lower rate when treating annual weeds below 6 inches in height (or length of summer in annual weeds). Use the higher rate as size of plants increases or as they approach flower or seedhead formation. Use the higher rate of this product for control or partial control of perennial species. Use the lower rates for suppression of growth. For best results, see the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the proper rate selection.

Use only in areas where bermudagrass is a desirable ground cover and where some temporary injury or discoloration can be tolerated.

RECOMMENDED RATES AND WEEDS CONTROLLED

Apply 32 to 64 fluid ounces of this product in 15 to 30 gallons of spray solution per acre. Rate recommendations for specific weed species are given below.

For the best rate recommendation for the mixture of weeds within your geographic area, contact your local sales representative.

Replace with the following pages:

BEST AVAILABLE COPY

6918
6

~~WEEDS CONTROLLED OR SUPPRESSED~~

Note: C = Control
S = Suppression

ANNUAL WEED SPECIES	CAMPAIGN* OZ/ACRE		
	32	48	64
Barley, little <i>Hordeum pusillum</i>	S	C	C
Bodilow, catchweed <i>Galium aparine</i>	S	C	C
Blugrass, annual <i>Poa annua</i>	S	C	C
Charvill <i>Chaerophyllum tanacetum</i>	S	C	C
Chickweed, common <i>Stellaria media</i>	S	C	C
Clown, crimson <i>Trifolium incarnatum</i>	S	S	S
Clown, hop <i>Trifolium spp.</i>	S	S	S
Craggrass <i>Digitaria spp.</i>	S	C	C
Fertail <i>Setaria spp.</i>	S	C	C
Geranium, Carolina <i>Geranium carolinianum</i>	•	S	S
Hemlock <i>Lamium amplexicaule</i>	•	S	C
Periwinkle <i>Cassia fasciculata</i>	S	C	C
Ragweed, common <i>Artemisia artemisiifolia</i>	S	S	C
Spindwell, corn <i>Lernaea arvensis</i>	S	C	C
Spurge, spotted <i>Euphorbia maculata</i>	S	C	C
Verbena, blue <i>Verbena hastata</i>	S	C	C
BIENNIAL WEED SPECIES			
Carrot, wild <i>Daucus carota</i>	•	S	C
PERENNIAL WEED SPECIES			
Fescue, tall <i>Festuca arundinacea</i>	S	S	S
Greenbrier <i>Smilax spp.</i>	•	•	S
Hemlock <i>Lonicera spp.</i>	S	S	C
Hemlock <i>Sorbus carolinianus</i>	•	S	S
Johnsongrass <i>Sorghum halepense</i>	S	S	C
Plantain, broadleaf <i>Plantago lanceolata</i>	•	S	S
Raspberry <i>Rubus spp.</i>	•	•	S

Replace with the following pages:

BEST AVAILABLE COPY

PERENNIAL WEED SPECIES			
<i>Ryegrass, Italian</i>	.	.	S
<i>Lolium multiflorum</i>	.	.	S
<i>Trisetospora</i>	.	.	S
<i>Corpus radicans</i>	S	C	C
<i>Vaccinium</i>	.	.	S
<i>Paspalum virgatum</i>	.	.	S
<i>Vicia sativa</i>	.	.	S

Replace with the following pages:

BEST AVAILABLE COPY

TANK MIXTURES

CAMPAIGN ALONE
CAMPAIGN + OUST TM
CAMPAIGN + ATRAZINE

BERMUDAGRASS

Read and carefully observe the label claims, cautions, statements, and all information on the labels of both products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Spring applications:

When applied as directed, this product will provide control or suppression of vines and many winter annual weeds for release of bermudagrass. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated.

Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.25 to 1 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

Summer applications:

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines for release of actively growing bermudagrass.

Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.25 to 2 ounces of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

For spring and summer applications to bermudagrass, use only in areas where bermudagrass is a desirable ground cover and where some temporary discoloration can be tolerated.

TM Oust is a TRADEMARK OF E.I. duPont de Nemours and Company.

BAHIAGRASS

Dormant applications:

When applied as directed, this product will provide control or suppression of vines and winter annual weeds for release of dormant bahiagrass. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated. To avoid delays in greenup and minimize injury, treat when bahiagrass is dormant.

Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.25 to 0.5 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

Active applications:

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines for release of bahiagrass.

Apply 1.5 to 2 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 1.5 to 2 pints of this product plus 0.25 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

TALL FESCUE

Spring applications:

When applied as directed, this product will provide control or suppression of vines and many winter annual weeds in tall fescue. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated. To minimize tall fescue injury, apply this product when tall fescue is 4 to 6 inches tall but prior to seedhead emergence.

Apply 2 to 3 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control of certain weeds. For this tank mixture, apply 2 pints of this product plus 0.25 ounces of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

Summer applications:

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines in tall fescue.

Apply 2 to 3 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control of certain weeds. For this tank mixture, apply 2 to 3 pints of this product plus 0.25 to 0.5 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct selection.

CAMPAIGN® plus ATRAZINE TANK MIXTURES

A tank mixture of Campaign plus atrazine may be used on bermudagrass or bahiagrass. For the proper application timing and Campaign rate selection see the directions for CAMPAIGN ALONE in the respective bermudagrass and bahiagrass sections above. For this tank mixture use 1.5 to 3 pounds of atrazine per acre. Refer to the atrazine label for weeds controlled by atrazine. Do not use this tank mixture on tall fescue.

RECOMMENDED RATES AND WEEDS CONTROLLED

Rate recommendations for specific weed species are given below.

Use the lower rate when treating annual weeds below 6 inches in height. Use the higher rate on weeds taller than 6 inches or as they approach flower or seedhead formation. Use the higher rate of this product for control or partial control of perennial species. Use the lower rates for growth suppression of perennials.

For the best rate recommendation on the mixture of weeds within your geographic area, contact your local Monsanto representative.

WEEDS CONTROLLED OR SUPPRESSED				
Note: C = Control S = Suppression				
ANNUAL WEED SPECIES	CAMPAIGN# 02/ADRE			
	32	40	44	46
Barley, wild <i>Hordeum periculum</i>	S	S	C	C
Bedstraw, catchweed <i>Galium aparine</i>	S	S	C	C
Bleeding heart <i>Pas. ovum</i>	S	S	C	C
Charoil <i>Chenopodium tentaculatum</i>	S	S	C	C
Chickweed, common <i>Stellaria media</i>	S	S	C	C
Claw, crimson <i>Trifolium incarnatum</i>	•	S	S	SC
Claw, hop <i>Trifolium spp.</i>	•	S	SC	SC
Cockspur <i>Digitaria spp.</i>	•	S	C	C
Festool <i>Setaria spp.</i>	•	S	C	C
Germinal, Carolina <i>Germinal carolinianum</i>	S	SC	SC	SC
Habit <i>Lamium amplexicaule</i>	S	SC	SC	C
Partridgefoot <i>Cassia fasciculata</i>	•	S	C	C
Ragweed, common <i>Ambrosia artemisiifolia</i>	•	S	SC	C
Spandwell, corn <i>Veronica arvensis</i>	S	S	C	C
Spurge, spotted <i>Euphorbia maculata</i>	•	S	C	C
Vervain, blue <i>Verbena hastata</i>	•	S	C	C
BIENNIAL WEED SPECIES				
Carrot, wild <i>Daucus carota</i>	•	•	S	C
PERENNIAL WEED SPECIES				
Fescue, tall <i>Festuca arundinacea</i>	S	S	S	S
Groundsley <i>Smilax spp.</i>	•	•	•	S
Monarda <i>Leonurus spp.</i>	•	S	S	C
Monarda <i>Salvia caroliniana</i>	•	•	S	S
Johnsongrass <i>Sorghum halepense</i>	•	S	S	C
Plains, broadleaf <i>Plantago lanceolata</i>	S	VS	SC	SC
Rumex <i>Rubus spp.</i>	•	•	S	

Add:
CAMPAIGN# ALONE

Change to:
Pints/Acre
1.5 2 3 4

BEST AVAILABLE COPY

Add:
Vetch • S C C
Vicia spp.

Bahiagrass S S S S
Paspalum notatum

Dock (early)
Rumex crispus • S C C

Peppervine • S C C
Ampelopsis arborea

PERENNIAL WEED
SPECIES

Styragma-Rubra	•	•	•
Lolium-mutillorum			
Trampoliner	•	•	S
Campis radicans			
Vanagras	S	C	C
Populus arctica			
Urtica-cannabina	•	•	•
Urtica-cannabina			

Replace with
Ryegrass
Lolium spp.

Delete

BEST AVAILABLE COPY

14 g 18
(14)

WEEDS CONTROLLED OR SUPPRESSED
CAMPAIGN plus OUST

Note: C = Control

S = Suppression

ANNUAL WEED SPECIES	CAMPAIGN (PINTS/ACRE)	1.5	2	3	3	4	4	4
	OUST (OZ/ACRE)	+	+	+	+	+	+	+
		$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
Barley, little <i>Hordeum pusillum</i>		C	C	C	C	C	C	C
Bedstraw, catchweed <i>Galium aparine</i>		C	C	C	C	C	C	C
Bluegrass, annual <i>Poa annua</i>		S	S	C	C	C	C	C
Chervil <i>Chaerophyllum tainturieri</i>		C	C	C	C	C	C	C
Chickweed, common <i>Stellaria media</i>		S	S	C	C	C	C	C
Clover, crimson <i>Trifolium incarnatum</i>		S	S	S	C	C	C	C
Clover, hop <i>Trifolium spp.</i>		•	S	C	C	C	C	C
Crabgrass <i>Digitaria spp.</i>		•	S	C	C	C	C	C
Foxtail <i>Setaria spp.</i>		•	S	C	C	C	C	C
Geranium, Carolina <i>Geranium carolinianum</i>		S	C	C	C	C	C	C
Henbit <i>Lamium amplexicaule</i>		S	C	C	C	C	C	C
Partridgepea <i>Cassia fasciculata</i>		•	S	C	C	C	C	C

Ragweed, common <i>Ambrosia artemisiifolia</i>	• S C C C C C C
Speedwell, corn <i>Veronica arvensis</i>	S S C C C C C
Spurge, spotted <i>Euphorbia maculata</i>	• S C C C C C C
Vervain, blue <i>Verbena hastata</i>	• S C C C C C C
Vetch <i>Vicia spp.</i>	• S C C C C C C

BIENNIAL WEED
SPECIES

Carrot, wild <i>Daucus carota</i>	• • S S C C C
--------------------------------------	---------------

PERENNIAL WEED
SPECIES

Bahiagrass <i>Paspalum notatum</i>	S S S S S S S
Dock, curly <i>Rumex crispus</i>	• S C C C C C
Fescue, tall <i>Festuca arundinacea</i>	S S S S S S S
Greenbriar <i>Smilax spp.</i>	• • • • S S S
Honeysuckle <i>Lonicera spp.</i>	• S C C C C C
Horsenettle <i>Solanum carolinense</i>	• • S S S S S
Johnsongrass <i>Sorghum halepense</i>	• S S S C C C
Peppervine <i>Ampelopsis arborea</i>	• S C C C C C
Plantain, buckhorn <i>Plantago lanceolata</i>	S S C C C C C
Raspberry <i>Rubus spp.</i>	• • • • S S S

Rvegrass, Italian
 Lolium spp.

• • S S C C C

Trumpet creeper
 Campsis radicans

• S C C C C C

Vaseygrass
 Paspalum urvillei

• S C C C C C

(

(

APPLICATION EQUIPMENT

This product may be applied using ground spray equipment. Use extreme care to avoid misting or drifting of herbicide solution onto foliage, green stems or fruit of desirable crops, trees, or plants during both growing and dormant periods since even very small quantities of spray can cause severe plant injury.

GROUND APPLICATION: Apply recommended rates of this product in 15 to 30 gallons of water per acre as a broadcast spray. For optimum spray distribution and coverage, use flat fan or low-volume flood nozzles. When using flood nozzles, space them no more than 40 inches apart and ensure double overlap of spray pattern. Refer to the manufacturer's recommendations for correct pressure and nozzle height above the target canopy. Avoid pressure and nozzles which produce fine droplets or mist.

MIXING INSTRUCTIONS

Fill the spray tank to about 3/4 of the desired volume with clean water. Add the recommended amount of this product, then complete the filling process while maintaining agitation. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source.

During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, terminate by-pass and return lines at the tank bottom and/or use an agriculturally approved anti-foam or defoaming agent.

Additional surfactant is not necessary for this formulation.

NOTE: Reduced results may occur if water containing silt is used, such as water from ponds and unlined ditches.

Add:
CAMPAIGN® ALONE

BEST AVAILABLE COPY

Add:
CAMPAIGN TANK MIXTURES WITH OUST OR ATRAZINE

Always predetermine the compatibility of labeled tank mixtures of this herbicide with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of Campaign herbicide with water as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with water and start agitation.
3. Add the required amount of Oust or atrazine slowly while maintaining agitation.
4. Continue filling the sprayer tank with water and add the required amount of Campaign herbicide near the end of the filling process.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near bottom of tank to minimize foaming.

SPRAYER CLEANUP

CLEAN THE ENTIRE SPRAYER AFTER APPLICATION OF THIS PRODUCT. Add clean water to the tank and thoroughly rinse the entire sprayer system, then fill the tank with water and ammonia. Add 1 quart of household ammonia per 25 gallons of water. Pump enough solution through the boom, boom and nozzles to fill these parts completely. Then fill the tank, close and leave for 24 hours before draining and rinsing thoroughly with water.

897.16-000.01/53

©MONSANTO COMPANY 1968

In case of emergency involving this product,
Call Collect, day or night, (314) 694-4000.

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.



Change to:

897.16-000.08/53

BEST AVAILABLE COPY