

CAMPAIGN

HERBICIDE by Monsanto

For control or suppression of emerged grasses, broadleaf weeds and vines in bermudagrass, bahiagrass, and tall fescue turf on roadsides, highway rights-of-way, ditchbanks and dry ditches

Complete Directions for Use

AVOID CONTACT WITH FOLIAGE OF CROP OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT

EPA Reg. No. 524-35

Campan® is a trademark of Monsanto Company

1969 1 897 1 1018 53

MONSANTO COMPANY 1984

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

524-351 PM 25 1064

Use only according to label instructions

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OF SET ALKALINE PROHIBITED

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms with the chemical description on the label and is reasonably fit for the purpose set forth in the Complete Directions for Use label booklet. Directions are contained in a booklet with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is subject to conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with product, other than those set forth in the Directions; application to or contact with desirable vegetation; unusual weather; weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied; as well as weather conditions which are outside the application ranges set forth in the Directions; application in any manner not explicitly set forth in the Directions; moisture conditions outside the moisture range specified in the Directions; or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this Limit Of Warranty and Liability which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children

DANGER!

CAUSES EYE BURNS.
HARMFUL IF SWALLOWED.
MAY CAUSE ALLERGIC SKIN REACTION

Do not get in eyes, on skin, or on clothing

Wear goggles or face shield and rubber gloves in transferring and mixing and when adjusting, repairing or cleaning equipment

524-351 PM 25 1064

Wash thoroughly with soap and water after handling.
FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Call a physician.

IF ON SKIN, immediately flush with plenty of water for at least 15 minutes while removing contaminated clothing. As soon as practicable, wash thoroughly with soap and water. Do not use oil-based or sensitized personal care products after contact and reuse of containers.

IF SWALLOWED, do not induce vomiting. Immediately call a physician. If unable to do so, call a poison control center or hospital. Do not give anything by mouth to an unconscious person. NEVER use a pump or syringe to force anything into the mouth.

Use of this product is restricted to the following uses: Control of weeds in bermudagrass, bahiagrass, and tall fescue turf on roadsides, highway rights-of-way, ditchbanks and dry ditches.

Environmental Hazards

Do not apply directly to water or wetland. Do not apply to marshes or other wetlands. Do not apply to water when the use of equipment will be required.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT (OR SPRAY SOLUTIONS OF THIS PRODUCT) IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT FOR PRESS STEEL CONTAINERS OR SPRAY TANKS). This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENTS*

Glyphosate [N-phosphonomethylglycine] as the isopropylamine salt	12.9%
2,4-D [2,4-dichlorophenoxyacetic acid] as the isopropylamine salt	20.6%
INERT INGREDIENTS	66.5%
	100.0%

*Contains 144 grams per litre or 1.2 lbs per U.S. gallon of the active ingredient, glyphosate, as its isopropylamine salt and 227 grams per litre or 1.9 lbs per U.S. gallon of the active ingredient, 2,4-D, as its isopropylamine salt. Equivalent to 106 grams per litre or 0.9 lb per U.S. gallon of the acid, glyphosate and 182 grams per litre or 1.5 lbs per U.S. gallon of the acid, 2,4-D.

Product protected by U.S. Patent No. 3,799,758 and U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent.

GENERAL INFORMATION

This product is a post-emergent herbicide for control or suppression of emerged weeds. This product is recommended for use on highway, roadside rights-of-way.

This product enters the plant through the foliage and moves throughout the plant. Visual effects of control are a gradual wilting or yellowing of the plant, which advances to complete browning of above ground growth and deterioration of affected underground plant parts.

BEST AVAILABLE COPY

Visible symptoms will usually develop on labeled weeds within 2 to 4 days after application, but may not occur for 7 or more days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay the visual effects of control.

APPLICATION PRECAUTIONS AND WARNINGS

- DO NOT APPLY IN THE VICINITY OF 2,4-D SENSITIVE CROPS SUCH AS COTTON, GRAPES, TOMATOES AND OTHER DESIRABLE VEGETATION.
- Application should be made only when there is no hazard from spray drift to susceptible crops or desirable vegetation. Heavy drift of spray which may not be visible may severely injure susceptible crops or desirable vegetation.
- The likelihood of injury occurring to adjacent crops from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions including lesser wind velocities will favor spray drift.
- Buyer and all users are responsible for all injury or damage in connection with the use or handling of mixtures of this herbicide or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops or other unintended consequences.

Campaign is subject to all state and county regulation for 2,4-D amine.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal. Do not store near fertilizers, seeds, insecticides or fungicides.

STORAGE

STORE ABOVE 40°F to keep product in solution. If crystals form, place in a warm room (72°F), allow the product to reach room temperature and roll or shake the container periodically until crystals have dissolved. Keep container closed to prevent spills and contamination.

DISPOSAL

Wastes of this pesticide may cause eye burns and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Empty container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

TIMING OF APPLICATION

This product should be applied postemergence to vigorously growing weeds at a rate recommended in the

"RECOMMENDED RATES AND WEEDS CONTROLLED" section of this label. Application should be delayed until maximum emergence of the target weeds but before mature seeds are produced.

Reduced control may result if treatments are made during poor growing conditions such as drought stress, disease or insect damage, or if weeds have been mowed, grazed or cut. Heavy growth of foliage on an overstory canopy may also reduce control.

Heavy fall cutting within 6 hours after application may reduce effectiveness.

TANK MIXTURES

CAMPAIGN[®] ALONE

CAMPAIGN plus OUST[™]

CAMPAIGN plus ATRAZINE

BERMUDAGRASS

Read and carefully observe the label claims, cautions, statements, and all information on the labels of both products used in these tank mixtures. Use according to the most restrictive label directions for each product in the mixture.

Spring Applications:

When applied as directed, this product will provide control or suppression of vines and many winter annual weeds for release of bermudagrass. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated.

Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.25 to 1 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

Summer Applications:

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines for release of actively growing bermudagrass.

Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.25 to 2 ounces of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

For spring and summer applications to bermudagrass use only in areas where bermudagrass is a desirable ground cover and where some temporary discoloration can be tolerated.

*Oust is a trademark of E.I. du Pont de Nemours and Company.

BERMUDAGRASS

Spring Applications:

When applied as directed, this product will provide control or suppression of vines and winter annual weeds for release of dormant bermudagrass. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated. To avoid delays in greenup and minimize injury, treat when bermudagrass is dormant.

Apply 2 to 4 pints of Campaign herbicide in 15 to 30 gallons of spray solution per acre. A tank mixture of Campaign plus Oust may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.25 to 0.5 ounce of Oust per acre. See the "RECOMMENDED RATES AND WEEDS CONTROLLED" section for the correct rate selection.

BEST AVAILABLE COPY

Active Applications

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines for release of bahiagrass.

Apply 1.5 to 2 pints of Campaign herbicide to bahiagrass in the form of spray solution per acre. A tank mixture of Campaign herbicide plus atrazine may be used to provide control or partial control of perennials in the form of spray solution per acre. Refer to the atrazine label for weed species controlled by atrazine. Do not use this tank mixture on tall fescue.

TALL FESCUE

Spring Applications

Apply 1.5 to 2 pints of Campaign herbicide to bahiagrass in the form of spray solution per acre. A tank mixture of Campaign herbicide plus atrazine may be used to provide control or partial control of perennials in the form of spray solution per acre. Refer to the atrazine label for weed species controlled by atrazine. Do not use this tank mixture on tall fescue.

Apply 2 to 3 pints of Campaign herbicide to bahiagrass in the form of spray solution per acre. A tank mixture of Campaign herbicide plus atrazine may be used to provide control or partial control of perennials in the form of spray solution per acre. Refer to the atrazine label for weed species controlled by atrazine. Do not use this tank mixture on tall fescue.

Summer Applications

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines in tall fescue.

Apply 2 to 3 pints of Campaign herbicide to bahiagrass in the form of spray solution per acre. A tank mixture of Campaign herbicide plus atrazine may be used to provide control or partial control of perennials in the form of spray solution per acre. Refer to the atrazine label for weed species controlled by atrazine. Do not use this tank mixture on tall fescue.

TANK MIXTURE

CAMPAIGN® plus ATRAZINE

A tank mixture of Campaign plus atrazine may be used on bermudagrass or bahiagrass for the proper application timing and Campaign rate selection see the directions for CAMPAIGN ALONE in the respective bermudagrass and bahiagrass sections above. For this tank mixture, use 1.5 to 3 pounds of atrazine per acre. Refer to the atrazine label for weed species controlled by atrazine. Do not use this tank mixture on tall fescue.

RECOMMENDED RATES AND WEEDS CONTROLLED

Rate recommendations for specific weed species are given below.

Use the lower rate when treating annual weeds below 6 inches in height. Use the higher rate on weeds taller than 6 inches or as they approach flower or seedhead formation. Use the higher rate of this product for control or partial control of perennial species. Use the lower rates for growth suppression of perennials.

For the best rate recommendation on the mixture of weeds within your geographic area, contact your local Monsanto representative.

WEEDS CONTROLLED OR SUPPRESSED CAMPAIGN™ ALONE

W = Control
S = Suppression

CAMPAIGN® PLUS ATRAZINE

ANNUAL WEED SPECIES	W	S	C	C	C	C
Barley, little						
Bedstraw, catchweed						
Bigleaf, annual						
Chenop						
Chickweed, common						
Clover, crimson						
Clover, hop						
Crabgrass						
Foxtail						
Geranium, Carolina						
Henbit						
Partridgepea						
Ragweed, common						
Speedwell, corn						
Spurge, spotted						
Vervain, blue						
Vetch						

BIENNIAL WEED SPECIES	W	S	C	C	C	C
Carrot, wild						

PERENNIAL WEED SPECIES	W	S	C	C	C	C
Bahiagrass						
Dock, early						
Fescue, tall						
Greenbrier						
Honeysuckle						
Morsonettle						

Johnsongrass						
Peppercorn						
Plaintain, buckhorn						
Raspberry						
Ryegrass						
Trumpet creeper						
Vaseygrass						

WEEDS CONTROLLED OR SUPPRESSED CAMPAIGN® plus ATRAZINE

ANNUAL WEED SPECIES	W	S	C	C	C	C
Barley, little						
Bedstraw, catchweed						
Bluegrass, annual						
Chenop						
Chickweed, common						
Clover, crimson						
Clover, hop						
Crabgrass						
Foxtail						
Geranium, Carolina						
Henbit						
Partridgepea						
Ragweed, common						
Speedwell, corn						
Spurge, spotted						
Vervain, blue						
Vetch						

BIENNIAL WEED SPECIES	W	S	C	C	C	C
Carrot, wild						

BEST AVAILABLE COPY

[illegible]

This product may be applied using ground spray equipment. Use extreme care to avoid misting or drifting of herbicide solution onto foliage, green stems or fruit of desirable crops, trees, or plants during both growing and dormant periods since even very small quantities of spray can cause severe plant injury.

MIXING INSTRUCTIONS

CAMPAIGN™ ALONE

- Additional surfactant is not necessary for this formulation.

897 16 000 CR 55

MONSANTO COMPANY 1980

In case of emergency involving this product
Call Collect, day or night. (214) 694-4000

MONSANTO COMPANY -
AGRICULTURAL PRODUCTS
ST LOUIS, MISSOURI, 63167 U.S.A.



BES 6 COPY