




For suppression of emerged field bindweed and control or suppression of annual weeds in fallow and reduced tillage small grains systems.

Complete Directions for Use

AVOID CONTACT WITH FOLIAGE OF CROP OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

EPA Reg. No 524-351

*Landmaster is a registered trademark of Monsanto Company.



ANNUAL WEED SPECIES	RATE PER ACRE	MAXIMUM HEIGHT
Barnyardgrass <i>Echinochloa crus-galli</i>	54 oz	6"
Buffalober <i>Solanum rostratum</i>		
Coatgrass <i>Aegilops cylindrica</i>		
Mustard, blue <i>Chorispora tenella</i>		
Panicum, fall <i>Panicum dichotomiflorum</i>		
Oats, wild <i>Avena fatua</i>	54 oz	12"
Witchgrass <i>Panicum capillare</i>		

*For improved control in no-till systems or heavy infestations or overwintered stands, use 54 ounces. For best results on light Kochia infestations, treat after the plant has passed through the woolly stage of growth and is 3 to 6 inches in height. When treating medium to heavy infestations or plants that are in the woolly stage (1 to 3 inches in height), add 2 fluid ounces of Banvel™ to the recommended rate of Landmaster BW. Refer to the Banvel label for planting, cropping and other restrictions. Follow all precautions on the Banvel label.

**Landmaster BW may also be used as a 1 percent solution on a spray-to-wet basis for annual weed control and field bindweed suppression. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Prepare the desired volume of spray by mixing the amount of Landmaster BW in clean water as shown in the following table:

SPRAY SOLUTIONS	
DESIRED VOLUME	AMOUNT OF LANDMASTER® BW
1 gallon	1-1/3 ounces
25 gallons	1 quart
100 gallons	1 gallon
2 tablespoons = 1 ounce	

*Banvel is a trademark of Sandoz, Inc.

POSTHARVEST APPLICATIONS

This product will provide control of weeds following grain harvest. Weeds should be allowed to regrow after damage incurred during harvest operations and to recover from environmental stress before application of this product. Weeds should be treated prior to the heading stage of annual grasses and before broadleaf weeds exceed 24 inches in height. Ammonium sulfate will improve performance on annual weeds under stress conditions.

Weeds controlled at the 40-ounce-per-acre rate include downy brome, green foxtail, stinkgrass, and volunteer wheat.

Weeds controlled at the 54-ounce-per-acre rate include field bindweed, Kochia, lambsquarters, mustard, pigweed and Russian thistle.

Weeds controlled at the 64-ounce-per-acre rate include barnyardgrass, sandbar, witchgrass, yellow foxtail and prickly lettuce.

BEST AVAILABLE COPY

INSERT "ECOFARMING SYSTEMS" SECTION.

INSERT THE FOLLOWING SECTION:

ECOFARMING SYSTEMS

The Ecofarming System consists of the following rotation:
winter wheat, corn/sorghum, ecofallow.

Use the following tank mixtures for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System.

- LANDMASTER Ⓞ BW at 54 to 64 ounces per acre
plus
- ATRAZINE at 0.75 to 1 pound a.i. per acre
plus
- LASSO Ⓞ EC at 2.5 to 3 quarts per acre

The above tank mixture should be applied in 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre. The liquid fertilizer may be diluted with water to achieve the proper carrier volume.

WEEDS CONTROLLED - The following weeds will be controlled up to a maximum height of 4 inches:

Brome, downy
Bromus tectorum

Cheat
Bromus secalinus

Foxtail, green
Setaria viridis

Foxtail, yellow
Setaria lutescens

Kochia*
Kochia scoparia

Lettuce, prickly
Lactuca serriola

Pigweed, redroot
Amaranthus retroflexus

Thistle, Russian
Salsola kali

Wheat, volunteer
Triticum aestivum

BEST AVAILABLE COPY

*For improved control of kochia, add 4 ounces per acre (0.125 pound a.i. per acre) of Banvel to the above tank mixture.

Risk of crop injury from 2,4-D or Banvel can be reduced by applying this treatment 7 to 14 days before planting.

Refer to the Lasso label for preemergence weed control achieved by this tank mixture.

Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products in these mixtures.

Ⓢ Lasso is a registered trademark of Monsanto Company.



RECOMMENDED RATES AND WEEDS CONTROLLED

For best results, apply this product after most weed seeds have germinated but before seedhead formation in grasses or flower bud formation in broadleaves.

When applied as directed, this product will provide control or suppression of the grass and broadleaf weed species listed below. Rates recommended are for maximum weed height at treatment time.

PERENNIAL WEED SPECIES	LANDMASTER® BW			
	RATE PER ACRE	LENGTH OF VINES/ TREATMENT HEIGHT		
Bindweed, field** <i>Convolvulus</i> <i>arvensis</i>	54 oz (Suppression only)	6 to 18"		
ANNUAL WEED SPECIES	RATE PER ACRE	MAXIMUM HEIGHT		
Furtail, green <i>Setaria verticis</i>	27 oz	12"		
Barley <i>Hordeum vulgare</i>	40 oz	6"		
Bromus, downy* <i>Bromus tectorum</i>				
Cheese* <i>Bromus secalinus</i>				
Furtail <i>Setaria spp</i>				
Kochia* <i>Kochia scoparia</i>				
Lettuce, prickly* <i>Lactuca scariola</i>				
Oats, wild <i>Avena fatua</i>				
Puncturevine <i>Tribulus terrestris</i>				
Portulaca, common <i>Portulaca oleracea</i>				
Cocklebur <i>Xanthium strumarium</i>			40 oz	12"
Lambquarters <i>Chenopodium album</i>				
Mustard, tansy <i>Descurainia pinnata</i>				
Mustard, tumble <i>Sisymbrium altissimum</i>				
Pigweed, redroot <i>Amaranthus retroflexus</i>				
Pigweed, smooth <i>Amaranthus hybridus</i>				
Rye <i>Secale cereale</i>				
Stinkgrass <i>Eragrostis chloranthera</i>				
Thistle, Russian <i>Salsola kali</i>				
Wheat <i>Triticum aestivum</i>				
Wildgrass <i>Panicum capillare</i>				

INSERT:

Spruce, Leafywee 54 oz post bloom
Euphorbia esula

BEST AVAILABLE COPY

ANNUAL WEED SPECIES	RATE PFR ACRE	MAXIMUM HEIGHT
Barryardgrass <i>Echinochloa crus-galli</i>	54 oz	6"
Buffalober <i>Solanum rostratum</i>		
Goatgrass <i>Aegilops cylindrica</i>		
Mustard, blue <i>Chorispora tenella</i>		
Panicum, fall <i>Panicum dichotomiflorum</i>		
Oats, wild <i>Avena fatua</i>	54 oz	12"
Witchgrass <i>Panicum capillare</i>		

*For improved control in no-till systems or heavy infestations or overwintered stands, use 54 ounces. For best results on light locoweed infestations, treat after the plant has passed through the woolly stage of growth and is 3 to 6 inches in height. When treating medium to heavy infestations or plants that are in the woolly stage (1 to 3 inches in height), add 2 fluid ounces of Banvel™ to the recommended rate of Landmaster BW. Refer to the Banvel label for planting, cropping and other restrictions. Follow all precautions on the Banvel label.

**Landmaster BW may also be used as a 1 percent solution on a spray-to-wet basis for annual weed control and field bindweed suppression. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Prepare the desired volume of spray by mixing the amount of Landmaster BW in clean water as shown in the following table:

SPRAY SOLUTIONS	
DESIRED VOLUME	AMOUNT OF LANDMASTER® BW
1 gallon	1-1/3 ounces
25 gallons	1 quart
100 gallons	1 gallon
2 tablespoons = 1 ounce	

™Banvel is a trademark of Santex, Inc.

POSTHARVEST APPLICATIONS

This product will provide control of weeds following grain harvest. Weeds should be allowed to regrow after damage incurred during harvest operations and to recover from environmental stress before application of this product. Weeds should be treated prior to the heading stage of annual grasses and before broadleaf weeds exceed 24 inches in height. Ammonium sulfate will improve performance on annual weeds under stress conditions.

Weeds controlled at the 40-ounce-per-acre rate include downy brome, green foxtail, stinkgrass, and volunteer wheat.

Weeds controlled at the 54-ounce-per-acre rate include field bindweed, lactuca, lambsquarters, mustard, pigweed and Russian thistle.

Weeds controlled at the 64-ounce-per-acre rate include barryardgrass, sandbar, witchgrass, yellow foxtail and prickly lettuce.

INSERT:

*** When applied for control in pastures, remove domestic livestock before application and wait 8 weeks before grazing or harvesting.

