YHOS TOURSON

IOPE.

JUN 17 1993

Ms. Lydia A. Suba Monsanto Company The Agricultural Group 700 14th Street, N.W. Suite #1100 Washington, D. C. 20005

Dear Ms. Suba:

Subject: Campaign Herbicide (Control Musk Thistle In Industrial, Rangeland, and Pasture Sites in Colorado, Idaho, Montana, Nebraska, Utah, And Wyoming) EPA Registration No. 524-351 Your Application Dated May 27, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable. Please submit five (5) copies of your final printed labeling, before you release the product for shipment. A stamped copy of the label is enclosed for your records.

The accceptance of your label does not relieve you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR Part 156, subpart K, 200 (c)(3), you are prohibited from distrubuting or selling any product within the scope of WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely,

J43

Robert J. Taylor Product Manager 25 Fungicide-Herbicide Branch Registration Division (H7505C)

### SUPPLEMENTAL LABELING

1-17:0

READ AND FOLLOW THE ENTIRE LABEL FOR CAMPAIGN® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label for Campaign herbicide, the label for any tank mix product and this supplement.

### CAMPAIGN<sup>®</sup> HERBICIDE BY MONSANTO

(EPA REG NO. 524-351)

CAMPAIGN HERBICIDE FOR THE CONTROL OF MUSK THISTLE AND OTHER LABELED WEEDS FOUND IN INDUSTRIAL, RANGELAND AND PASTURE SITES IN COLORADO, IDAHO, MONTANA, NEBRASKA, UTAH AND WYOMING.

Keep out of reach of children.

DANGER!

X

In case of emergency involving this product, Call Collect, day or night, (314)-694-4000.

<sup>\*</sup>Campaign is a registered trademark of Monsanto Company.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

This product is recommended for musk thistle control in rangeland, pasture and noncrop areas. It is also recommended for the control of those weeds listed on the product label booklet in these sites.

Do not graze lactating dairy animals on treated grass within 7 days after application. Animals being finished for slaughter that are grazing in the treated area within 30 days of treatment must be removed from the treated area 3 days before slaughter. Do not cut forage for hay within 30 days of application. When this product is used for spot treatments where less than 10 percent of the total grazed area will be treated, there is no grazing restriction. When using tank mixtures, refer to the tank mix product labe? for grazing restrictions for that product. Follow the mcct restrictive label.

Best control of musk thistle is obtained when applications are made while plants are in the rosette stage of growth. Applications should be made in the spring prior to bolting or in the fall prior to soil freeze-up. Partial control or suppression may be obtained with treatments made from bolting through flowering. For best results, plants should be actively growing and not under stress and free of dust layers on leaves.

For broadcast applications, apply 16 to 32 fluid ounces per acre of this product alone or as a tank mixture with either 4 to 8 fluid ounces of Banvel<sup>M</sup>, 4 to 8 fluid ounces of Tordon<sup>M</sup> 22K or 0.1 to 0.3 ounces of Escort<sup>M</sup>.

For spray-to-wet applications, apply this product as a 1 to 2 percent solution alone or as a tank mixture with 1/2 to 1 percent Banvel, 1/2 to 1 percent Tordon 22K or with Escort at a rate of 1 ounce of product per 100 gallons of spray solution.

Read and carefully observe the label directions, cautionary statements and all information on the labels of each product used in these tank mixtures.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label for Campaign before using this product. Those terms apply to this supplemental labeling and, if those are not acceptable, return the product unopened at once.

<sup>TM</sup>Banvel is a trademark of Sandoz, Inc. <sup>TM</sup>Escort is a trademark of E. I. du Pont de Nemours and Company. <sup>TM</sup>Tordon is a trademark of DowElanco Products Company.

**©MONSANTO COMPANY 1993** 

1

• .

MONSANTO COMPANY AGRICULTURAL PRODUCTS ST. LOUIS, MISSOURI 63167

ACCEPTED with COMMENTS in EPA Letter Dated: ビッチョウ

JUN 1 7 1993

Under the Federal Insectivide, Fungicide, and Rodenticide Act anended, for the pesticide registered under EPA Reg. No.

524-351

JUN 17 1993

Dr. Sheila A. Schuette Monsanto Company The Agricultural Group 700 14th Street, N.W. Suite #1100 Washington, D. C. 20005

Dear Dr. Schuette:

Subject: Landmaster BW Herbicide (Add Pre-Harvest Wheat) EPA Registration No. 524-351 Your Application Dated May 7, 1993

The amendmet referred to above, submitted in connection with registration under FIFRA sec. 3(C)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.

2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

The accceptance of your label does not reliave you of any obligation to comply with the Worker Protection Standard (WPS). Under the WPS labeling regulations at 40 CFR Part 156, subpart K, 200 (c)(3), you are prohibited from distrubuting or selling any product within the scope of WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

Sincerely,

Robert J. Taylor Product Manager 25 Fungicide-Herbicide Branch Registration Division (H7505C) 4076

# SUPPLEMENTAL LABELING

こうもら

READ THE ENTIRE LABEL FOR LANDMASTER<sup>®</sup> BW HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for Landmaster BW and this supplement.

LANDMASTER' BW

Herbicide by Monsanto

ACCEPTED with CONSTENTS in EPA Letter Dated

BPA Reg. No. 524-351

JUN 1 7 1993

Urder the Faleral Insettlede FOR PREHARVEST APPLICATIONS IN WHEAT Funded and holdswidtle Act

registered under EPA Bog. No.

524-451

Keep out of reach of children.

# DANGER!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

•Landmaster is a registered trademark of Monsanto Company.

## DIRECTIONS FOR USB

t is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for Landmaster BW herbicide for essential product performance information.

### RECOMMENDATIONS

This product controls annual and perennial weeds listed on this label prior to the harvest of wheat. Make applications after the hard-dough stage (30% or less grain moisture) and at least 7 days prior to harvest. This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section of this label for instructions for ground and aerial applications.

in strong

For specific rates of application and instructions for control of annual and perennial weeds, see the "WEEDS CONTROLLED" section of this label.

For suppression of field bindweed, apply 54 ounces of this product in 3 to 10 gallons of water per acre.

DO NOT APPLY MORE THAN 84 FLUID OUNCES PER ACRE OF THIS PRODUCT FOR PREHARVEST APPLICATIONS.

Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.

Do not feed treated straw to livestock.

Note: It is not recommended that wheat grown for seed be treated because a reduction in germination or vigor may occur.

Read the "Limit of Warranty and Liability" in the label booklet for Roundup herbicide before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

•Monsanto Company 1992

e

MAP

MONSANTO COMPANY AGRICULTURAL PRODUCTS T. LOUIS, MISSOURI 63167