


READ AND FOLLOW THE ENTIRE LABEL FOR CAMPAIGN® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label for Campaign herbicide, the label for any tank mix product and this supplement.



EPA Reg No. 524-351

CAMPAIGN HERBICIDE FOR THE CONTROL OF MUSK THISTLE AND OTHER LABELED WEEDS FOUND IN INDUSTRIAL, RANGELAND AND PASTURE SITES IN COLORADO, IDAHO, IOWA, KANSAS, MONTANA, NEBRASKA, NORTH DAKOTA, OREGON, UTAH, WASHINGTON AND WYOMING.

Keep out of reach of children.

DANGER!

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

*Campaign is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

This product is recommended for musk thistle control in rangeland, pasture and noncrop areas. It is also recommended for the control of those weeds listed on the product label booklet in these sites.

Do not graze lactating dairy animals on treated grass within 7 days after application. Animals being finished for slaughter that are grazing in the treated area within 30 days of treatment must be removed from the treated area 3 days before slaughter. Do not cut forage for hay within 30 days of application. When this product is used for spot treatments where less than 10 percent of the total grazed area will be treated,

there is no grazing restriction. When using tank mixtures, refer to the tank mix product label for grazing restrictions for that product. Follow the most restrictive label.

Best control of musk thistle is obtained when applications are made while plants are in the rosette stage of growth. Applications should be made in the spring prior to bolting or in the fall prior to soil freeze-up. Partial control or suppression may be obtained with treatments made from bolting through flowering. For best results, plants should be actively growing and not under stress and free of dust layers on leaves.

For broadcast applications, apply 16 to 32 fluid ounces per acre of this product alone or as a tank mixture with either 4 to 8 fluid ounces of Banvel™, 4 to 8 fluid ounces of Tordon™ 22K or 0.1 to 0.3 ounces of Escort™.

For spray-to-wet applications, apply this product as a 1 to 2 percent solution alone or as a tank mixture with 1/2 to 1 percent Banvel, 1/2 to 1 percent Tordon 22K or

with Escort at a rate of 1 ounce of product per 100 gallons of spray solution.

Read and carefully observe the label directions, cautionary statements and all information on the labels of each product used in these tank mixtures.

Read "LIMIT OF WARRANTY AND LIABILITY" in the label for Campaign before using this product. Those terms apply to this supplemental labeling and, if those are not acceptable, return the product unopened at once.

™Banvel is a trademark of Sandoz, inc.
™Escort is a trademark of E. I. du Pont de Nemours and Company.
™Tordon is a trademark of DowElanco Products Company.

© MONSANTO COMPANY 1994

MAF-3531.01-1/25/94

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

BEST AVAILABLE COPY