

524-351

8/19/2008

1/44



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:
524-351

Date of Issuance:
AUG 19 2008

NOTICE OF PESTICIDE:
 _ Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Name of Pesticide Product:
Land Master BW Herbicide

Name and Address of Registrant (include ZIP Code):
Monsanto Company
1300 "I" St NW, Suite 450 East
Washington, D.C. 20005

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(6)(2)(C) provided you agree in writing to:


1. "PELIGRO" must be added to Page 22 of the label following the word "DANGER."
2. Per the acute toxicity review, text in **bold type** must be added to the First Aid statements currently on the label and the based on toxicity ranking the statements should be placed in the following order:

IF IN EYES:
 Hold eye open and rinse slowly and gently with water for 15-20 minutes.
 Remove contact lenses, if present, after the first 15* minutes, then continue rinsing eye.
 Call a poison control center or doctor for treatment advice.

IF SWALLOWED:
 Call a poison control center or doctor immediately for treatment advice.
 Have person sip a glass of water if able to swallow.
 Do not induce vomiting unless told to by a poison control center or doctor.
Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING:
 Take off contaminated clothing.
 Rinse skin immediately with plenty of water for 15-20 minutes.
 Call a poison control center or doctor for treatment advice.

Signature of Approving Official:
 Joanne I. Miller
 Product Manager 23
 Herbicide Branch
 Registration Division (7505P)



Date:
AUG 19 2008

IF INHALED:

Move person to fresh air.

If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

Call a poison control center or doctor for further treatment advice.

*Note: Per the acute toxicity review, contact lenses may be removed after the first 5 minutes of rinsing instead of 15-20 minutes as specified on the label. The label may be revised to reflect this change.

3. Per the acute toxicity review, the Hazards to Humans and Domestic Animals must be revised to read:

“DANGER

Corrosive. Causes irreversible eye damage. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Do not get in eyes or on clothing. Avoid contact with skin. Avoid breathing spray mist or vapor.”

4. Per the acute toxicity review, the PPE section must be revised to read:

“Some materials that are chemical-resistant to this product are *made of any waterproof material*. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks,
- chemical-resistant gloves,*
- goggles or face shield, and*
- chemical-resistant apron when mixing, loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.”

The text “For applicators applying this product from a tractor, plane or helicopter that has a completely enclosed cab, eye protection is not required, but must be immediately available for use in an emergency, such as a spill or equipment breakdown” appearing on Page 22 of the label is unnecessary and should be deleted from the label.

5. The following engineering control text is missing from Sublabel III and must be added to the label:

“Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40CFR 170.240 (d)(6).”

6. Per the 2,4-D RED label table, the text in **bold type** below must be added to the following User Safety Requirements on Page 22 of the label:

“...If no such instructions for washables **exist**, use detergent and hot water...”

7. The text “Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking, or drinking. Wash hands and arms before using the toilet. After work, remove all clothing and shower using soap and water. If pesticide gets on skin, wash immediately with soap and water” on Page 22 of the label must be replaced with the following User Safety Recommendations text and must be placed in a box:

“User Safety Recommendations

User should wash hands before, eating, drinking, chewing gum, tobacco, of using the toilet.

User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.”

8. Because this product may be used on sod farms, an Agricultural Use Requirements box must be added to Sublabel III and must contain the REI text and early entry PPE as listed in Sublabel I and the 2,4-D RED.

9. A Non-Agricultural Use Requirements box must be added to label with the following entry restriction text:

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to product agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow people (or pets) to enter the treated area until sprays have dried.”

10. Per the product chemistry review, the Agency recommends that the text “Do not store near fertilizers, seeds, insecticides or fungicides” be moved to the “Pesticide Storage” section of the label.

4/44

11. The text below conflicts with the required spray drift text. The label must be revised in Supplemental Labels II and IV:

“Coarse droplets in the 300 to 500 (VMD) micron range are recommended” versus “When applying sprays that contain 2,4-D as the sole active ingredient...apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter or 385 microns or greater for spinning atomizer nozzles” as required in the 2,4-D RED.

“Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety” versus “

Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety” as required in the 2,4-D RED.

“...1) Do not apply within 100 feet of any desirable vegetation or crops. 2) If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops. 3) Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet” versus “... Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas...within 250 feet downwind...” as required in the 2,4-D RED.

12. Reference to chemigation must be deleted from the spray drift section because the label prohibits application through any type of irrigation system.

13. Revisions are needed to the directions for use:

Sublabel I

Farmsteads, Ditches and Fencerows:

-The text “Use a 1 percent solution of this product for spray-to-wet, spot-spraying applications. Spray coverage should be uniform and complete. Do not spray to the point of runoff” should be deleted from the label because it conflicts with rate restrictions specified in the RED for such noncrop sites.

-Per the RED, the following rate restrictions must be added section under the heading “Farmsteads, Ditches and Fencerows:”

“Annual and perennial weeds:

Limited to 2 applications per year.

*Maximum of 2.0 lbs ae per acre per application.

Woody plants:

Limited to 1 application per year.

*Maximum of 4.0 lbs ae per acre per year.”

*The label rate of 108 fluid ounces (1.26 lbs ae) per acre specified on the label may replace the rates above.

Either the ditchbank use must be deleted from the label or the ditch bank use must be moved to a separate section and include the following text as required in the 2,4-D RED:

“Postemergence:

Limited to 2 applications per season.

Maximum of 2.0 lbs ae/acre per application.

Minimum of 30 days between applications.

Spot treatment permitted.

Do not use on small canals with a flow rate less than 10 cubic feet per second (CFS) where water will be used for drinking purposes. CFS may be estimated by using the formula below. The approximate velocity needed for the calculation can be determined by observing the length of time that it takes a floating object to travel a defined distance.

Divide the distance (ft.) by the time (sec.) to estimate velocity (ft. per sec.). Repeat 3 times and use the average to calculate CFS.

$$\text{Average Width (ft.)} \times \text{Average Depth (ft.)} \times \text{Average Velocity (ft. per sec.)} = \text{CFS}$$

For ditchbank weeds:

Do not allow boom spray to be directed onto water surface.

Do not spray across stream to opposite bank.

For shoreline weeds:

Allow no more than 2 foot overspray onto water.”

Sublabels II:

-As noted in item 14 above, the spray drift text appearing in this section of the label must be revised.

-The text “[Insert Specific Use Directions Here]” is unacceptable. Use directions must be listed for the Agency to review.

-The text “For specific rates of application and instructions for control of various annual and perennial weeds, see the Weed Control tables in the label pamphlet for [INSERT BRAND NAME]” appearing on Pages 18 and 19 does not provide enough information to evaluate the directions for use for this sublabel.

Sublabel III:

- The following rate restrictions must be added to the application restriction text currently on the label for ditchbanks appearing on Page 28 of the label:

“Postemergence:

Do not use on small canals with a flow rate less than 10 cubic feet per second (CFS) where water will be used for drinking purposes. CFS may be estimated by using the formula below. The approximate velocity needed for the calculation can be determined by observing the length of time that it takes a floating object to travel a defined distance.

Divide the distance (ft.) by the time (sec.) to estimate velocity (ft. per sec.). Repeat 3 times and use the average to calculate CFS.

Average Width (ft.) x Average Depth (ft.) x Average Velocity (ft. per sec.) = CFS”

-Application restriction text including maximum application rates must be added to the noncrop section of the label (...” airports, parking areas, railroads, rights-of-way, roadsides, utility or industrial sites”) as specified in the 2,4-D RED:

Annual and perennial weeds:

Limited to 2 applications per year.

Maximum of 2.0 lbs ae per acre per application.

Minimum of 30 days between applications.

Woody plants:

Limited to 1 application per year.

Maximum of 4.0 lbs ae per acre per year.”

-The label must clearly specify the following rate restrictions for all ornamental turf:

“Limited to 2 applications per year.

Maximum of 1.5 lbs ae per acre per application.”

-The label must be clearly specify the following rate restrictions for turf grown for seed or sod:

“Limited to 2 applications per year.

Maximum of 2.0 lbs ae per acre per application.

Minimum of 21 days between applications.”

-For stem injections:

Add the following restrictions: “Limit to one injection per year.”

Sublabel IV:

-As noted in item 14 above, the spray drift text appearing in this section of the label must be revised.

-The text “[Insert Specific Use Directions Here]” is unacceptable. Use directions must be listed for the Agency to review.

14. Add the following statements to the labeling: Use of this product in certain of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al.v. EPA, C01-0132C, (W.D. WA). For further information, please refer to EPA Web Site:<http://www.epa.gov/espp>.
15. When Final Printed labels are prepared assure that all front panel text required by 40 CFR Part 156 appears on the front panel including the Signal Word DANGER and Keep Out of the Reach of Children.
16. Delete the phrase “General Use”. This is an implied safety claim.

17. Assure that the required acid equivalents per acre (lbs ae/A) restrictions are expressed as product volume or product weight per unit area that are in the same units as the registered application rate already on the label. Assure that the maximum application restrictions are identical to the highest currently registered rate for each appropriate application site. Finally, ensure that the most restrictive of the required limitations for the active ingredients in this product are placed on the labeling.

Submit one copy of the revised final printed label for the record before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

Enclosure

8/44

I. MAIN LABEL FOR CROP, FALLOW AND REDUCED TILLAGE SYSTEMS

ACCEPTED with COMMENTS in EPA Letter Dated

AUG 19 2008

[INSERT BRAND NAME]

Herbicide

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

524-351

For suppression of emerged field bindweed and control or suppression of other weeds in fallow and reduced tillage systems

Complete Directions for Use

EPA Reg. No. 524-351

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUITS OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.

Read the entire label before using this product.

Use only according to label instructions.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Refillable Container Label Statement: THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A MONSANTO REPACKAGING OR TOLL REPACKAGING AGREEMENT. Non-Refillable Container Label Statement: THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING

INGREDIENTS

ACTIVE INGREDIENTS*:

Table with 2 columns: Ingredient Name and Percentage. Rows include Glyphosate (12.9%), 2,4-D (20.6%), OTHER INGREDIENTS (66.5%), and a total of 100.0%.

*Contains 144 grams per liter or 1.2 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt and 227 grams per liter or 1.9 pounds per U.S. gallon of the active ingredient 2,4-

9/44

D, in the form of its isopropylamine salt. Equivalent to 108 grams per liter or 0.9 pound per U.S. gallon of the acid, glyphosate, and 182 grams per liter or 1.5 pounds per U.S. gallon of the acid, 2,4-D.

No license granted under any non-U.S. patent(s).

IMPORTANT PHONE NUMBERS

1. FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE, 1-800-332-3111.
2. IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL COLLECT, DAY OR NIGHT. (314) 694-4000.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

DANGER! PELIGRO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

CAUSES IRREVERSIBLE EYE DAMAGE.

HARMFUL IF SWALLOWED OR INHALED.

Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to by a poison control center or doctor.
IF IN EYES:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advice.
IF ON SKIN:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> ▪ Move person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. ▪ Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

- Have product container or label with you when calling a poison control center or doctor, or going for treatment.
- You may also contact (314) 694-4000, collect day or night, for emergency medical treatment information.
- This product is identified as {INSERT BRAND NAME}, EPA Registration No. 524-351.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Personal Protective Equipment (PPE):

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, flaggers and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks and protective eyewear. Chemical resistant gloves are required when applying with any handheld nozzle or equipment. A chemical resistant apron plus chemical resistant gloves are required when mixing, loading, cleaning up spills or equipment or when otherwise exposed to this product's concentrate.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. Pilot must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as spill or equipment breakdown.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas..

This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. Application around a cistern or well may result in contamination of drinking water or ground water.

11/44

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas that may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Monsanto Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statement on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

Storage and Disposal

Do not contaminate water, foodstuffs feed or seed by storage or disposal. Do not store near fertilizers, seeds, insecticides or fungicides.

Keep container closed to prevent spills and contamination.

PESTICIDE STORAGE: STORE ABOVE 40°F to keep product in solution. If crystals form, place in a warm room (72°F), allow the product to reach room temperature and roll or shake the container periodically until crystals have dissolved.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

See individual container label for disposal information.

[Alternate Container Label Storage and Disposal Statements by Container Type]:

[FOR PLASTIC ONE-WAY CONTAINERS & BOTTLES]

Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

{FOR ONE-WAY DRUMS}

Do not reuse container. Return container per the Monsanto container return program. If not returned, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

{FOR METAL CONTAINERS (non-aerosol)}

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

{FOR REFILLABLE PORTABLE (MINI-BULK) CONTAINERS}

This container must only be refilled with pesticide product. **Do not reuse this container for any other purpose.**

Final disposal must be in compliance with state and local regulations. If not refilled, returned, or recycled, triple rinse or pressure rinse, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Do not transport this container if it is damaged or leaking. If the container is damaged, leaking or obsolete, or to obtain information about recycling portable refillable containers, contact Monsanto Company at 800-768-6387.

Users: When the container is empty, replace the cap and seal all openings that have been made during usage, and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, triple rinse or pressure rinse the empty container and offer it for recycling if available.

Refillers: Do not reuse this mini-bulk container except for refill in accordance with a valid Monsanto Repackaging or Toll Repackaging Agreement. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting.

{FOR REFILLABLE STATIONARY BULK CONTAINERS}

This container must only be refilled with pesticide product. **Do not reuse this container for any other purpose.**

Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices.

Final disposal must be in compliance with state and local regulations. If not refilled, triple rinse or pressure rinse container and offer for recycling or reconditioning if possible. If burned, stay out of smoke.

GENERAL INFORMATION

This product is a postemergence herbicide for control or suppression of emerged weeds in fallow and reduced tillage systems, prior to planting or emergence of wheat, barley, corn, oats, rye or sorghum (grain or forage), and as a spot treatment in corn, sorghum, forage grasses or forage legumes.

Do not harvest or feed treated vegetation for 8 weeks after application. Following spot treatment in forage grasses or legumes, allow 30 days before harvesting or grazing domestic livestock.

This product enters the plant through the foliage and moves throughout the plant. Visual effects of control are a gradual wilting or yellowing of the plant, which advances to complete browning of above-ground growth and deterioration of affected underground plant parts. Visible symptoms will usually develop within 2 to 4 days after application, but may not occur for 7 or more days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay the visual effects of control.

**APPLICATION PRECAUTIONS
AND RESTRICTIONS**

- Do not plant any crop other than wheat, barley, corn, oats, rye or sorghum (grain or forage) for 3 months after treatment.

- Applications of this product after planting and prior to crop emergence may cause crop injury if rainfall or environmental conditions delaying crop emergence are experienced.
- This product is recommended for control of emerged weeds prior to establishment of labeled crops. Large amounts of green or decaying vegetation left standing or incorporated into the seedbed may enhance the development of disease in newly planted crops. This may result in poor emergence and/or stands, especially under cool and/or wet conditions.
- Spraying early to control young weeds before dense stands develop or light cultivation to assist weed decay will favor preparation of suitable seedbeds.
- In reduced tillage and no-till systems, ensure good seed to soil contact and proper seeding depth.
- Do not cut, feed or forage vegetation from treated areas for 8 weeks following application.
- **DO NOT APPLY IN THE VICINITY OF 2,4-D SENSITIVE CROPS SUCH AS COTTON, GRAPES, TOMATOES AND OTHER DESIRABLE VEGETATION.**
- Applications should be made only when there is no hazard from spray drift, since very small quantities of spray, which may not be visible, may severely injure susceptible crops or desirable vegetation.

Movement of this product on soil particles during windstorms may cause damage to susceptible plants that are contacted. This hazard is reduced if rainfall occurs shortly after application.

- When this label recommends a tank mixture with a generic active ingredient such as diuron, atrazine, or dicamba, the user is responsible for ensuring that the mixture product's label allows the specific application.
- Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this herbicide or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

[INSERT BRAND NAME] is subject to all state and county regulations for 2,4-D amine.

GENERAL USE INSTRUCTIONS

This product should be applied postemergence to vigorously growing weeds when they have reached the recommended size given in the "RECOMMENDED RATES AND WEEDS CONTROLLED" section of this label. Application should be delayed until maximum emergence of the target weeds, but before weeds exceed the maximum size recommended. For annual weeds, allow 1 day after treatment before tillage. For field bindweed, allow at least 7 days after treatment before tillage.

Reduced control may result if treatments are made during poor growing conditions such as drought stress, disease or insect damage or if weeds have been mowed, grazed or cut. Heavy dust on foliage or an overstory canopy covering targeted weeds may also reduce control.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application will wash this product off the foliage and a repeat treatment will be required.

RECOMMENDED RATES AND WEEDS CONTROLLED

For best results, apply this product after most weed seeds have germinated but before seedhead formation in grasses or flower bud formation in broadleaves.

When applied as directed, this product will provide control or suppression of the grass and broadleaf weed species listed below. Rates recommended are for maximum weed height at treatment time.

[INSERT BRAND NAME]

PERENNIAL WEED SPECIES	RATE PER ACRE (FLUID OUNCES)	LENGTH OF VINES/ TREATMENT HEIGHT
------------------------	------------------------------	-----------------------------------

Bindweed, field** <i>Convolvulus arvensis</i>	54 oz (suppression only)	6" to 18"
Spurge, leafy	54 oz (suppression only)	post bloom

ANNUAL WEED SPECIES	RATE PER ACRE	MAXIMUM HEIGHT
------------------------	---------------	----------------

Foxtail, green <i>Setaria viridis</i>	27 oz	12"
---	-------	-----

Barley <i>Hordeum vulgare</i>	40 oz	6"
Brome, downy* <i>Bromus tectorum</i>		
Cheat* <i>Bromus secalinus</i>		
Foxtail <i>Setaria spp.</i>		
Kochia* <i>Kochia scoparia</i>		
Lettuce, prickly* <i>Lactuca serriola</i>		
Oats, wild <i>Avena fatua</i>		
Puncturevine <i>Tribulus terrestris</i>		
Purslane, common <i>Portulaca oleracea</i>		

Cocklebur <i>Xanthium strumarium</i>	40 oz	12"
--	-------	-----

Lambsquarters <i>Chenopodium album</i>		
Mustard, tansy <i>Descurainia pinnata</i>		
Mustard, tumble <i>Sisymbrium altissimum</i>		
Pigweed, redroot <i>Amaranthus retroflexus</i>		
Pigweed, smooth <i>Amaranthus hybridus</i>		
Rye <i>Secale cereale</i>		
Stinkgrass <i>Eragrostis ciliaris</i>		
Thistle, Russian <i>Salsola kali</i>		
Wheat <i>Triticum aestivum</i>		

Barnyardgrass <i>Echinochloa crus-galli</i>	54 oz	6"
---	-------	----

Buffalobur

Solanum rostratum

Goatgrass

Aegilops cylindrical

Mustard, blue

Chorispora tenella

Panicum, fall

Panicum dichotomiflorum

Witchgrass

Panicum capillare

Oats, wild

Avena fatua

54 oz

12"

- * For improved control in no-till systems or heavy infestations or overwintered stands, use 54 fluid ounces. For best results on light kochia infestations, treat after the plant has passed through the woolly stage of growth and is 3 to 6 inches in height. When treating medium to heavy infestations or plants that are in the woolly stage (1 to 3 inches in height), add 2 fluid ounces of Banvel[®] to the recommended rate of this product. Refer to the Banvel label for planting, cropping and other restrictions. Follow all precautions on the Banvel label.
- ** This product may also be used as a 1 percent solution on a spray-to-wet basis for annual weed control and field bindweed suppression. Spray coverage should be uniform and complete. Do not spray to the point of runoff.

Prepare the desired volume of spray by mixing the amount of [INSERT BRAND NAME] in clean water as shown in the following table:

SPRAY SOLUTIONS

DESIRED VOLUME	AMOUNT OF [INSERT BRAND NAME]
1 gallon	1.3 fluid ounces
25 gallons	1 quart
100 gallons	1 gallon

2 tablespoons = 1 fluid ounce

SPOT TREATMENT

Applications in growing crops must be made prior to heading of small grains and grain sorghum (milo) and silking of corn.

For applications in forage grasses and legumes, apply in areas where the movement of domestic livestock can be controlled. No more than one-tenth of any acre should be treated at one time. Do not treat the same area more than once in each growing season. Remove domestic livestock before application and wait 30 days after application before grazing livestock or harvesting.

ECOFARMING SYSTEMS

The Ecofarming System consists of the following rotation: winter wheat, corn/sorghum, ecofallow.

Use the following tank mixtures for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System. Make no more than one preemergence treatment with 2,4-D per season.

16/44

[INSERT BRAND NAME] at 54 to 64 fluid ounces per acre

plus

ATRAZINE at 0.75 to 1 pound active ingredient per acre

The above tank mixture should be applied in 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre. The liquid fertilizer may be diluted with water to achieve the desired carrier volume.

WEEDS CONTROLLED - The following weeds, up to a maximum height of 4 inches, will be controlled:

Brome, downy

Bromus tectorum

Cheat

Bromus secalinus

Foxtail, green

Setaria viridis

Foxtail, yellow

Setaria lutescens

Kochia*

Kochia scoparia

Lettuce, prickly

Lactuca serriola

Pigweed, redroot

Amaranthus retroflexus

Thistle, Russian

Salsola kali

Wheat, volunteer

Triticum aestivum

* For improved control of kochia, add 4 fluid ounces per acre (0.125 pound a.i. per acre) of Banvel to the above tank mixture.

Risk of crop injury from 2,4-D or Banvel can be reduced by applying this treatment 7 to 14 days before planting.

Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products in these mixtures.

PREHARVEST APPLICATIONS TO WHEAT

This product provides weed control when applied prior to harvest of wheat. Make a single application of no more than 42 fluid ounces after the hard-dough stage of grain (30 percent or less grain moisture) and at least 14 days prior to harvest of grain, hay or straw.

This product may be applied using either aerial or ground spray equipment. See the "APPLICATION EQUIPMENT AND TECHNIQUES" section for instructions for ground and aerial applications.

Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.

NOTE: It is not recommended that wheat grown for seed be treated because a reduction in germination or vigor may occur.

POSTHARVEST APPLICATIONS

This product will provide control of weeds following grain harvest. Weeds should be allowed to regrow after damage incurred during harvest operations and to recover from environmental stress before application of this product. Weeds should be treated prior to the heading stage of annual grasses and before broadleaf weeds exceed 24 inches in height. Ammonium sulfate will improve performance on annual weeds under stress conditions.

Weeds controlled with 40 fluid ounces per acre include downy brome, green foxtail, stinkgrass and volunteer wheat.

17/44

Weeds controlled with 54 fluid ounces per acre include field bindweed, kochia, lambsquarters, mustard, pigweed and Russian thistle.

Weeds controlled with 64 fluid ounces per acre include barnyardgrass, sandbur, witchgrass, yellow foxtail and prickly lettuce.

Make up to 2 treatments per year, as needed, observing at least a 30-day retreatment interval. Do not plant any crop other than wheat, barley, corn, oats, rye or sorghum (grain or forage) for 3 months after treatment.

FARMSTEADS, DITCHES AND FENCEROWS

When applied as directed, this product will control downy brome, bulbous bluegrass, kochia, tumble mustard, tansy mustard and prickly lettuce, and provide suppression of crested wheatgrass, smooth bromegrass and field bindweed.

Use a 1 percent solution of this product for spray-to-wet, spot-spraying applications. Spray coverage should be uniform and complete. Do not spray to the point of runoff. Treat when weeds are small, actively growing and free of dust. Use 108 fluid ounces of this product in 5 to 10 gallons of water per acre for broadcast boom applications.

For ditch banks, make no more than 2 applications per season, observing at least a 30-day retreatment interval. Do not allow boom spray to be directed onto water surface. Do not spray across stream to opposite bank. Spot treatments are permitted.

For shoreline weeds, allow no more than 2-foot overspray onto water.

AID TO TILLAGE

This product, used in conjunction with preplant and conventional fallow tillage practices will provide control of downy brome, cheat, volunteer wheat, tansy mustard and foxtail. Apply 27 fluid ounces of this product in 3 to 10 gallons of water per acre to weeds that are actively growing. Treat when weeds are less than 6 inches in height. Application must be followed by conventional tillage practices before regrowth of the treated plant occurs. Allow at least 1 day after application before tillage.

AMMONIUM SULFATE

The addition of 1 to 2 percent dry ammonium sulfate by weight (or liquid equivalent) or 8.5 to 17 pounds per 100 gallons of water may increase the performance of [INSERT BRAND NAME] and [INSERT BRAND NAME] tank mixtures on annual weeds. The improvement in performance may be apparent where environmental stress is a concern. Low-quality ammonium sulfate may contain material that will not readily dissolve which could result in nozzle tip plugging. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet line. ENSURE THAT AMMONIUM SULFATE IS COMPLETELY DISSOLVED IN THE SPRAY TANK BEFORE ADDING HERBICIDES. THOROUGHLY RINSE THE SPRAY SYSTEM WITH CLEAN WATER AFTER USE TO REDUCE CORROSION.

NOTE: Compatibility problems may occur at carrier volumes below 5 gallons per acre.

APPLICATION EQUIPMENT

Do not apply this product through any type of irrigation system.

This product may be applied using ground, aerial spray or hand-held equipment. Use extreme care to avoid misting or drifting of herbicide solution onto foliage, green stems or fruit of desirable crops, trees, or plants during both growing and dormant periods since even very small quantities of spray can cause severe plant injury.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial, airblast, and chemigation) can influence

pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 miles per hour. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 miles per hour, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use, or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables or tobacco. Small amounts of spray drift that may not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

GROUND BROADCAST SPRAY EQUIPMENT: Apply recommended rates of this product in 3 to 10 gallons of water per acre as a broadcast spray. For optimum spray distribution and coverage, use flat-fan or low volume flood nozzles. When using flood nozzles, space them no more than 40 inches apart and ensure double overlap of spray pattern. Refer to the manufacturer's recommendations for correct pressure and nozzle height above the target canopy. Avoid pressure and nozzles that produce fine droplets or mist. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Use appropriate marking devices to ensure uniform spray coverage and best results from [INSERT BRAND NAME] herbicide.

HAND-HELD and HIGH-VOLUME SPRAY EQUIPMENT (use coarse sprays only): Mix this product in clean water and apply to foliage of vegetation to be controlled. For applications made on a spray-to-wet basis, spray coverage should be uniform and complete. Do not spray to the point of runoff. Use a 1 percent solution for annual weed control and field bindweed suppression.

AERIAL EQUIPMENT: Apply the recommended rates of this product in 3 to 5 gallons of water per acre as a broadcast spray. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

The boom length must not exceed 75 percent of the wingspan or 90 percent of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications. When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

MIXING INSTRUCTIONS

[INSERT BRAND NAME] ALONE

Fill the spray tank to about 3/4 of the desired volume with clean water. Add the recommended amount of this product, then complete the filling process while maintaining agitation. Remove the hose from the mix tank immediately after filling to avoid siphoning back into the carrier source. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, terminate by-pass and return lines at the tank bottom and/or use an agriculturally approved anti-foam or defoaming agent.

Additional surfactant is not necessary for this formulation.

NOTE: Reduced control may occur if water containing soil is used, such as water from ponds and unlined ditches.

TANK MIXTURES

Always predetermine the compatibility of labeled tank mixtures of this herbicide with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of [INSERT BRAND NAME] with water as follows:

1. Place a 20- to 35-mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with water and start agitation.
3. If a wettable powder is used, make a slurry with the water carrier, and add it SLOWLY through the screen into the tank. Continue agitation.
4. If a flowable formulation is used, premix one part flowable with one part water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
5. If an emulsifiable concentrate formulation is used, premix one part emulsifiable concentrate with two parts water. Add diluted mixture SLOWLY through the screen into the tank. Continue agitation.
6. Continue filling the sprayer tank with water and add the required amount of this product near the end of the filling process.
7. Add individual formulations to the spray tank as follows: wettable powder, flowable, emulsifiable concentrate, drift control additive, water soluble liquid.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle should be no finer than 100-mesh and in-line strainers should be no finer than 50-mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat-fan nozzles.

Check the label of all products used in the tank mix for cleaning instructions. Clean as per the hardest material to remove.

SPRAYER CLEANUP

CLEAN THE ENTIRE SPRAYER AFTER APPLICATION OF THIS PRODUCT. Failure to clean the sprayer thoroughly may result in injury to desirable crops that are subsequently sprayed.

First, add clean water to the tank and thoroughly rinse the entire sprayer system. Secondly, fill the tank with water and ammonia. Add 1 quart of household ammonia per 25 gallons of water. Pump enough

solution through the hoses, boom and nozzles to fill these parts completely. Then fill the tank, close and leave for 24 hours before draining and rinsing thoroughly with water.

Application or use of other agricultural chemicals with the equipment used for this product may result in injury to desirable vegetation.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE. The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this Limit of Warranty and Liability which may not be varied by any verbal or written agreement.

[INSERT BRAND NAME] is a registered trademarks of Monsanto Technology LLC.

All other trademarks are the property of their respective owners.

© [DATE] MONSANTO COMPANY

ST. LOUIS, MISSOURI 63167 USA

II. SUPPLEMENTAL LABELING FOR CROP, FALLOW AND REDUCED TILLAGE SYSTEMS
--

Table of Contents: Fallow and Reduced Tillage Systems Supplemental Labeling

	Name	Approval Date
A	FOR AERIAL APPLICATION IN ARKANSAS ONLY	December 12, 2003
B	FOR USE WITH HOODED SPRAYERS IN CORN	June 19, 1996
C	FOR USE WITH HOODED SPRAYERS IN MILO	June 19, 1996

22/44

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR [INSERT BRAND NAME] HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using [INSERT BRAND NAME] as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

[INSERT BRAND NAME]

Herbicide

EPA Reg. No. 524-351

Keep out of reach of children.

DANGER! PELIGRO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

[INSERT BRAND NAME] is a registered trademark of Monsanto Technology LLC.

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See the "GENERAL INFORMATION" and "MIXING INSTRUCTIONS" sections of the label booklet for [INSERT BRAND NAME] for essential product performance information.

[INSERT SPECIFIC USE DIRECTIONS HERE]

Read the "Limit of Warranty and Liability" in the label booklet for [INSERT BRAND NAME] before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© [DATE] MONSANTO COMPANY

ST. LOUIS, MISSOURI 63167

A. FOR AERIAL APPLICATION IN ARKANSAS ONLY

USE DIRECTIONS

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the recommended rate of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75 percent of the length of the wingspan or rotor. In many cases, reducing this distance to 65 percent of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing airflow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge.

Do not apply this product when winds are in excess of 10 miles per hour.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 miles per hour.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

Read and carefully observe the label claims, cautionary statements, and all information on the label for **[INSERT BRAND NAME]**.

24/44

B. FOR USE WITH HOODED SPRAYES IN CORN

USE INSTRUCTIONS

This product may be used through hooded sprayers for weed control between the rows of corn. Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to corn that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 42 fluid ounces of this product per acre per application.
- Corn must be at least 12 inches tall, measured without extending leaves.
- Do not apply after tasseling.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 miles per hour.
- Maximum wind speed: 10 miles per hour.
- Use low-drift nozzles.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

For specific rates of application and instructions for control of various annual and perennial weeds, see the Weed Control tables in the label pamphlet for **[INSERT BRAND NAME]**.

Do not graze or feed corn forage or fodder following applications of this product through hooded sprayers.

Do not make more than one application per growing season.

C. FOR USE WITH HOODED SPRAYES IN MILO

USE INSTRUCTIONS

This product may be used through hooded sprayers for weed control between the rows of grain sorghum (milo). Only hooded sprayers that completely enclose the spray pattern may be used.

A hooded sprayer is a type of shielded applicator. The spray pattern is completely enclosed on the top and all 4 sides by a hood, thereby shielding the crop from the spray solution. This equipment must be set up and operated in a manner that avoids bouncing or raising the hoods off the ground in any way. If the hoods are raised, spray particles may escape and come into contact with the crop, causing damage or destruction of the crop. The spray hoods must be operated on the ground or skimming across the ground. Tractor speed must be adjusted to avoid bouncing of the spray hoods. Avoid operation on rough or sloping ground where the spray hoods might be raised off the ground.

When applying to milo that is grown on raised beds, ensure that the hood is designed to completely enclose the spray solution. If necessary, extend the front and rear flaps of the hoods to reach the ground in deep furrows.

Follow these requirements:

- The spray hoods must be operated on the ground or skimming across the ground.
- Do not apply more than 54 ounces of this product per acre per application.
- Milo must be at least 12 inches tall, measured without extending leaves. Treat before milo sends tillers between the drill rows. If such tillers are contacted with the spray solution, the main plant may be killed.
- Do not apply to milo during boot, flowering or early dough stages.
- Leave at least an 8 inch untreated strip over the drill row. For example, if the crop row width is 38 inches, the maximum width of the spray hood should be 30 inches.
- Maximum tractor speed: 5 miles per hour.
- Maximum wind speed: 10 miles per hour.
- Use low-drift nozzles.

Crop injury may occur when the foliage of treated weeds comes into direct contact with leaves of the crop. Do not apply this product when the leaves of the crop are growing in direct contact with weeds to be treated. Droplets, mist, foam or splatter of the herbicide solution may contact the crop and cause discoloration, stunting or destruction.

Contact of this product in any manner to any vegetation to which treatment is not intended may cause damage. Such damage shall be the sole responsibility of the applicator.

For specific rates of application and instructions for control of various annual and perennial weeds, see the Weed Control tables in the label booklet for **[INSERT BRAND NAME]**.

Do not graze or feed milo forage or fodder following applications of this product through hooded sprayers.

Do not make more than one application per growing season.

26/44

III. MAIN LABEL FOR NON-CROP, RANGELAND, AND INDUSTRIAL USES

Note : Part III. "NON-CROP, RANGELAND, AND INDUSTRIAL USES" of this Master Label consists only of approved use-sites that are outside the scope of the Worker Protection Standard (WPS), 40 CFR part 170. No WPS Agricultural Use Requirements or Non-Agricultural Use Requirements boxes are required for these uses, and no WPS boxes will be included on any final printed labeling for products under this registration that are limited solely to these Part III. use-sites.

[INSERT BRAND NAME]

Herbicide by Monsanto

A broad-spectrum postemergence herbicide for non-crop, rangeland and industrial weed control.

Complete Directions for Use

EPA Reg. No. 524-351

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUITS OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION IS LIKELY TO RESULT.

Read the entire label before using this product.

Use only according to label instructions.

Read the "LIMIT OF WARRANTY AND LIABILITY" statement at the end of the label before buying or using. If terms are not acceptable, return at once unopened.

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. SEE INDIVIDUAL CONTAINER LABEL FOR REPACKAGING LIMITATIONS.

Refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION. IT IS INTENDED THAT REPACKAGING BE ONLY IN ACCORDANCE WITH A MONSANTO REPACKAGING OR TOLL REPACKAGING AGREEMENT.

Non-Refillable Container Label Statement:

THIS IS AN END-USE PRODUCT. MONSANTO DOES NOT INTEND AND HAS NOT REGISTERED IT FOR REFORMULATION OR REPACKAGING

CONTENTS

- 1 1.0 **INGREDIENTS**
- 2 2.0 **IMPORTANT PHONE NUMBERS**
- 3 3.0 **PRECAUTIONARY STATEMENTS**

- 3.1 Hazards to Humans and Domestic Animals
- 3.2 Environmental Hazards
- 3.3 Physical or Chemical Hazards
- 4 4.0 **STORAGE AND DISPOSAL**
- 5 5.0 **GENERAL INFORMATION**
- 6 6.0 **MIXING**
 - 6.1 Mixing with Water
 - 6.2 Tank Mixing Procedure
- 7 7.0 **APPLICATION EQUIPMENT AND TECHNIQUES**
 - 7.1 Aerial Application Equipment
 - 7.2 Ground Broadcast Equipment
 - 7.3 Equipment Cleaning
- 8 8.0 **SITE AND USE RECOMMENDATIONS**
 - 8.1 Bermudagrass
 - 8.2 Bahiagrass
 - 8.3 Tall Fescue
 - 8.4 Rangelands
 - 8.5 Hollow Stem Injection
- 9 9.0 **WEEDS CONTROLLED**
- 10 10.0 **LIMIT OF WARRANTY AND LIABILITY**

1.0 INGREDIENTS

ACTIVE INGREDIENTS*:

Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt	12.9%
2,4-D, 2,4-dichlorophenoxyacetic acid, in the form of its isopropylamine salt	20.6%
OTHER INGREDIENTS.....	<u>66.5%</u>
	100.0%

28/44

*Contains 144 grams per liter or 1.2 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt and 227 grams per liter or 1.9 pounds per U.S. gallon of the active ingredient 2,4-D, in the form of its isopropylamine salt. Equivalent to 108 grams per liter or 0.9 pound per U.S. gallon of the acid, glyphosate, and 182 grams per liter or 1.5 pounds per U.S. gallon of the acid, 2,4-D.

No license granted under any non-U.S. patent(s).

2.0 IMPORTANT PHONE NUMBERS

1. FOR PRODUCT INFORMATION OR ASSISTANCE IN USING THIS PRODUCT, CALL TOLL-FREE, 1-800-332-3111.
2. IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT, OR FOR MEDICAL ASSISTANCE, CALL COLLECT, DAY OR NIGHT, (314) 694-4000

3.0 PRECAUTIONARY STATEMENTS

3.1 Hazards to Humans and Domestic Animals

Keep out of reach of children.

DANGER!

CAUSES IRREVERSIBLE EYE DAMAGE.

HARMFUL IF SWALLOWED OR INHALED.

Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

All mixers, loaders, applicators, flaggers and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks and protective eyewear. Chemical resistant gloves are required when applying with any handheld nozzle or equipment. A chemical resistant apron plus chemical resistant gloves are required when mixing, loading, cleaning up spills or equipment or when otherwise exposed to this product's concentrate. For applicators applying this product from a tractor, plane or helicopter that has a completely enclosed cab, eye protection is not required, but must be immediately available for use in an emergency, such as a spill or equipment breakdown..

Wash hands, face and arms with soap and water as soon as possible after mixing, loading, or applying this product. Wash hands, face and arms with soap and water before eating, smoking, or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. If pesticide gets on skin, wash immediately with soap and water.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining protective equipment. If no such instructions are available for washables, use detergent and hot water. Keep and wash protective equipment and clothing separately from other laundry. After each day of use, clothing or protective equipment must not be reused until it has been cleaned.

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to by a poison control center or doctor.
IF IN EYES:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 15 minutes, then continue rinsing eye.

	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor for treatment advice.
IF ON SKIN:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> ▪ Move person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. ▪ Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<ul style="list-style-type: none"> • Have product container or label with you when calling a poison control center or doctor, or going for treatment. • You may also contact (314) 694-4000, collect day or night, for emergency medical treatment information. • This product is identified as [INSERT BRAND NAME], EPA Registration No. 524-351. 	

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

3.2 Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

Drift or runoff may be hazardous to aquatic organisms in water adjacent to treated areas.

This chemical has properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination. Application around a cistern or well may result in contamination of drinking water or ground water.

3.3 Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas that may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This product can only be used in accordance with the Directions for Use on this label or in separately published Monsanto Supplemental Labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations. Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

4.0 STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Do not store near fertilizers, seeds, insecticides or fungicides.

PESTICIDE STORAGE: STORE ABOVE 40°F to keep product in solution. If crystals form, place in a warm room (72°F), allow the product to reach room temperature and roll or shake the container periodically until crystals have dissolved.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

See individual container label for disposal information.

[Alternate Container Label Storage and Disposal Statements by Container Type]:

[FOR PLASTIC ONE-WAY CONTAINERS & BOTTLES]

Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[FOR ONE-WAY DRUMS]

Do not reuse container. Return container per the Monsanto container return program. If not returned, triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[FOR METAL CONTAINERS (non-aerosol)]

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

[FOR REFILLABLE PORTABLE (MINI-BULK) CONTAINERS]

This container must only be refilled with pesticide product. **Do not reuse this container for any other purpose.**

Final disposal must be in compliance with state and local regulations. If not refilled, returned, or recycled, triple rinse or pressure rinse, puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Do not transport this container if it is damaged or leaking. If the container is damaged, leaking or obsolete, or to obtain information about recycling portable refillable containers, contact Monsanto Company at 800-768-6387.

Users: When the container is empty, replace the cap and seal all openings that have been made during usage, and return the container to the point of purchase, or to an alternate location designated by the manufacturer at the time of purchase of this product. If not returned, triple rinse or pressure rinse the empty container and offer it for recycling if available.

Refillers: Do not reuse this mini-bulk container except for refill in accordance with a valid Monsanto Repackaging or Toll Repackaging Agreement. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting.

[FOR REFILLABLE STATIONARY BULK CONTAINERS]

This container must only be refilled with pesticide product. **Do not reuse this container for any other purpose.**

Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices.

Final disposal must be in compliance with state and local regulations. If not refilled, triple rinse or pressure rinse container and offer for recycling or reconditioning if possible. If burned, stay out of smoke.

5.0 GENERAL INFORMATION

Product Description:

This product is a broad-spectrum postemergence herbicide for control or suppression of emerged weeds growing in non-crop, rangeland, and industrial areas.

Time to Symptoms: This product enters the plant through the foliage and moves throughout the plant. Visual effects of control are a gradual wilting or yellowing of the plant, which advances to complete browning of aboveground growth and deterioration of affected underground plant parts. Visible symptoms will usually develop on labeled weeds within 2 to 4 days after application, but may not occur for 7 or more days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay the visual effects of control.

Stage and Condition of Weeds: This product should be applied postemergence to vigorously growing weeds at a rate recommended in the "RECOMMENDED RATES AND WEEDS CONTROLLED" section of this label. Application should be delayed until maximum emergence of the target weeds but before mature seeds are produced.

Reduced control may result if treatments are made during poor growing conditions such as drought stress, disease or insect damage or if weeds have been mowed, grazed or cut. Heavy dust on foliage or an overstory canopy covering targeted weeds may also reduce control.

Rainfastness: Rainfall occurring within 6 hours after application may reduce effectiveness.

Tank Mixing: When tank mixing with [INSERT BRAND NAME] follow a label approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used. Use according to the most restrictive label directions for each product in the mixture.

When this label recommends a tank mixture with a generic active ingredient such as diuron, atrazine, or dicamba, the user is responsible for ensuring that the mixture product's label allows the specific application.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this label. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance.

PRECAUTIONS AND RESTRICTIONS

DO NOT APPLY IN THE VICINITY OF 2,4-D SENSITIVE CROPS SUCH AS COTTON, GRAPES, TOMATOES AND OTHER DESIRABLE VEGETATION.

Applications should be made only when there is no hazard from spray drift since very small quantities of spray, which may not be visible, may severely injure susceptible crops or desirable vegetation.

The likelihood of injury occurring to adjacent crops from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions including lesser wind velocities will favor spray drift.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this herbicide or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

[INSERT BRAND NAME] is subject to all state and county regulations for 2,4-D amine.

6.0 MIXING

6.1 Mixing with Water

Fill the spray tank to about 3/4 of the desired volume with clean water. Add the recommended amount of this product, then complete the filling process while maintaining agitation. Remove the hose from the mixing tank immediately after filling to avoid siphoning back into the carrier source.

During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, terminate by-pass and return lines at the tank bottom and/or use an agriculturally approved anti-foam or defoaming agent.

Additional surfactant is not necessary for this product.

NOTE: Reduced results may occur if water, containing soil is used, such as water from ponds and unlined ditches.

6.2 Tank Mixing Procedure

[INSERT BRAND NAME] Tank Mixtures With Oust® XP, dicamba, Escort® XP, Tordon™ and Transline™

Refer to the "Tank Mixing" section of "GENERAL INFORMATION" for additional precautions.

Always predetermine the compatibility of labeled tank mixtures of this herbicide with water carrier by mixing small proportional quantities in advance.

Mix labeled tank mixtures of [INSERT BRAND NAME] with water as follows:

1. Place a 20- to 35-mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with water and start agitation.
3. Add the required amount of Oust XP, dicamba, Escort XP, Tordon or Transline slowly while maintaining agitation.
4. Continue filling the sprayer tank with water and add the required amount of this product near the end of the filling process.

Maintain good agitation at all times until the contents of the tank are sprayed. If the spray mixture is allowed to settle, thorough agitation is required to resuspend the mixture before spraying is resumed. Keep by-pass line on or near bottom of tank to minimize foaming.

7.0 APPLICATION EQUIPMENT AND TECHNIQUES

This product may be applied using ground or aerial spray equipment. Use extreme care to avoid misting or drifting of herbicide solution onto foliage, green stems or fruit of desirable crops, trees, or plants during both growing and dormant periods since even very small quantities of spray can cause severe plant injury.

SPRAY DRIFT MANAGEMENT

AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended.

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, aerial, airblast, and chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 15 miles per hour. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 miles per hour, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use, or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables or tobacco. Small amounts of spray drift that may not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

7.1 Aerial Equipment

DO NOT APPLY THIS PRODUCT USING AERIAL SPRAY EQUIPMENT EXCEPT UNDER CONDITIONS AS SPECIFIED WITHIN THIS LABEL.

Use the recommended rates of this product in 3 to 5 gallons of water per acre as a broadcast spray. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Avoid direct application to any body of water.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

Ensure uniform application --To avoid streaked, uneven or overlapped application, use appropriate marking devices.

The boom length must not exceed 75 percent of the wingspan or 90 percent of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Pilot must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].

Aircraft Maintenance

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR ARE MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

7.2 Ground Broadcast Equipment

Apply recommended rates of this product in 15 to 30 gallons of water per acre as a broadcast spray. For optimum spray distribution and coverage, use flat fan or low-volume flood nozzles. When using flood nozzles, space them no more than 40 inches apart and ensure double overlap of spray pattern. Refer to the manufacturer's recommendations for correct pressure and nozzle height above the target canopy. Avoid pressure settings and nozzles that produce fine droplets or mist. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

Use appropriate marking devices to ensure uniform spray coverage and best results from this product.

7.3 Equipment Cleaning

CLEAN THE ENTIRE SPRAYER AFTER APPLICATION OF THIS PRODUCT. Add clean water to the tank and thoroughly rinse the entire sprayer system, then fill the tank with water and ammonia. Add 1 quart of household ammonia per 25 gallons of water. Pump enough solution through the hoses, boom and nozzles to fill these parts completely. Then fill the tank, close and leave for 24 hours before draining and rinsing thoroughly with water.

8.0 SITE AND USE RECOMMENDATIONS

[INSERT BRAND NAME] may be used for general postemergence control or suppression of emerged weeds. This product is recommended for use on weeds growing in areas such as airports, apartment complexes, cemeteries, commercial sites, Conservation Reserve Program (CRP) areas, ditch banks, dry ditches, dry canals, fencerows, golf courses, highway rights-of-way, industrial sites specified on this label, landscape areas, lumberyards, manufacturing sites, municipal sites, natural areas, office complexes, parking areas, pastures, petroleum tank farms and pumping installations, railroads, rangeland, recreational areas, rights-of-way, roadsides, sod or turf seed farms, sports complexes, storage areas, substations, utility sites, warehouse areas, and wildlife management areas.

Precautions, Restrictions:

- For ditch banks: Make no more than 2 applications per season, observing at least a 30-day retreatment interval. Each application is limited to no more than 4 pints. Do not allow boom spray to be directed onto water surface. Do not spray across stream to opposite bank. Spot treatments are permitted.
- For shoreline weeds: allow no more than 2 foot overspray onto water.
- For airports, parking areas, railroads, rights-of-ways, roadsides, utility or industrial sites specified on this label: make only one application per year for the control of woody plants and no more than two applications, observing at least a 30-day interval, for control on annual or perennial weeds.

8.1 Bermudagrass

When treating bermudagrass, make no more than 2 applications per year and observe at least a 21-day retreatment interval.

Dormant Bermudagrass

Fine Turf (Highly Maintained Turf) When applied as directed, this product will provide control or suppression of vines and many winter annual grasses and broadleaf weeds for effective release of dormant bermudagrass in highly maintained turf. Treat only when turf is dormant and prior to spring greenup. For best results, treat winter annuals when plants are in an early growth stage but after most have germinated. Apply 2 to 4 pints of this product in 15 to 30 gallons of spray solution per acre. **DO NOT** apply tank mixtures of this product plus Outrider or Oust XP in highly maintained turfgrass areas. See the "WEEDS CONTROLLED" section for the correct rate selection.

Coarse Turf (Low Maintenance Turf) When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines for release of actively growing bermudagrass.

Apply 2 to 4 pints of this product in 15 to 30 gallons of spray solution per acre. A tank mixture of this product plus Outrider, Oust XP or Escort XP may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.75 to 2.0 ounces of Outrider or 0.25 to 1 ounce of Oust XP or Escort XP per acre. See the "WEEDS CONTROLLED" section for the correct rate selection.

For control of annual ryegrass (*Lolium multiflorum*) in dormant coarse bermudagrass turf, a tank mixture with Roundup PRO Concentrate herbicide or Roundup PRO herbicide may be used. See rate chart below for proper rate selection.

	[INSERT BRAND NAME] Rate (pints/acre)			
	1.5	2.0	3.0	4.0
	Rate of Roundup brand product to add (ounces/acre)			
Roundup PRO Concentrate	18	16	8	4
Roundup PRO	21	19	9	5

Actively Growing Bermudagrass

For spring and summer applications to bermudagrass, apply 2 to 4 pints of this product in 15 to 30 gallons of spray solution per acre. A tank mixture of this product plus Outrider, Oust XP or Escort XP may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.75 to 2.0 ounces of Outrider or 0.25 to 1 ounce of Oust XP or Escort XP per acre. Use only in areas where bermudagrass is a desirable groundcover and where some temporary discoloration can be tolerated. See the "WEEDS CONTROLLED" section for the correct rate selection.

8.2 Bahiagrass

When treating bahiagrass, make no more than 2 applications per year and observe at least a 21-day retreatment interval.

Dormant Bahiagrass

When applied as directed, this product will provide control or suppression of vines and winter annual weeds for release of dormant bahiagrass. For best results on winter annuals, treat when plants are in an early growth stage but after most plants have germinated. To avoid delays in greenup and to minimize injury, treat when bahiagrass is dormant.

Apply 2 to 4 pints of this product in 15 to 30 gallons of spray solution per acre. A tank mixture of this product plus Outrider or Oust XP may be used to provide control or suppression of certain weeds. For this tank mixture, apply 2 to 4 pints of this product plus 0.75 to 2.0 ounces of Outrider or 0.25 to 0.5 ounce of Oust XP per acre. See the "WEEDS CONTROLLED" section for the correct rate selection.

Actively Growing Bahiagrass

When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines for release of bahiagrass.

Apply 1.5 to 2 pints of this product in 15 to 30 gallons of spray solution per acre. A tank mixture of this product plus Outrider or Oust XP may be used to provide control or suppression of certain weeds. For this tank mixture, apply 1.5 to 2 pints of this product plus 0.75 to 2.0 ounces of Outrider or 0.25 ounce of Oust XP per acre. See the "WEEDS CONTROLLED" section for the correct rate selection.

8.3 Tall Fescue

When treating tall fescue, make no more than 2 applications per year and observe at least a 21-day retreatment interval.

Spring Applications. When applied as directed, this product will provide control or suppression of vines and many winter annual weeds in tall fescue. For best results on winter annuals, treat when plants are in an early growth stage but after most have germinated. To minimize tall fescue injury, apply this product when tall fescue is 4 to 6 inches tall but prior to seedhead emergence.

Apply 2 to 3 pints of this product in 15 to 30 gallons of spray solution per acre. A tank mixture of this product plus Oust XP may be used to provide control of certain weeds. For this tank mixture, apply 2 pints of this product plus 0.25 ounce of Oust XP per acre. See the "WEEDS CONTROLLED" section for the correct rate selection.

Summer Applications. When applied as directed, this product will provide control or suppression of several grasses, broadleaf weeds, and vines in tall fescue.

Apply 2 to 3 pints of this product in 15 to 30 gallons of spray solution per acre. A tank mixture of this product plus Oust XP may be used to provide control of certain weeds. For this tank mixture, apply 2 to 3 pints of this product plus 0.25 to 0.5 ounce of Oust XP per acre. See the "WEEDS CONTROLLED" section for the correct selection.

8.4 RANGELANDS

Preventing viable seed production is key to the successful control and invasion of annual grassy weeds in rangelands. Follow-up applications in sequential years should eliminate most of the viable seeds.

Grazing of treated areas should be delayed to encourage growth of desirable perennials. Allowing desirable perennials to flower and reseed in the treated area will encourage successful transition.

PRECAUTIONS, RESTRICTIONS: Do not use ammonium sulfate when spraying rangeland grasses with this product. Do not make more than one application per year. Do not cut grass or forage for hay.

Postemergence Applications to Control Grass Weeds

When applied as directed, this product will control or suppress many annual grass weeds growing in perennial cool and warm season grass rangelands.

Apply 40 to 54 fluid ounces of this product to control or suppress many weeds, including downy brome, cheat grass, cereal rye and jointed goatgrass in rangelands. Apply when most mature brome plants are in early flower and before the plants including seedheads turn color. Allowing for secondary weed flushes to occur in the spring following rain events further depletes the seed reserve, and encourages perennial grass conversion on weedy sites. Fall applications are possible, and recommended where spring moisture is usually limited and fall germination allows for good weed growth.

For medusahead control, apply 54 fluid ounces when the medusahead has reached the 3-leaf stage. Delaying applications beyond this stage will result in reduced or unacceptable control. Fire may be useful in eliminating the thatch layer produced by slow decaying culms prior to application. Allow new growth to occur before spraying after a burn. Repeat applications in subsequent years may be necessary to eliminate the seedbank before reestablishing desirable perennial grasses in medusahead-dominated rangelands.

Slight discoloration of the desirable grasses may occur, but they will regreen and regrow under moist soil conditions as effects of this product wear off.

Postemergence Applications to Control Noxious Weeds

When applied as directed, this product will provide control or suppression of noxious weeds growing in rangelands, perennial grasslands, roadsides, or similar industrial sites specified on this label. For best results on biennial or perennial noxious weeds, treat early season when target plants are in the rosette stage of growth, except as described below. See specific weed control recommendations listed below.

Apply 40 to 54 fluid ounces of this product per acre at recommended water volumes.

Leafy Spurge. Apply 40 to 54 fluid ounces of this product alone, or in mixtures with 4 to 6 fluid ounces of Banvel herbicide, or 4 to 8 fluid ounces of Tordon herbicide at early flower bud stage. This application works best in areas of greater than 20-inch average annual rainfall or when local conditions provide adequate soil moisture to promote vigorous growth.

Perennial Pepperweed. Apply 40 to 54 fluid ounces of this product per acre alone or in combination with 0.5 ounce per acre of Escort from early bud stage through full flower.

Yellowstar Thistle, Spotted Knapweed, Diffuse Knapweed, Musk Thistle, and Common Teasel. Apply 40 to 54 fluid ounces of this product alone or in combination with 0.25 to 0.5 pint Transline herbicide, or 4 to 8 fluid ounces of Tordon herbicide per acre at the rosette stage through early bolting.

Note: When applied over desirable perennial grasses, these mixtures may result in temporary grass leaf discoloration. Therefore, where desirable perennial grasses are present and slight and temporary discoloration cannot be tolerated, it is recommended that a rate of 40 fluid ounces per acre not be exceeded.

8.5 Hollow Stem Injection

This product may be applied through hand-held injection devices that deliver recommended amounts of this product into targeted hollow-stem plants growing in any non-crop, rangeland, or industrial sites specified on this label. For control of the weeds listed below, inject 6 mL/stem of this product between the second and third internodes of the stem.

Japanese Knotweed, *Polygonum cuspidatum*.

Bohemian Knotweed, *Polygonum bohemicum*.

Note: Based on the maximum annual use rate of 2,4-D for these non-crop sites, the combined total for all treatments must not exceed 10.8 quarts per acre of this product. At 6 mL/stem, 10.8 quarts should treat about 1700 stems per acre.

9.0 WEEDS CONTROLLED

Rate recommendations for specific weed species in non-crop and industrial sites are given below.

Use the lower rate when treating annual weeds below 6 inches in height. Use the higher rate on weeds taller than 6 inches or as they approach flower or seedhead formation. Use the higher rate of this product for control or partial control of perennial species. Use the lower rates for growth suppression of perennials.

For the best rate recommendation on the mixture of weeds within your geographic area, contact your local Monsanto representative.

**WEEDS CONTROLLED OR SUPPRESSED
[INSERT BRAND NAME] ALONE**

Note: C = Control
S = Suppression

ANNUAL WEED SPECIES	[INSERT BRAND NAME] pints/acre			
	1.5	2	3	4
Barley, little <i>Hordeum pusillum</i>	S	S	C	C
Barnyardgrass (< 6 inches in height) <i>Echinochloa Crus-galli</i>	S	S	S	C
Bedstraw, catchweed <i>Galium aparine</i>	S	S	C	C
Bluegrass, annual <i>Poa annua</i>	S	S	C	C
Brome, downy <i>Bromus tectorum</i>	S	S	C	C

Brome, Japanese <i>Bromus, japonicus</i>	S	S	C	C
Cheat <i>Bromus secalinus</i>	S	S	C	C
Chervil <i>Chaerophyllum tainturieri</i>	S	S	C	C
Chickweed, common <i>Stellaria media</i>	S	S	C	C
Clover, crimson <i>Trifolium incarnatum</i>	.	S	S	C
Clover, hop <i>Trifolium spp.</i>	.	S	C	C
Crabgrass <i>Digitaria spp.</i>	.	S	C	C
Foxtail <i>Setaria spp.</i>	.	S	C	C
Geranium, Carolina <i>Geranium carolinianum</i>	S	C	C	C
Henbit <i>Lamium amplexicaule</i>	S	C	C	C
Horseweed/Marestail <i>Conyza canadensis</i>	S	C	C	C
Oats, wild <i>Avena fatua</i>	S	S	C	C
Panicum, fall (< 6 inches in height) <i>Panicum dichotomiflorum</i>	.	S	S	C
Partridgepea <i>Cassia fasciculata</i>	.	S	C	C
Purple Deadnettle <i>Lamium purpureum</i>	S	C	C	C
Ragweed, common <i>Ambrosia artemisiifolia</i>	.	S	C	C
Sowthistle, annual (rosette stage only) <i>Sonchus oleraceus</i>	.	S	S	C
Speedwell, corn <i>Veronica arvensis</i>	S	S	C	C
Spurge, spotted <i>Euphorbia maculata</i>	.	S	C	C
Vervain, blue <i>Verbena hastata</i>	.	S	C	C
Vetch <i>Vicia spp.</i>	.	S	C	C

BIENNIAL WEED SPECIES

Carrot, wild <i>Daucus carota</i>	.	.	S	C
Thistle, musk (rosette stage only) <i>Carduus nutans</i>	.	S	S	C

PERENNIAL WEED SPECIES

Bahiagrass <i>Paspalum notatum</i>	S	S	S	S
Bindweed, field <i>Convolvulus arvensis</i>	.	.	.	S

Dock, curly <i>Rumex crispus</i>	•	S	C	C
Fescue, tall <i>Festuca arundinacea</i>	S	S	S	S
Greenbriar <i>Smilax spp.</i>	•	•	•	S
Honeysuckle <i>Lonicera spp.</i>	•	S	S	C
Horsenettle <i>Solanum carolinense</i>	•	•	S	S
Johnsongrass <i>Sorghum halepense</i>	•	S	S	C
Peppervine <i>Ampelopsis arborea</i>	•	S	C	C
Plantain, buckhorn <i>Plantago lanceolata</i>	S	S	C	C
Raspberry <i>Rubus spp.</i>	•	•	•	S
Ryegrass <i>Lolium spp.</i>	•	•	•	S
Trumpet creeper <i>Campsis radicans</i>	•	•	•	S
Vaseygrass <i>Paspalum urvillei</i>	•	S	C	C

**WEEDS CONTROLLED OR SUPPRESSED
[INSERT BRAND NAME] plus OUST**

Note: C= Control S = Suppression

ANNUAL WEED SPECIES	[INSERT BRAND NAME] (PINTS/ACRE)	1.5	2	3	3	4	4	4
	OUST XP (OZ/ACRE)	+	+	+	+	+	+	+
		1/4	1/4	1/4	1/2	1/4	1/2	1
Barley, little <i>Hordeum pusillum</i>		C	C	C	C	C	C	C
Bedstraw, catchweed <i>Galium aparine</i>		C	C	C	C	C	C	C
Bluegrass, annual <i>Poa annua</i>		S	S	C	C	C	C	C
Chervil <i>Chaerophyllum tainturieri</i>		C	C	C	C	C	C	C
Chickweed, common <i>Stellaria media</i>		S	S	C	C	C	C	C
Clover, crimson <i>Trifolium incarnatum</i>		S	S	S	C	C	C	C
Clover, hop <i>Trifolium spp.</i>		•	S	C	C	C	C	C
Crabgrass <i>Digitaria spp.</i>		•	S	C	C	C	C	C
Foxtail <i>Setaria spp.</i>		•	S	C	C	C	C	C
Geranium, Carolina <i>Geranium carolinianum</i>		S	C	C	C	C	C	C
Henbit <i>Lamium amplexicaule</i>		S	C	C	C	C	C	C
Partridgepea <i>Cassia fasciculata</i>		•	S	C	C	C	C	C
Ragweed, common		•	S	C	C	C	C	C

<i>Ambrosia artemisiifolia</i>							
Speedwell, corn	S	S	C	C	C	C	C
<i>Veronica arvensis</i>							
Spurge, spotted	•	S	C	C	C	C	C
<i>Euphorbia maculata</i>							
Vervain, blue	•	S	C	C	C	C	C
<i>Verbena hastata</i>							
Vetch	•	S	C	C	C	C	C
<i>Vicia spp.</i>							

BIENNIAL WEED SPECIES

Carrot, wild	•	•	S	S	C	C	C
<i>Daucus carota</i>							

PERENNIAL WEED SPECIES

Bahiagrass	S	S	S	S	S	S	S
<i>Paspalum notatum</i>							
Dock, curly	•	S	C	C	C	C	C
<i>Rumex crispus</i>							
Fescue, tall	S	S	S	S	S	S	S
<i>Festuca arundinacea</i>							
Greenbriar	•	•	•	•	S	S	S
<i>Smilax spp.</i>							
Honeysuckle	•	S	C	C	C	C	C
<i>Lonicera spp.</i>							
Horsenettle	•	•	S	S	S	S	S
<i>Solanum carolinense</i>							
Johnsongrass	•	S	S	S	C	C	C
<i>Sorghum halepense</i>							
Peppervine	•	S	C	C	C	C	C
<i>Ampelopsis arborea</i>							
Plantain, buckhorn	S	S	C	C	C	C	C
<i>Plantago lanceolata</i>							
Raspberry	•	•	•	•	S	S	S
<i>Rubus spp.</i>							
Ryegrass, Italian	•	•	S	S	C	C	C
<i>Lolium spp.</i>							
Trumpetcreeper	•	S	C	C	C	C	C
<i>Campsis radicans</i>							
Vaseygrass	•	S	C	C	C	C	C
<i>Paspalum urvillei</i>							

10.0 LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label pamphlet ("Directions") when used in accordance with those Directions under the conditions described therein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

To the extent consistent with applicable law, buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at

4/44

the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

This Company does not warrant any product reformulated or repackaged from this product except in accordance with this Company's stewardship requirements and with express written permission from this Company.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW,, IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this Limit of Warranty and Liability which may not be varied by any verbal or written agreement.

Campaign is a registered trademark of Monsanto Technology LLC.

All other trademarks are the property of their respective owners.

In case of an emergency involving this product,

Call Collect, day or night, (314) 694-4000.

© [DATE] MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167 USA

IV. SUPPLEMENTAL LABELING FOR NON-CROP, RANGELAND, AND INDUSTRIAL USES

Table of Contents: Non-crop, Rangeland, and Industrial Uses Supplemental Labeling

	Name	Approval Date
A	FOR AERIAL APPLICATION IN ARKANSAS ONLY	December 17, 2003

43/44

SUPPLEMENTAL LABELING

READ THE ENTIRE LABEL FOR [INSERT BRAND NAME] BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using [INSERT BRAND NAME] as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

[INSERT BRAND NAME]

Herbicide

EPA Reg. No. 524-351

Keep out of reach of children.

DANGER! PELIGRO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

[INSERT BRAND NAME] is a registered trademark of Monsanto Technology LLC.

In case of an emergency involving this product, Call Collect, day or night, 314-694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of herbicide application.

AVOID CONTACT OF HERBICIDE WITH FOLIAGE, GREEN STEMS, EXPOSED NON-WOODY ROOTS OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, BECAUSE SEVERE INJURY OR DESTRUCTION MAY RESULT.

See "GENERAL INFORMATION" and "MIXING" sections of the label booklet for [INSERT BRAND NAME] for essential product performance information.

[INSERT SPECIFIC USE DIRECTIONS HERE]

Read the "Limit of Warranty and Liability" in the label booklet for [INSERT BRAND NAME] before using. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

© [DATE] MONSANTO COMPANY

ST. LOUIS, MISSOURI 63167

44/44

A. FOR AERIAL APPLICATION IN ARKANSAS ONLY

USE DIRECTIONS

AVOID DRIFT. DO NOT APPLY INTO STILL AIR WHERE THERE IS A TEMPERATURE INVERSION LAYER LOW ENOUGH FOR FINE SPRAY PARTICLES TO BECOME SUSPENDED AND MOVE OUTSIDE THE TARGET AREA WHEN THE INVERSION LAYER MOVES. DO NOT APPLY WHEN WINDS ARE GUSTY OR UNDER ANY OTHER CONDITION THAT FAVORS DRIFT. DRIFT IS LIKELY TO CAUSE DAMAGE TO ANY VEGETATION CONTACTED. TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION, APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Use the recommended rate of this product in 3 to 15 gallons of water per acre.

Use sufficient carrier volume and appropriate equipment set-up to form droplets large enough to avoid drift potential. Coarse droplets in the 300 to 500 (VMD) micron range are recommended.

Applications should typically be made with the nozzle release point at 8 to 15 feet above the top of the target plants unless a greater height is required for aircraft safety.

The distance of the outermost nozzles on the boom must not exceed 75 percent of the length of the wingspan or rotor. In many cases, reducing this distance to 65 percent of the length of the wingspan or rotor will improve drift control without affecting the swath width.

Nozzles must always discharge backward parallel with the air stream and never discharge downwards more than 45 degrees on fixed wing aircraft or forward of the prevailing airflow on rotary winged aircraft. Avoid the use of nozzles with wide-angle discharge.

Do not apply this product when winds are in excess of 10 miles per hour.

Do not apply when there is a low-level inversion where fine spray particles could be suspended in still air and move outside the target area when the inversion layer moves. These conditions may occur when wind speeds are less than 2 miles per hour.

Use the following guidelines when applications are made near crops or other desirable vegetation:

1. Do not apply within 100 feet of any desirable vegetation or crops.
2. If wind up to 5 miles per hour is blowing toward desirable vegetation or crops, do not apply within 500 feet upwind of the desirable vegetation or crops.
3. Winds blowing from 5 to 10 miles per hour toward desirable vegetation or crops will likely require buffer zones in excess of 500 feet.

Read and carefully observe the label claims, cautionary statements, and all information on the label for [INSERT BRAND NAME].