

JAN/SAN™ HERBICIDE

by MONSANTO

Read the entire label before using this product. Use only according to label instructions. Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

HARMFUL IF SWALLOWED. MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes, or clothing.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Call a physician if irritation persists. Remove product from skin or clothing. **IF SWALLOWED,** this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Call a physician.

Personal Protective Equipment

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

In case of a human health emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Mix, store or apply this product ONLY in stainless steel, aluminum, fiberglass plastic or plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

KEEP PEOPLE OFF TREATED AREAS UNTIL SPRAY SOLUTION HAS DRIED.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: Store this product in its original container and in a secured storage area.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

This product is a nonselective herbicide that is diluted and applied to the foliage of actively growing weeds as a spot or broadcast application. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms may require a week or more to appear, with burndown usually occurring in 2 to 4 weeks. Symptoms are a gradual wilting and yellowing of the sprayed plant followed by deterioration of both shoots and roots. This product has no herbicidal activity in the soil and will not wash or leach to affect nearby vegetation.

This product controls many weeds around buildings, vacant lots, storage and recreational areas, gravel or ground bark mulch, shade trees, ornamentals and shrubs; along fences, sidewalks, driveways and patios; and prior to planting turfgrasses or shrubs listed on the label.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental garden or parks, or on golf courses or lawns and grounds.

When used as a spot treatment in lawns, all vegetation contacted will be damaged.

THIS PRODUCT CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. AVOID SPRAY DRIFT CONTACT WITH DESIRABLE LAWN GRASSES, FLOWERS, VEGETABLES, SHRUBS OR TREES. DO NOT CONTACT GREEN BARK OF TREES OR SHRUBS. IF DESIRABLE VEGETATION IS CONTACTED, WASH IMMEDIATELY WITH WATER.

WEEDS CONTROLLED: When applied as recommended under the conditions described, this product controls many common weeds including the following:

ANNUALS: Barnyardgrass, Bluegrass (annual), Chickweed, Crabgrass, Foxtail, Lambsquarters (common), Lettuce (prickly), Mustard (tansy), Pannycress (field), Pigweed (redroot, smooth), Ragweed (common), Ryegrass (annual), Sandbur (field), Shepherds-purse, Sowthistle.

PERENNIALS: Bahiagrass, Bermudagrass, Bindweed (field), Bluegrass (Kentucky), Fescue (tall), Johnsongrass, Nutsedge (purple, yellow), Orchardgrass, Broffegrass (smooth), Clover (red, white), Clover (red, white), Dallisgrass, Dandelion, Fescue species, Quackgrass, Ryegrass (perennial), Thistle (Canada), Timothy, Vaseygrass.

BRUSH: Alder, Blackberry, Cherry, Dewberry, Elderberry, Honeysuckle, Kudzu, Oak, Poison Ivy, Poison Oak, Raspberry, Rose (multiflora), Sumac, Trumpet-creeper, Willow.

CLEANING EQUIPMENT: Triple-rinse spray tank and flush all sprayer components with water to remove residues of this product. After thorough cleaning, equipment may be used to apply other products.

LAWN RENOVATION: Best results are obtained when at least one mowing is skipped prior to treatment. This provides more leaf surface to intercept the spray. For areas where renovation and reseeding is desired, thorough spray coverage of all existing vegetation is necessary. Delay planting or sodding of treated areas for at least 7 days after application.

ORNAMENTALS: Any ornamental species may be planted in treated areas 7 days or more after application. Apply this product as recommended to control undesirable plants. To avoid injury on desirable plants, prevent spray or drift contact with foliage, stems or green bark. Do not spray into or over the top of, growing ornamentals.

MIXING AND SPRAYING: Do not apply this product through any type of irrigation system.

Hand-held Sprayers - Add 3 to 6 fluid ounces (6 to 12 tbsp.) of this product to 1 gallon of clean water. Use the higher recommended rate for control of listed perennials and brush. Use pump-up-type or hand-trigger-type sprayers. USE OF HOSE-END SPRAYERS OR SPRINKLER-TYPE DEVICES MAY RESULT IN POOR AND/OR ERRATIC RESULTS.

Ensure uniform and complete spray coverage, but do not spray plants to the point that the spray solution runs off. Use a coarse spray. Do not spray during windy conditions, as spray drift may cause damage to any vegetation contacted.

WHEN TO APPLY: Apply to vigorously growing plants. Apply to annual weeds any time after emergence. For best results, apply to perennial weeds after seedheads, flowers or berries appear. On woody brush or trees, apply after leaves have fully expanded. Applications made at less than the recommended stage of growth may result in reduced control. After mowing, clipping or tilling, allow weeds to regrow to the recommended growth stage before spraying this product.

Rainfall or watering within 6 hours after application can wash the product off plant foliage which may require retreatment. Delay mowing, clipping, tilling, planting or sodding of treated areas for at least 7 days after application.

application. This allows time for this product to move within the plant for most effective results. SINCE THIS PRODUCT DOES NOT WORK IN SOIL, RETREATMENTS WILL BE NECESSARY TO CONTROL NEW WEEDS EMERGING FROM SEED.

One application will control most weeds listed. Symptoms on certain tough perennial weeds and woody brush and tree species may be slow to develop. If control is not achieved within 4 weeks, allow time for the sprayed weeds to regrow and make a repeat application. In some cases, control of woody brush and trees treated in the fall may not be fully apparent until the following season.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGE RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

ACTIVE INGREDIENT:

Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt.....18.0%

INERT INGREDIENTS:.....82.0%

100.0%

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license is granted under any non-U.S. patent(s).


MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

™Trademark of Monsanto Company.
© MONSANTO COMPANY 1994

NET GAL

EPA Est. 524-LA-1

21027S1-1L



Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-370

524-370

08/01/2000

117

Carefully read and follow detailed instructions in pamphlet.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION! HARMFUL IF SWALLOWED. MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing.

IF SWALLOWED, this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Mix, store or apply this product ONLY in stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers. DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. Use of this product in such containers could result in the formation of an explosive hydrogen gas mixture which could flash or explode if ignited by open flames or any other ignition source.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

Trademark of Monsanto Company
©MONSANTO COMPANY 1991
MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.
EPA Reg. No. 524-370
EPA Est. 524-LA-1
897.18-000.63L-2



▲ Cut or tear along perforation

Roundup

CONCENTRATE

by Monsanto

Makes Up To 36 Gallons

Grass & Weed Killer

- COVERS UP TO 10,800 SQ. FT.
- KILLS UNWANTED LABELED GRASSES AND WEEDS SYSTEMATICALLY—ROOTS AND ALL
- DOES NOT HAVE SOIL ACTIVITY
- SUITABLE FOR USE AROUND FLOWER BEDS, TREES, SHRUBS, FENCES AND DRIVEWAYS
- USE TO PREPARE GARDEN SITES
- MIX 6 TO 12 TABLESPOONS (3-6 OZ) PER GALLON

Store in secured place in original container.
Read the entire label before using this product.
Use only according to label instructions.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions. If these terms are not acceptable, return at once unopened.

Keep out of reach of children.

CAUTION! Read precautions on back panel

ACTIVE INGREDIENTS:

*Glyphosate, N-(phosphonomethyl)-glycine, in the form of its isopropylamine salt	18%
INERT INGREDIENTS:	82%
	100%

NET 108.8 fl. oz.
(3 qts. 12.8 oz.) 897.18-000.63-1

2/17

KEEP PEOPLE AND PETS OFF TREATED AREAS UNTIL SPRAY SOLUTION HAS DRIED.

DOMESTIC ANIMALS: This product is considered to be relatively nontoxic to dogs and other domestic animals. However, ingestion of this product or large amounts of freshly sprayed vegetation may result in temporary gastrointestinal irritation (vomiting, diarrhea, colic, etc.) If such symptoms are observed, provide the animal with plenty of fluids to prevent dehydration. Call a veterinarian if symptoms persist for more than 24 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Roundup L&G is a nonselective herbicide that is diluted and applied to the foliage of actively growing weeds as a spot or broadcast application. It is absorbed by the leaves and moves throughout the stem and roots to control the entire plant. Visible symptoms may require a week or more to appear, with burndown usually occurring in 2 to 4 weeks. Symptoms are a gradual wilting and yellowing of the sprayed plant followed by deterioration of both shoots and roots. This product has no herbicidal activity in the soil and will not wash or leach to affect nearby vegetation.



1

This product controls many weeds around buildings, vacant lots, storage and recreational areas, gravel or ground bark mulch, shade trees, ornamentals and shrubs; along fences, sidewalks, driveways and patios; and prior to planting turf-grasses, shrubs, flower beds or vegetables listed on the label. When used as a spot treatment in lawns, all vegetation contacted will be damaged.

THIS PRODUCT CAN INJURE OR DESTROY ALL VEGETATION CONTACTED. AVOID SPRAY DRIFT CONTACT WITH DESIRABLE LAWN GRASSES, FLOWERS, VEGETABLES, SHRUBS OR TREES. DO NOT CONTACT GREEN BARK OF TREES OR SHRUBS. IF DESIRABLE VEGETATION IS CONTACTED, WASH IMMEDIATELY WITH WATER.

For more product information, call toll-free 1-800-225-2883.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Store this product in its original container and in a secured storage area. In case of SPILL or LEAK, soak up with paper towels and discard in trash.

DISPOSAL:

To dispose of unused product, securely wrap original container in several layers of newspaper and discard in trash.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is discarded.

Do not reuse container. Rinse thoroughly before discarding in trash.

2

WEEDS CONTROLLED

When applied as recommended under the conditions described, Roundup L&G controls many common weeds including the following:

Annuals

Barnyardgrass	Pennycress (field)
Bluegrass (annual)	Pigweed (redroot, smooth)
Chickweed	Ragweed (common)
Crabgrass	Ryegrass (annual)
Foxtail	Sandbur (field)
Lambsquarter (common)	Shepherdspurse
Lettuce (prickly)	Southistle
Mustard (tansy)	

Perennials

Bahiagrass	Fescue (tall)
Bermudagrass	Johnsongrass
Bindweed (field)	Nutsedge (purple, yellow)
Bluegrass (Kentucky)	Orchardgrass
Bromegrass (smooth)	Quackgrass
Clover (red, white)	Ryegrass (perennial)
Dallisgrass	Thistle (Canada)
Dandelion	Timothy
Fescue species	Vaseygrass

Brush

Alder	Poison Ivy
Blackberry	Poison Oak
Cherry	Raspberry
Dewberry	Rose (multiflora)
Elderberry	Sumac
Honeysuckle	Trumpetcreeper
Kudzu	Willow
Oak	

3

MIXING AND

Do not apply to

Hand-held applicators of this product are not recommended. Use only hose-end applicators. RESULTS.

Ensure uniform coverage of spray plants. Do not use a coarse spray nozzle to avoid spray drift.

CLEANING

Triple-rinse applicator with water. Thoroughly clean applicator products.

WHEN TO

Apply to vegetation anytime after flowering. On woody plants, apply when expanded. Apply to vegetation at mowing, clipping, or other recommended

Rainfall or watering within 6 hours after application can wash the product off plant foliage which may require retreatment. Delay mowing, clipping, tilling, planting or sodding of treated areas for at least 7 days after application. This allows time for this product to move within the plant for most effective results. **SINCE THIS PRODUCT DOES NOT WORK IN SOIL, RETREATMENTS WILL BE NECESSARY TO CONTROL NEW WEEDS EMERGING FROM SEED.**

One application will control most weeds listed. Symptoms on certain tough perennial weeds and woody brush and tree species may be slow to develop. If control is not achieved within 4 weeks, allow time for the sprayed weeds to regrow and make a repeat application. In some cases, control of woody brush and trees treated in the fall may not be fully apparent until the following season.

LAWN RENOVATION

Best results are obtained when at least one mowing is skipped prior to treatment. This provides more leaf surface to intercept the spray. For areas where renovation and reseedling is desired, thorough spray coverage of all existing vegetation is necessary. Delay planting or sodding of treated areas for at least 7 days after application.

ORNAMENTALS

Any ornamental species may be planted in treated areas 7 days or more after application. Apply this product as recommended to control undesirable vegetation growing around ornamentals, trees or shrubs. Avoid spray contact with any desirable plants. To avoid injury on desirable plants, prevent spray or drift contact with foliage, stems or green bark. Do not spray into, or over the top of, growing ornamentals.


TREES (Apple, Citrus, Nut, Pear) AND GRAPEVINES

Apply this product to control undersirable plants growing around these trees and vines. Spray contact, other than with mature bark of the main trunk, can injure the tree or vine. Allow 21 days between application and harvest.

BERRIES, SMALL FRUITS & VEGETABLE GARDENS

The following vegetables may be planted in treated areas 7 days or more after application: asparagus, beans, beet greens, beets, blackberry, blueberry, boysenberry, broccoli, cabbage, carrot, cauliflower, celery, chicory, corn, cranberry, cucumber, currant, dewberry, elderberry, eggplant, garlic, gourds, gooseberry, horseradish, huckleberry, Jerusalem artichoke, kale, lentils, lettuce, loganberry, melons, mustard greens, okra, ollalieberry, onion, parsnips, peas, peppers, Irish and sweet potatoes, pumpkin, radish, raspberry (black, red), rutabagas, spinach, squash (summer, winter), strawberry, tomatillo, tomato*, watercress, watermelon and turnips. In the treated area, do not plant any vegetables not listed on this label for one year following application. Do not apply this product in or around growing vegetables.

*Use on tomatoes is restricted to direct seeded tomatoes only.



**ROUNDUP
L&G CONCENTRATE
GRASS & WEED KILLER**

Lawn and Garden formulation which controls labeled weeds and grasses.

Read the entire label before using this product. Use only according to label instructions.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS. IF THESE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.

REFORMULATION OR REPACKAGING IS PROHIBITED.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt 18.0%
INERT INGREDIENTS: 82.0%
100.0%

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

©MONSANTO COMPANY 1991

MONSANTO COMPANY
 AGRICULTURAL PRODUCTS
 ST. LOUIS, MISSOURI 63167 U.S.A.



Carefully read and follow detailed instructions in pamphlet.
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION! HARMFUL IF SWALLOWED. MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing.

IF SWALLOWED, this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Mix, store or apply this product ONLY in stainless steel, aluminum, fiberglass, plastic or plastic lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** Use of this product in such containers could result in the formation of an explosive hydrogen gas mixture which could flash or explode if ignited by open flames or any other ignition source.

See pamphlet for additional information on storage and disposal.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

© Trademark of Monsanto Company ©MONSANTO COMPANY 1991
 MONSANTO COMPANY • AGRICULTURAL PRODUCTS
 ST. LOUIS, MISSOURI 63167 U.S.A.
 EPA Reg. No. 524-370

▲ Cut or tear along perforation

Printed in U.S.A. by the following: 10/11/88

4/17

5/17

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE DIRECTIONS IN THE LABEL AND OBSERVE ALL PRECAUTIONS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION HARMFUL IF SWALLOWED. MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing. **IF SWALLOWED,** this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

PRECAUTION: SEE "DIRECTIONS FOR USE" FOR IMPORTANT CONTAINER PRECAUTIONS, and inside label for storage and disposal information.

EPA Reg. No. 524-370

EPA Est. (G)56100-LA-001; (L)524-LA-1; (I)239-1A-3. Lot number prefix G, L, or I indicates appropriate establishment number.



PULL HERE TO OPEN/PRESS TO RESEAL

897-18-000-84-2X3
MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

Kills Weeds
Roots and All
With No
Soil Activity

WEED & GRASS KILLER

Roundup

CONCENTRATE

Keep out of reach of children. **Makes up to 5 Gallons**
Covers up to 1,500 Sq. Ft.

CAUTION Read precautions and directions in and on back panel.

ACTIVE INGREDIENT:
Glyphosate, N-(phosphono-
methyl)glycine, in the form
of its isopropylamine salt..... 18.0%
INERT INGREDIENTS..... 82.0%
100.0%

NET 16 fl oz
897-18-000-84-1

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS. IF THESE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.


Environmental Hazards
Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

REFORMULATION OR REPACKAGING FOR SALE IS PROHIBITED.

©Trademark of Monsanto Company
©MONSANTO COMPANY 1993

THIS PANEL IS INTENTIONALLY LEFT BLANK.



**ROUNDUP® CONCENTRATE
USE DIRECTIONS TABLE**

Follow mixing rate recommendations in this Use Directions Table. Ensure uniform and complete spray coverage. Sprayed plants usually show effects within 7 days, with complete kill in 2 to 4 weeks. For best results, apply to actively growing undesirable weeds, grasses, trees, brush and plants in warm, sunny weather. If not completely killed within 4 weeks, repeat application. In some cases, control of brush and trees sprayed in the fall may not be fully apparent until the following season. Sprayed areas can be reseeded or replanted, 7 days after spraying.

USES	SPRAYING INSTRUCTIONS	WEEDS CONTROLLED	MIXING RATE
SPOT SPRAYING/TRIM and EDGE Undesirable plants in and around: <ul style="list-style-type: none"> • fences • trees* • driveways • flower beds • walkways • shrubs 	Apply to actively growing plants. Apply to annual weeds anytime after emergence. For perennial weeds, apply after seed heads or flowers appear. Avoid spray contact with foliage, stems or green bark of desirable plants.	Most annual and perennial weeds and grasses such as chickweed, ragweed, dandelion (Complete list follows this table.)	ANNUALS - 3 fl oz (6 tbsp) per 1 gallon water PERENNIALS and BRUSH - 6 fl oz (12 tbsp) per 1 gallon water 1) Use clean water. 2) Ensure uniform and complete spray coverage. Position sprayer tip 1-2 feet from weeds.

*Note: If Roundup is used to control undesirable plants around fruit or nut trees or grapevines, allow 21 days before eating the fruits or nuts.

WEEDS CONTROLLED

When applied as recommended under the conditions described in the Use Directions Table, Roundup Concentrate controls many common weeds including the following:

Annuals	
Barnyardgrass	Pennycress (field)
Bluegrass (annual)	Pigweed (redroot, smooth)
Chickweed	Ragweed (common)
Crabgrass	Ryegrass (annual)
Foxtail	Sandbur (field)
Lambsquarter (common)	Shepherdspurse
Lettuce (prickly)	Sowthistle
Mustard (fancy)	
Perennials	
Bahiagrass	Fescue (tall)
Bermudagrass	Johnsongrass
Bindweed (field)	Nutsedge (purple, yellow)
Bluegrass (Kentucky)	Orchardgrass
Bromegrass (smooth)	Quackgrass
Clover (red, white)	Ryegrass (perennial)
Dallisgrass	Thistle (Canada)
Dandelion	Timothy
Fescue species	Vaseygrass
Brush	
Alder	Poison Ivy
Blackberry	Poison Oak
Cherry	Raspberry
Dewberry	Rose (multiflora)
Elderberry	Sumac
Honeysuckle	Trumpet creeper
Kudzu	Willow
Oak	

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE FOLLOWING DIRECTIONS AND OBSERVE ALL PRECAUTIONS.**

Roundup® Concentrate Weed and Grass Killer, when absorbed by contacted foliage, is carried throughout the stems and roots to give complete kill of tough annual and perennial weeds and grasses. Roundup is broken down in the soil by microbes. Once in contact with soil, it has no more weed killing activity and will not move in or on the soil to affect nearby plants.

HOW TO USE

IMPORTANT: This product is an all-purpose weed, grass, and brush killer which can injure or kill all plants contacted (virtually anything that is green and growing). **APPLY ONLY WHEN AIR IS CALM. AVOID SPRAY CONTACT ON DESIRABLE PLANTS. IF NECESSARY, SHIELD DESIRABLE PLANTS TO PREVENT CONTACT WITH SPRAY.** If spray or drift accidentally contacts desirable plants, wash off immediately with water. Rainfall within 6 hours could affect performance. Avoid mowing, cutting or otherwise disturbing sprayed plants for at least 7 days. People and pets should stay off sprayed area until thoroughly dry to ensure that product is not tracked into areas where desirable plants are growing.

SPRAY EQUIPMENT

Use pump-up-type or hand-trigger type sprayers to apply this product. Triple-rinse sprayer and flush all sprayer components with water after use to remove residues of this product. After very thorough cleaning, equipment may be used to apply other products. Apply rinse waters to undesirable plants not previously sprayed.

WARNING: Mix, spray, apply and store solutions of this product only in stainless steel, aluminum, fiberglass, plastic or plastic lined metal sprayers. **DO NOT MIX, APPLY, OR STORE THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) SPRAYERS.** Because the product and such containers can react to produce a highly combustible mixture of hydrogen gas. This buildup of hydrogen gas in a closed container can create enough pressure to cause the container to rupture, or can flash or explode, causing serious personal injury, if ignited by open flame, spark, welders torch, lit cigarette or any other ignition source.

Do not apply this product through any type of irrigation system, nor use in hose-end sprayers or sprinkler-type devices. These applications may result in poor and/or erratic results.

USES	SPRAYING INSTRUCTIONS	WEEDS CONTROLLED	MIXING RATE
LAWN RENOVATION • Replacement of worn out lawns • Changing to more desirable grass	For best results on lawns, skip at least one mowing prior to spraying. Delay reseeding, sprigging, tilling or sodding for 7 days after spraying. Early spring or fall spraying recommended.	Most annual and perennial weeds and grasses such as chickweed, ragweed, dandelion (Complete list follows this table.)	ANNUALS - 3 fl oz (6 tbsp) per 1 gallon water PERENNIALS and BRUSH - 6 fl oz (12 tbsp) per 1 gallon water
GARDEN/ORNAMENTAL BED PREPARATION FOR PLANTING: • Flowers • Trees, Shrubs • Vegetables • Berries and small fruits	Any ornamental may be planted 7 days after spraying. Vegetables, berries and small fruits may be planted 7 days after spraying. (Complete list follows this table.)		1) Use clean water. 2) Ensure uniform and complete spray coverage. Position sprayer tip 1-2 feet from weeds.
BRUSH CONTROL TO KILL: • Woody brush • Trees • Vines	Spray when leaves are fully expanded (spring-early summer) or when flowers or berries appear (mid-summer - fall). Spray trees and brush up to 5 feet in height. For taller plants, cut down and spray regrowth.	Most brush, such as poison ivy, blackberry and kudzu (Complete list follows.)	

VEGETABLES - SMALL FRUITS AND BERRIES

The following vegetables, small fruits and berries may be planted (by seed, rooted cuttings or transplants) 7 days after spraying. If it is desired to plant vegetables, small fruits and berries other than those in the below list, wait for 30 days after spraying.

VEGETABLES

ARTICHOKE, JERUSALEM	LEEK
ASPARAGUS	LENTILS
BEANS (ALL)	LETTUCE
BEEF GREENS	MANGO MELON
BEETS (RED, SUGAR)	MELONS (ALL)
BROCCOLI (ALL)	MUSKMELON
BRUSSELS SPROUTS	MUSTARD GREENS
CABBAGE (ALL)	OKRA
CANTALOUPE	ONION
CARROT	RAPE GREENS
CAULIFLOWER	PARSLEY
CASABA MELON	PARSNIPS
CELERY	PEAS (ALL)
CHARD, SWISS	PEPPER (ALL)
CHICORY	PERSIAN MELON
COLLARDS	POTATO (IRISH, SWEET)
CORN	PUMPKIN
CRENSHAW MELON	RADISH
CUCUMBER	RHUBARB
EGGPLANT	RUTABAGA
ENDIVE	SPINACH (ALL)
GARLIC	SQUASH (SUMMER, WINTER)
GOURDS	TOMATILLO
GROUND CHERRY	TOMATO*
HONEYDEW MELON	TURNIP
HONEY BALL MELON	WATERCRESS
HORSERADISH	WATERMELON
KALE	YAMS
KOHLRABI	

*Tomatoes from seed only. Wait 30 days for rooted transplants.

SMALL FRUITS AND BERRIES

BLACKBERRY
BLUEBERRY
BOYSENBERRY
CRANBERRY
CURRANT
DEWBERRY
ELDERBERRY

GOOSEBERRY
HUCKLEBERRY
LOGANBERRY
OLALLIEBERRY
RASPBERRY (BLACK, RED)
STRAWBERRY

STORAGE AND DISPOSAL

Storage: Store this product only in its original container and in a secured storage area. Protect container from freezing.

PRODUCT DISPOSAL:

Monsanto is committed to responsible environmental practices and recommends that all of the product be used up, carefully following all label directions and precautions.

If necessary to dispose of partially filled product container, then securely wrap it in several layers of newspaper and discard in trash.

EMPTY CONTAINER:

Do not reuse container. Emptied container retains vapor and product residue. Observe all labeled safeguards until container is discarded.

Securely wrap empty container in several layers of newspaper and discard in trash.

SPILL OR LEAK:

In case of spill or leak, soak up with paper towels and discard in trash.

For more product information or questions, call toll-free 1-800-225-2883, 8:00AM - 4:00PM Pacific Standard Time. MON-FAY-FRIDAY

THIS PANEL IS INTENTIONALLY LEFT BLANK.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE DIRECTIONS IN THE LABEL AND OBSERVE ALL PRECAUTIONS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals
CAUTION HARMFUL IF SWALLOWED. MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing. IF SWALLOWED, this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

PRECAUTION: SEE "DIRECTIONS FOR USE" FOR IMPORTANT CONTAINER PRECAUTIONS, and inside label for storage and disposal information.

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.
EPA Reg. No. 524-370

EPA Est. (G)56100-LA-001; (L)524-LA-1; (I)239-IA-3. Lot number prefix G, L, or I indicates appropriate establishment number.

897.18-000.84-3X3

8/17

Kills Weeds
Roots and All
With No
Soil Activity

WEED & GRASS

KILLER Roundup®

CONCENTRATE

Keep out of reach of children.

CAUTION Read precautions and directions
in and on back panel.

ACTIVE INGREDIENT:
 Glyphosate, N-(phosphonomethyl)glycine,
 in the form of its isopropylamine salt 18.0%
 INERT INGREDIENTS 82.0%
 100.0%

Makes up to 10 Gallons
Covers up to 3,000 Sq. Ft.

NET 32 fl oz
897.18-001.82-1

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE DIRECTIONS IN THE LABEL AND OBSERVE ALL PRECAUTIONS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION HARMFUL IF SWALLOWED. MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing.

IF SWALLOWED, this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product, call Collect, day or night, (314) 694-4000.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS. IF THESE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.

Environmental Hazards:

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

PRECAUTION: SEE "DIRECTIONS FOR USE" FOR IMPORTANT CONTAINER PRECAUTIONS, and inside label for storage and disposal information.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

REFORMULATION OR REPACKAGING FOR SALE IS PROHIBITED.


©Trademark of Monsanto Company
©MONSANTO COMPANY 1993

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167
U.S.A.
EPA Reg. No. 524-370

EPA Est. (G)56100-LA-001;
(L)524-LA-1; (I)239-IA-3

Lot number prefix G.L. or I indicates appropriate establishment number.

897.18-000 82-2X3



PULL HERE TO OPEN/PRESS TO RESEAL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. **READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE FOLLOWING DIRECTIONS AND OBSERVE ALL PRECAUTIONS.**

Roundup® Concentrate Weed and Grass Killer, when absorbed by contacted foliage, is carried throughout the stems and roots to give complete kill of tough annual and perennial weeds and grasses. Roundup is broken down in the soil by microbes. Once in contact with soil, it has no more weed killing activity and will not move in or on the soil to affect nearby plants.

HOW TO USE

IMPORTANT: This product is an all-purpose weed, grass, and brush killer which can injure or kill all plants contacted (virtually anything that is green and growing). **APPLY ONLY WHEN AIR IS CALM. AVOID SPRAY CONTACT ON DESIRABLE PLANTS. IF NECESSARY, SHIELD DESIRABLE PLANTS TO PREVENT CONTACT WITH SPRAY.** If spray or drift accidentally contacts desirable plants, wash off immediately with water. Rainfall within 6 hours could affect performance. Avoid mowing, cutting or otherwise disturbing sprayed plants for at least 7 days. People and pets should stay off sprayed area until thoroughly dry to ensure that product is not tracked into areas where desirable plants are growing.

SPRAY EQUIPMENT

Use pump-up-type or hand-trigger type sprayers to apply this product. Triple-rinse sprayer and flush all sprayer components with water after use to remove residues of this product. After very thorough cleaning, equipment may be used to apply other products. Apply rinse waters to undesirable plants not previously sprayed.

WARNING: Mix, spray, apply and store solutions of this product only in stainless steel, aluminum, fiberglass, plastic or plastic-lined steel sprayers. **DO NOT MIX, APPLY, OR STORE THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) SPRAYERS,** because the product and such containers can react to produce a highly combustible mixture of hydrogen gas. This buildup of hydrogen gas in a closed container can create enough pressure to cause the container to rupture, or can flash or explode, causing serious personal injury, if ignited by open flame, spark, welders torch, lit cigarette or any other ignition source.

Do not apply this product through any type of irrigation system, nor use in hose-end sprayers or sprinkler-type devices. These applications may result in poor and/or erratic results.

**ROUNDUP® CONCENTRATE
USE DIRECTIONS TABLE**

Follow mixing rate recommendations in this Use Directions Table. Ensure uniform and complete spray coverage. Sprayed plants usually show effects within 7 days, with complete kill in 2 to 4 weeks. For best results, apply to actively growing undesirable weeds, grasses, trees, brush and plants in warm, sunny weather. If not completely killed within 4 weeks, repeat application. In some cases, control of brush and trees sprayed in the fall may not be fully apparent until the following season. Sprayed areas can be reseeded or replanted, 7 days after spraying.

USES	SPRAYING INSTRUCTIONS	WEEDS CONTROLLED	MIXING RATE
SPOT SPRAYING/ TRIM and EDGES Undesirable plants in and around: • fences • trees • driveways • flower beds • walkways • shrubs	Apply to actively growing plants. Apply to annual weeds anytime after emergence. For perennial weeds, apply after seed heads or flowers appear. Avoid spray contact with logs, stems or green bark of desirable plants.	Most annual and perennial weeds and grasses such as chickweed, ragweed, and dandelion. (Complete list follows this table.)	ANNUALS 3 tps (6 tbs) per 1 gallon water PERENNIALS and BRUSH 6 fl oz (12 tbs) per 1 gallon water 1) Use clean water. 2) Ensure uniform and complete spray coverage. Position sprayer tip 1-2 feet from weeds.

*Note: If Roundup is used to control undesirable plants around fruit or nut trees or grapevines, allow 21 days before eating the fruits or nuts.

USE DIRECTIONS TABLE (CONTINUED)

USES	SPRAYING INSTRUCTIONS	WEEDS CONTROLLED	MIXING RATE
LAWN RENOVATION • Replacement of worn out lawns • Changing to more desirable grass	For best results on lawns, skip at least one mowing prior to spraying. Delay reseeding, sprigging or sodding for 7 days after spraying. Earliest spring or fall spraying recommended.	Most annual and perennial weeds and grasses such as chickweed, pigweed, dandelion (Complete list follows this table.)	ANNUALS - 3 fl oz (6 tbsp) per 1 gallon water PERENNIALS and BRUSH - 6 fl oz (12 tbsp) per 1 gallon water
GARDEN/ORNAMENTAL BED PREPARATION FOR PLANTING: • Flowers • Trees, Shrubs • Vegetables • Berries and small fruits	Any ornamental may be planted 7 days after spraying. Vegetables, berries and small fruits may be planted 7 days after spraying. (Complete list follows this table.)		1) Use clean water. 2) Ensure uniform and complete spray coverage. Position sprayer tip 1-2 feet from weeds.
BRUSH CONTROL TO KILL: • Woody brush • Trees • Vines	Spray when leaves are fully expanded (spring-early summer) or when flowers or berries appear (mid-summer - fall). Spray trees and brush up to 5 feet in height. For taller plants, cut down and spray regrowth.	Most brush, such as poison ivy, blackberry and kudzu (Complete list follows.)	

WEEDS CONTROLLED

When applied as recommended under the conditions described in the Use Directions Table, Roundup Concentrate controls many common weeds including the following:

Annuals

- Barnyardgrass
- Bluegrass (annual)
- Chickweed
- Crabgrass
- Foxtail
- Lambsquarter (common)
- Lettuce (prickly)
- Mustard (tansy)

- Pennycress (field)
- Pigweed (red root, smooth)
- Ragweed (common)
- Ryegrass (annual)
- Sanibar (field)
- Shepherdspurse
- Sowthistle

Perennials

- Bahiagrass
- Bermudagrass
- Bindweed (field)
- Bluegrass (Kentucky)
- Bromegrass (smooth)
- Clover (red, white)
- Dallisgrass
- Dandelion
- Fescue species

- Fescue (tall)
- Johnsongrass
- Nutsedge (purple, yellow)
- Orchardgrass
- Quackgrass
- Ryegrass (perennial)
- Timothy (Canada)
- Timothy
- Vaseygrass

Brush

- Alder
- Blackberry
- Cherry
- Chokeberry
- Elderberry
- Honeysuckle
- Kudzu
- Oak

- Poison Ivy
- Poison Oak
- Raspberry
- Rose (multiflora)
- Sumac
- Trumpet creeper
- Willow

VEGETABLES, SMALL FRUITS AND BERRIES

The following vegetables, small fruits and berries may be planted (by seed, rooted cuttings or transplants) 7 days after spraying. If it is desired to plant vegetables, small fruits and berries other than those in the below list, wait for 30 days after spraying.

VEGETABLES

ARTICHOKE, JERUSALEM
 ASPARAGUS
 BEANS (ALL)
 BEET GREENS
 BEETS (RED, SUGAR)
 BROCCOLI (ALL)
 BRUSSELS SPROUTS
 CABBAGE (ALL)
 CANTALOUPE
 CARROT
 CAULIFLOWER
 CASABA MELON
 CELERY
 CHARD, SWISS
 CHICORY
 COLLARDS
 CORN
 CRENSHAW MELON
 CUCUMBER
 EGGPLANT
 ENDIVE
 GARLIC
 GOURDS
 GROUND CHERRY
 HONEYDEW MELON
 HONEY BALL MELON
 HORSERADISH
 KALE
 KOHLRABI

LEEK
 LENTILS
 LETTUCE
 MANGO MELON
 MELONS (ALL)
 MUGKMELO
 MUSTARD GREENS
 OKRA
 ONION
 RAPE GREENS
 PARSLEY
 PARSNIPS
 PEAS (ALL)
 PEPPER (ALL)
 PERSIAN MELON
 POTATO (IRISH, SWEET)
 PUMPKIN
 RADISH
 RHUBARB
 RUTABAGA
 SPINACH (ALL)
 SQUASH (SUMMER, WINTER)
 TOMATILLO
 TOMATO
 TURNIP
 WATERCRESS
 WATERMELON
 YAMS

SMALL FRUITS AND BERRIES

BLACKBERRY
 BLUEBERRY
 BOYSENBERRY
 CRANBERRY
 CURRANT
 DEWBERRY
 ELDERBERRY

GOOSEBERRY
 HUCKLEBERRY
 LOGANBERRY
 OLLALIEBERRY
 RASPBERRY (BLACK, RED)
 STRAWBERRY

*Tomatoes from seed only. Wait 30 days for rooted transplants.

STORAGE AND DISPOSAL

Storage: Store this product only in its original container and in a secured storage area. Protect container from freezing.

PRODUCT DISPOSAL:

Monsanto is committed to responsible environmental practices and recommends that all of the product be used up, carefully following all label directions and precautions.

If necessary to dispose of partially filled product container, then securely wrap it in several layers of newspaper and discard in trash.

EMPTY CONTAINER:


Do not reuse container. Empty container retains vapor and product residue. Observe all labeled safeguards until container is discarded.

Securely wrap empty container in several layers of newspaper and discard in trash.

SPILL OR LEAK:

In case of spill or leak, soak up with paper towels and discard in trash.

For more product information or questions, call toll-free 1-800-225-2883, 8:00AM - 4:00PM Pacific Standard Time, MONDAY-FRIDAY



READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE DIRECTIONS IN THE USE DIRECTIONS TABLE AND OBSERVE ALL PRECAUTIONS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION HARMFUL IF SWALLOWED. MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing.

IF SWALLOWED, this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS. IF THESE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.

Environmental Hazards

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

PRECAUTION: SEE "DIRECTIONS FOR USE" FOR IMPORTANT CONTAINER PRECAUTIONS, and inside label for storage and disposal information.



This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

REFORMULATION OR REPACKAGING FOR SALE IS PROHIBITED.

©Trademark of Monsanto Company ©MONSANTO COMPANY 1993
MONSANTO COMPANY, AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.
EPA Reg. No. 524-370

EPA Est. (G)56100-LA-001; (L)524-LA-1; (I)239-1A-3
Lot number prefix G, L, or I indicates appropriate establishment number.

897.18-000.82-3X3



THIS PAGE IS INTENTIONALLY LEFT BLANK.

13/17

Kills Brush
Roots and All
With No
Soil Activity

BRUSH
KILLER

Roundup[®]

CONCENTRATE

Keep out of reach of children.

CAUTION Read precautions and directions in and on back panel.

ACTIVE INGREDIENT:

Glyphosate, N-(phosphonomethyl)glycine, in the form of its isopropylamine salt	18.0%
INERT INGREDIENTS	82.0%
	100.0%

Makes up to 8 Gallons
Covers up to 2,400 Sq. Ft.

NET 32 fl oz

897.18-100.11-1

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE DIRECTIONS IN THE LABEL AND OBSERVE ALL PRECAUTIONS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION HARMFUL IF SWALLOWED. MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing. IF SWALLOWED, this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product. Call Collect, day or night, (314) 694-4000.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS. IF THESE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.

Environmental Hazards

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

PRECAUTION: SEE "DIRECTIONS FOR USE" FOR IMPORTANT CONTAINER PRECAUTIONS, and inside label for storage and disposal information.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

REFORMULATION OR REPACKAGING FOR SALE IS PROHIBITED.

©Trademark of Monsanto Company

©MONSANTO COMPANY 1993

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.

EPA Est. (G)56100-LA-001;
(L)524-LA-1, (I)239-IA-3

Lot number prefix G,L, or I indicates appropriate establishment number.

EPA Reg. No. 524-370



897.18-100.11-2X3

PULL HERE TO OPEN/PRESS TO RESEAL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE FOLLOWING DIRECTIONS AND OBSERVE ALL PRECAUTIONS.

Roundup® Concentrate Brush Killer, when absorbed by contacted foliage, is carried throughout the stems and roots to give complete kill of woody brush, trees and vines. Roundup is broken down in the soil by microbes. Once in contact with soil, it has no more weed killing activity and will not move in or on the soil to affect nearby plants.

HOW TO USE

IMPORTANT: This product is an all-purpose weed, grass, brush, tree and vine killer which can injure or kill all plants contacted (virtually anything that is green and growing). APPLY ONLY WHEN AIR IS CALM. AVOID SPRAY CONTACT ON DESIRABLE PLANTS. IF NECESSARY, SHIELD DESIRABLE PLANTS TO PREVENT CONTACT WITH SPRAY. If spray or drift accidentally contacts desirable plants, wash off immediately with water. Rainfall within 6 hours could affect performance. Avoid mowing, cutting or otherwise disturbing sprayed plants for at least 2 to 4 weeks, or until visible effects of brownout or defoliation are observed. People and pets should stay off sprayed area until thoroughly dry to ensure that product is not tracked into areas where desirable plants are growing.

SPRAY EQUIPMENT

Use pump-up-type or hand-trigger type sprayers to apply this product. Triple-rinse sprayer and flush all sprayer components with water after use to remove residues of this product. After very thorough cleaning, equipment may be used to apply other products. Apply rinse waters to undesirable plants not previously sprayed.

WARNING: Mix, spray, apply and store solutions of this product only in stainless steel, aluminum, fiberglass, plastic or plastic-lined steel sprayers. DO NOT MIX, APPLY, OR STORE THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) SPRAYERS, because the product and such containers can react to produce a highly combustible mixture of hydrogen gas. This buildup of hydrogen gas in a closed container can create enough pressure to cause the container to rupture, or can flash or explode, causing serious personal injury, if ignited by open flame, spark, welders torch, lit cigarette or any other ignition source.

Do not apply this product through any type of irrigation system, nor use in hose-end sprayers or sprinkler-type devices. These applications may result in poor and/or erratic results.

ROUNDUP® CONCENTRATE BRUSH KILLER USE DIRECTIONS TABLE

Follow mixing rate recommendations in this Use Directions Table. Ensure uniform and complete spray coverage. Sprayed plants usually show effects such as gradual yellowing and wilting within 7-14 days, with brownout and defoliation occurring by 4 to 8 weeks. For best results, apply to actively growing undesirable woody brush, trees, vines and plants in warm, sunny weather. If not completely killed within 8 weeks, repeat application. In some cases, control of brush and trees sprayed in the fall may not be fully apparent until the following season. Sprayed areas can be reseeded or replanted, when undesirable plants are dead.

USES	SPRAYING INSTRUCTIONS	PLANTS CONTROLLED	MIXING RATE
WOODY BRUSH, TREE AND VINE KILL AND CONTROL: <ul style="list-style-type: none"> • Around houses, buildings, wooded lots, vacant lots, storage and recreational areas • Along fences and driveways 	Apply to vigorously growing woody plants. Spray when leaves are fully expanded (spring-early summer) or when flowers or berries appear (mid-summer - fall). Spray trees and brush up to 5 feet in height. For taller plants, cut down and spray regrowth.	Most brush such as poison ivy, blackberry and kudzu (Complete list follows).	4-6* fl oz (8-12 tbsp) per 1 gallon water 1) Use clean water. 2) Ensure uniform and complete spray coverage. Position sprayer tip 1-2 feet from weeds. *For tough to control plants.

ACCEPTED

AUG - 1 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No 524-370

BRUSH AND WOODY PLANTS

When applied as recommended under the conditions described in the Directions for Use Table, Roundup Concentrate Brush Killer CONTROLS or PARTIALLY CONTROLS the following woody brush and vines.

Alder	Eucalyptus	Raspberry
Ash	Hazel	Rose (multiflora)
Aspen	Hawthorn	Sage
Blackberry	Honeysuckle	Sagebrush
Broom (French, Scotch)	Kudzu	Salmonberry
Buckwheat	Locust	Sassaparilla
Caenothus	Maple	Sumac
Chamise	Oak	Sweetgum
Cherry	Persimmon	Thimbleberry
Coyote Brush	Poison Ivy	Tree Tobacco
Dewberry	Poison Oak	Trumpet creeper
Elderberry	Poplar	Willow
Elm		

STORAGE AND DISPOSAL

Storage: Store this product only in its original container and in a secured storage area. Protect container from freezing.

PRODUCT DISPOSAL:

Monsanto is committed to responsible environmental practices and recommends that all of the product be used up, carefully following all label directions and precautions.

If necessary to dispose of partially filled product container, then securely wrap it in several layers of newspaper and discard in trash.

EMPTY CONTAINER:

Do not reuse container. Emptied container retains vapor and product residue. Observe all labeled safeguards until container is discarded.

Securely wrap empty container in several layers of newspaper and discard in trash.

SPILL OR LEAK:

In case of spill or leak, soak up with paper towels and discard in trash.

For more product information or questions, call toll-free 1-800-225-2883, 8:00AM - 4:00PM Pacific Standard Time. MONDAY-FRIDAY

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO THE DIRECTIONS IN THE USE DIRECTIONS TABLE AND OBSERVE ALL PRECAUTIONS.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION HARMFUL IF SWALLOWED, MAY CAUSE EYE IRRITATION. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing.

IF SWALLOWED, this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

NOTICE: BUYER ASSUMES ALL RESPONSIBILITY FOR SAFETY AND USE NOT IN ACCORDANCE WITH DIRECTIONS. IF THESE TERMS ARE NOT ACCEPTABLE, RETURN AT ONCE UNOPENED.

Environmental Hazards

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

PRECAUTION: SEE "DIRECTIONS FOR USE" FOR IMPORTANT CONTAINER PRECAUTIONS, and inside label for storage and disposal information.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

REFORMULATION OR REPACKAGING FOR SALE IS PROHIBITED.

©Trademark of Monsanto Company ©MONSANTO COMPANY 1993

MONSANTO COMPANY AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.
EPA Reg. No. 524-370

EPA Est. (G)56100-LA-001; (L)524-LA-1; (I)239-IA-3
Lot number prefix G, L, or I indicates appropriate establishment number.

897.18-100.11-3X3

COMPADRE

HERBICIDE BY **Monsanto**

COMPLETE DIRECTIONS FOR USE AS A CUT STUMP AND TREE INJECTION TREATMENT IN NONCROP AREAS SUCH AS FORESTRY, UTILITY RIGHTS-OF-WAY, ROADSIDES AND SIMILAR INDUSTRIAL OR NONCROP AREAS.

AVOID CONTACT WITH FOLIAGE, GREEN STEMS, EXPOSED NONWOODY ROOTS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

HARMFUL IF SWALLOWED.

MAY CAUSE EYE IRRITATION.

Avoid contact with skin, eyes or clothing.

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water. Get medical attention if irritation persists. Remove product from skin or clothing.

IF SWALLOWED, this product will cause gastrointestinal tract irritation. Immediately dilute by swallowing water or milk. Get medical attention.

In case of an emergency involving this product, Call Collect, day or night, (314) 694-4000.

Environmental Hazards

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers.

DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine, in the form of the isopropylamine salt..... 18.0%

INERT INGREDIENTS..... 82.0%
100.0%

*Contains 193 grams per litre or 1.61 pounds per U.S. gallon of the active ingredient, glyphosate, N-(phosphonomethyl) glycine, in the form of the isopropylamine salt. Equivalent to 143 grams per litre or 1.19 pounds per U.S. gallon of the acid, glyphosate.

This product is protected by U.S. Patent No. 4,405,531. Other patents pending. No license granted under any non-U.S. patent(s).

(Continued on back panel)

©MONSANTO COMPANY 1993

*Registered trademark of Monsanto Company.



MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 USA

EPA Reg. No. 524-37C

Part 1 of 2

EPA Est. 524-LA-1

897.12-100.09/53L

NET 2.5 GAL

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: STORE ABOVE 10°F (-12°C) TO KEEP PRODUCT FROM CRYSTALLIZING.

Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68° F (20° C) for several days to redissolve and shake or roll to mix well before using.

DISPOSAL: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

It is a violation of Federal law to use this product in any manner inconsistent with the labeling.

This label should be in the possession of the user at the time of the herbicide application.

NOTE: When treating trees originating from root suckers, or where root grafting has occurred, product may move through the roots to damage untreated adjacent trees.

CUT STUMP APPLICATIONS

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. This product should not be diluted with water. Apply this ready-to-use product to the freshly cut surface **IMMEDIATELY AFTER CUTTING TREES.**

Cut vegetation close to the soil surface.

DELAY IN APPLYING THIS PRODUCT MAY RESULT IN REDUCED PERFORMANCE.

AVOID WASH-OFF OF PRODUCT FROM TREATED STUMPS BY RAINFALL OR OTHER MEANS WITHIN 6 HOURS OF APPLICATION.

For best results, trees should be cut during periods of active growth and full leaf expansion.

TREE INJECTION APPLICATIONS

Apply the equivalent of 2 ml of this product for each 2 inches of trunk diameter. This is best achieved by applying this product either to a continuous frill around the tree or to cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying to a continuous frill or more closely spaced cuttings.

Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frilling or cutting. In species such as this, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted product. For best results, applications should be made during periods of active growth and after full leaf expansion.

WOODY BRUSH AND TREES CONTROLLED

When applied as directed under the conditions described, this product will control or partially control most woody brush and tree species, some of which are listed below.

- | | |
|-------------------------------|--------------------------------|
| Alder | Oak |
| <i>Ainus spp.</i> | <i>Quercus spp.</i> |
| Coyotebrush | Poplar |
| <i>Baccharis consanguinea</i> | <i>Populus spp.</i> |
| Dogwood | Saltcedar |
| <i>Cornus spp.</i> | <i>Tamarix spp.</i> |
| Eucalyptus | Sweetgum |
| <i>Eucalyptus spp.</i> | <i>Liquidambar styraciflua</i> |
| Hickory | Sycamore |
| <i>Carya spp.</i> | <i>Platanus occidentalis</i> |
| Madrone | Tan oak |
| <i>Arbutus menziesii</i> | <i>Lithocarpus densiflorus</i> |
| Maple | Willow |
| <i>Acer spp.</i> | <i>Salix spp.</i> |

WETLAND SITES

Do not apply this product directly to water.

This product may be applied as directed around water and in wetland areas, including land adjacent to and surrounding domestic water supply reservoirs, supply streams, lakes and ponds.

Consult local public water control authorities before applying this product around public water. Permits may be required to treat in such areas.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. **NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE.** This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company including, but not limited to incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE, SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this **LIMIT OF WARRANTY AND LIABILITY** which may not be varied by any verbal or written agreement.

