

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR BUCKLE® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label for Buckle herbicide and this supplement.

BUCKLE®
Herbicide by Monsanto

EPA Reg. No. 524-375

FOR DISTRIBUTION AND USE ONLY WITHIN THE FOLLOWING NORTH DAKOTA COUNTIES: BURLEIGH, EMMONS, GRANT, KIDDER, MCLEAN, MERCER, MORTON, OLIVER, SHERIDAN AND SIOUX.

GRANULAR SELECTIVE HERBICIDE FOR FALL PREPLANT APPLICATION TO CONTROL WILD OATS AND PIGEONGRASS IN SPRING WHEAT.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, (314) 694-4000.

®Buckle is a registered trademark of Monsanto Company.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

This label must be in the possession of the user at the time of herbicide application.

Apply Buckle at a rate of 10 pounds per acre in the fall prior to spring planting of Spring Wheat.

Incorporate fall Buckle applications within 24 hours with a field cultivator. For the best results a second incorporation should be done in the spring at a right angle with a field cultivator with harrow attachment.

Buckle applications in the fall should only be made to fields fallowed the current calendar year.

CONTINUED

NOTIFICATION
LABEL NOT REVIEWED
PER PR NOTICE 88-6

DATE. OCT 31 1990

SUPPLEMENTAL LABELING
Buckle® Herbicide by Monsanto
Page 2

Do not fall apply Buckle for spring wheat on soil that has been treated with trifluralin during that calendar year.

Spring wheat varieties identified as susceptible to early injury from preplant applications of Buckle include: Alex, Amidon, Ellar, Era, Erik, Lew and Newana.

To minimize stress and allow for uniform germination, a seeding depth of no more than 1.5 to 2 inches is recommended for semi-dwarf spring wheat varieties.

DO NOT APPLY TO HILLTOPS.

Seed treatments are recommended to reduce potential stress to germinating seedlings from disease or insects.

Do not use on fields if the field will be irrigated during the spring wheat growing season.

NOTE: Slight stand reductions do not normally affect yield.

DO NOT APPLY BUCKLE TO A STRIP OR AREA IN THE FIELD THAT WILL SERVE AS AN UNTREATED COMPARISON FOR WEED CONTROL AND CROP TOLERANCE.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label for Buckle before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

®Buckle is a registered trademark of Monsanto Company.

© Monsanto Company 1990

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167

