

PM 25

J 14 - 375

11/17/97

1/2

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR BUCKLE® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label for Buckle herbicide and this supplement.

ACCEPTED

NOV 17 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-375



BUCKLE®

GRANULAR HERBICIDE BY **Monsanto**

EPA Reg. No. 524-375

®Buckle is a registered trademark of Monsanto Company.

FOR USE WITHIN NORTH DAKOTA EXCLUDING THE FOLLOWING COUNTIES: CASS, GRAND FORKS, PEMBINA, RICHLAND, TRAILL, AND WALSH.

THIS PRODUCT MAY ALSO BE USED IN THE FOLLOWING SOUTH DAKOTA COUNTIES: CORSON, DEWEY, HARDING, PERKINS, AND ZIEBACH.

GRANULAR SELECTIVE HERBICIDE FOR SPRING PREPLANT APPLICATION TO CONTROL WILD OATS AND PIGEONGRASS IN SPRING WHEAT.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with the labeling.

This label must be in possession of the user at the time of herbicide application.

Apply Buckle® at a rate of 10 pounds per acre in the spring prior to planting Spring Wheat.

Incorporate spring Buckle applications within 24 hours with a field cultivator with harrow attachment. A second incorporation at a right angle to the first will provide the best results. Delay the second incorporation or spring wheat planting for a minimum of 5 days following the first incorporation. Do not set tillage equipment to incorporate soil deeper than the first incorporation.

Do not apply Buckle for spring wheat on soil that has been treated with 0.5 lbs a/acre or greater of trifluralin, or other dinitroaniline herbicide, the previous year.

NOTE: Spring wheat varieties identified as susceptible to early injury from preplant applications of Buckle include: Alex, Amidon, Butte 86, Ellar, Era, Erik, Lew and Newana.

To minimize stress and allow for uniform germination, a seeding depth of no more than 1.5 to 2 inches is recommended. Delay spring planting of spring wheat until the average soil temperature at the 2-inch depth is 45 degrees F or greater.

DO NOT APPLY BUCKLE TO HILLTOPS.

Seed treatments are recommended to reduce potential stress to germinating seedlings from disease or insects.

Do not use on fields if the field is or will be irrigated in the current growing season.

NOTE: Slight stand reductions do not normally affect yield.

DO NOT APPLY BUCKLE TO A STRIP OR AREA IN THE FIELD THAT WILL SERVE AS AN UNTREATED COMPARISON FOR WEED CONTROL AND CROP TOLERANCE.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label for Buckle herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

© Monsanto Company 1997

Monsanto Company
St. Louis, Missouri 63167

9/25/97

29003V1-5

PM 25

524-375


11/17/97

2/2

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL FOR BUCKLE® HERBICIDE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label for Buckle herbicide and this supplement.



BUCKLE®

GRANULAR HERBICIDE BY **Monsanto**

EPA Reg. No. 524-375

®Buckle is a registered trademark of Monsanto Company.

FOR USE ONLY WITHIN THE FOLLOWING COUNTIES IN NORTH AND SOUTH DAKOTA:

North Dakota:

Adams, Billings, Bottineau, Bowman, Burleigh, Burke, Divide, Dunn, Emmons, Grant, Golden Valley, Hettinger, Kidder, McHenry, McKenzie, McLean, Mercer, Morton, Mountrail, Oliver, Renville, Sheridan, Sioux, Slope, Stark, Ward, and Williams.

South Dakota:

Corson, Dewey, Harding, Perkins, and Ziebach.

GRANULAR SELECTIVE HERBICIDE FOR FALL PREPLANT APPLICATION TO CONTROL WILD OATS AND PIGEONGRASS IN SPRING WHEAT.

Keep out of reach of children.

CAUTION!

In case of emergency involving this product, Call Collect, day or night, (314) 694-4000.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the labeling.

This label must be in possession of the user at the time of herbicide application.

Apply Buckle® at a rate of 10 pounds per acre in the fall prior to spring planting of Spring Wheat. Average soil temperature at the 2-inch depth must be 40 degrees F or less.

Incorporate fall Buckle applications within 24 hours with a field cultivator. For the best results a second incorporation should be done in the spring at a right angle with a field cultivator with harrow attachment. If no spring tillage is anticipated, both incorporations should be done in the fall with a minimum of 5 days between first and second incorporations. For best results, care must be

taken to avoid working the soil deeper than the first tillage operation.

Do not fall-apply Buckle for spring wheat on soil that has been treated with 0.5 lbs ai/acre or greater of trifluralin, or other dinitroaniline herbicide, during that calendar year.

NOTE: Spring wheat varieties identified as susceptible to early injury from preplant applications of Buckle include: Alex, Amidon, Butte 86, Ellar, Era, Erik, Lew and Newana.

To minimize stress and allow for uniform germination, a seeding depth of no more than 1.5 to 2 inches is recommended. Delaying spring planting until soil temperatures have reached 45 degrees F at the two inch depth will reduce the potential for crop injury.

DO NOT APPLY TO HILLTOPS.

Seed treatments are recommended to reduce potential stress to germinating seedlings from disease or insects.

Do not use on fields if the field will be irrigated during the spring wheat growing season.

Note: Slight stand reductions do not normally affect yield.

DO NOT APPLY BUCKLE TO A STRIP OR AREA IN THE FIELD THAT WILL SERVE AS AN UNTREATED COMPARISON FOR WEED CONTROL AND CROP TOLERANCE.

Read the "LIMIT OF WARRANTY AND LIABILITY" in the label for Buckle herbicide before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

© Monsanto Company 1997

MONSANTO COMPANY
ST. LOUIS, MISSOURI 63167

9/25/97

29003V1-4