




For control or suppression of emerged weeds in fallow and reduced tillage small grain systems and CRP or set-aside acres.

Complete Directions for Use

EPA Reg No 524-376

AVOID CONTACT WITH FOLIAGE OF CROP OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

*Landmaster is a registered trademark of Monsanto Company.



listed on the atrazine and Bladex labels

*Bladex is a trademark of E. I. du Pont de Nemours and Company

ANNUAL WEEDS CONTROLLED

Barnyardgrass	Pigweed, redroot
<i>Echinochloa crus-galli</i>	<i>Amaranthus retroflexus</i>
Brome, downy	Sandbur, field
<i>Bromus tectorum</i>	<i>Cenchrus spp</i>
Foxtail, green	Stinkgrass
<i>Setaria viridis</i>	<i>Eragrostis ciliaris</i>
Kochia	Thistle, Russian
<i>Kochia scoparia</i>	<i>Salsola kali</i>
Lambsquarters	Wheat
<i>Chenopodium album</i>	<i>Triticum aestivum</i>
Lettuce, prickly	Witchgrass
<i>Lactuca serriola</i>	<i>Panicum capillare</i>
Mustard, lamby	
<i>Descurainia pinnata</i>	

Weeds controlled with 40 ounces of this product plus 1 pound or less of the active ingredient, atrazine, per acre include downy brome, green foxtail, stinkgrass and volunteer corn.

Additional weeds controlled with 54 ounces of this product plus 2 pounds or less of the active ingredient, atrazine, per acre or 2.4 to 4 pounds of the active ingredient, cyanazine, in Bladex per acre include kochia, lambsquarters, mustard, pigweed, prickly lettuce, and Russian thistle.

Additional weeds controlled with 64 ounces of this product plus 3 pounds or less of the active ingredient, atrazine, per acre include sandbur, witchgrass, and yellow foxtail. Barnyardgrass will be suppressed at the 64 ounce per acre rate and will require 86 ounces for control.

The addition of 2 percent dry ammonium sulfate by weight is recommended to increase the performance of this product plus atrazine or Bladex tank mixtures.

In Oregon and Washington, do not exceed 1 pound of the active ingredient, atrazine, per acre.

Consult the atrazine and Bladex labels (90 DF 80W 41) for use rates, soil type, planting, cropping and other restrictions, as well as other precautionary statements and use according to the most restrictive label.

These tank mixtures may be applied with ground or aerial equipment. See the "APPLICATION EQUIPMENT" section for instructions.

POSTHARVEST APPLICATIONS

This product will provide control of weeds following grain harvest. Weeds should be allowed to regrow after damage incurred during harvest operations and to recover from environmental stress before application of this product. Weeds should be treated prior to the heading stage of annual grasses and before broadleaf weeds exceed 24 inches in height. Ammonium sulfate will improve performance on annual weeds under stress conditions.

Weeds controlled at the 40-ounce-per-acre rate include downy brome, green foxtail, stinkgrass, volunteer wheat.

Weeds controlled at the 54-ounce-per-acre rate include kochia, lambsquarters, mustard, pigweed and Russian thistle.

Weeds controlled at the 64-ounce-per-acre rate include barnyardgrass, sandbur, witchgrass, yellow foxtail and prickly lettuce.

INSERT "ECOFARMING SYSTEMS" SECTION.

INSERT THE FOLLOWING SECTION:

ECOFARMING SYSTEMS

The Ecofarming System consists of the following rotation:
winter wheat, corn/sorghum, ecofallow.

Use the following tank mixtures for control of emerged annual weeds before planting corn or sorghum in the Ecofarming System.

- LANDMASTER ® II at 54 to 64 ounces per acre
plus
- ATRAZINE at 0.75 to 1 pound a.i. per acre
plus
- LASSO ® EC at 2.5 to 3 quarts per acre

The above tank mixture should be applied in 28-0-0 or 32-0-0 liquid fertilizer carrier at 20 to 30 gallons per acre. The liquid fertilizer may be diluted with water to achieve the proper carrier volume.

WEEDS CONTROLLED - The following weeds will be controlled up to a maximum height of 4 inches:

Brome, downy
Bromus tectorum

Cheat
Bromus secalinus

Foxtail, green
Setaria viridis

Foxtail, yellow
Setaria lutescens

Kochia*
Kochia scoparia

Lettuce, prickly
Lactuca scariola

Pigweed, redroot
Amaranthus retroflexus

Thistle, Russian
Salsola kali

Wheat, volunteer
Triticum aestivum

*For improved control of kochia, add 4 ounces per acre (0.125 pound a.i. per acre) of Banvel to the above tank mixture.

4 9 4

Risk of crop injury from 2,4-D or Banvel can be reduced by applying this treatment 7 to 14 days before planting.

Refer to the Lasso label for preemergence weed control achieved by this tank mixture.

Refer to the specific product labels for crop rotation restrictions and cautionary statements for all products used in these tank mixtures.

Ⓢ Lasso is a registered trademark of Monsanto Company.