

2/8/2

1/3/2

GENERAL INFORMATION

Ranger® is a postemergence herbicide used in agricultural systems for control of quackgrass and other weeds listed on this label.

This product enters the plant through the foliage and moves throughout the plant. Visual effects of control are a gradual wilting or yellowing of the plant, which advances to complete browning of above-ground growth and deterioration of affected underground plant parts. Visible symptoms will usually develop on labeled weeds within 2 to 4 days after application, but may not occur for 7 or more days. Extremely cool or cloudy weather following treatment may slow activity of this product and delay the visual effects of control.

This product does not provide residual weed control. For subsequent residual weed control, follow a label approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the label of all herbicides used.

APPLICATION PRECAUTIONS AND WARNINGS

- Apply only when there is no hazard from spray drift since very small quantities of spray which may not be visible may severely injure susceptible crops or desirable vegetation.
- The likelihood of injury according to adjacent crops from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions including lesser wind velocities will favor spray drift.
- Do not feed or forage vegetation from treated areas within 8 weeks after application.
- Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product with herbicides or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops or other unintended consequences.

Timing of Application

Preventative spray should be applied postemergence to actively growing weeds when they have reached the 3- to 4-leaf stage and are at least 6 to 8 inches high. For quackgrass control, apply 3 pints per acre of this label. For other weeds, refer to the specific product label.

For control of quackgrass and other annual weeds, apply 3 pints per acre of this product when the plants have reached the 3- to 4-leaf stage and are at least 6 to 8 inches high. For quackgrass control, apply 3 pints per acre of this product when the plants have reached the 3- to 4-leaf stage and are at least 6 to 8 inches high. For other weeds, refer to the specific product label.

WEEDS CONTROLLED

FOR CONTROL OF QUACKGRASS AND ASSOCIATED ANNUAL WEEDS, APPLY 3 PINTS PER ACRE OF THIS

PRODUCT IN RECOMMENDED WATER VOLUMES WHEN THE PLANTS HAVE REACHED THE 3- TO 4-LEAF STAGE AND ARE AT LEAST 6 TO 8 INCHES HIGH. FOR QUACKGRASS CONTROL IN SODS WHICH HAVE BEEN ESTABLISHED FOR 3 YEARS OR LONGER, USE 5 TO 6 PINTS PER ACRE. THIS PRODUCT MAY BE TANK-MIXED WITH 2,4-D OR BANYEL FOR FORAGE LEGUME CONTROL IN SOD SITUATIONS. REFER TO THE SPECIFIC PRODUCT LABELS FOR USE RATES, CROP ROTATION RESTRICTIONS AND CAUTIONARY STATEMENTS.

When applied as directed under the conditions described, this product controls the following annual and perennial weeds listed below:

RECOMMENDED RATES AND WEEDS CONTROLLED

ANNUAL WEED SPECIES	MAXIMUM HEIGHT	RATE PER ACRE
Festail <i>Setaria</i> spp.	12"	12 oz
Barryardgrass <i>Echinochloa crus-galli</i>	6"	18 oz
Bluegrass, annual <i>Poa annua</i>		
Brome, downy* <i>Bromus tectorum</i>		
Mustard, blue <i>Chorispora tenella</i>		
Mustard, taney <i>Descurainia pinnata</i>		
Mustard, tumbled <i>Sisymbrium albidum</i>		
Mustard, wild <i>Sinapis arvensis</i>		
Spurry, umbelliform <i>Holcus umbelliformis</i>		
Barley <i>Hordeum vulgare</i>	12"	
Rye <i>Secale cereale</i>		
Sandspur, field <i>Callipepla</i> spp.		
Shattercane <i>Sorghum bicolor</i>		
Stinkgrass <i>Tripsacum daniellii</i>		
Wheat <i>Triticum aestivum</i>		
Mint <i>Mentha</i> spp.		
Bluegrass, bulbous <i>Lolium perenne</i>		
Chick <i>Gallinula</i> spp.		
Chickweed, mouseear <i>Malva</i> spp.		
Corn, volunteer <i>Zea mays</i>		
Groundsel, common <i>Helianthus annuus</i>		
Moroccan Bladder <i>Plantago lanceolata</i>		

BEST AVAILABLE COPY

ANNUAL WEED SPECIES	MAXIMUM HEIGHT	RATE PER ACRE
Pennycress, field <i>Thlaspi arvense</i>	6"	24 oz
Rocket, London <i>Sisymbrium irio</i>		
Ryegrass, Italian <i>Lolium multiflorum</i>		
Shepherdspurs <i>Caposela bursa-pastoris</i>		
Buttercup <i>Ranunculus</i> spp.	12"	
Cocklebur <i>Xanthium pensylvanicum</i>		
Crabgrass <i>Digitaria</i> spp.		
Dwarfandelion <i>Rigida cespitosa</i>		
Falsesun, smallseed <i>Camelina microcalpa</i>		
Lambquarters, common <i>Chenopodium album</i>		
Oats, wild <i>Avena fatua</i>		
Panicum, tall <i>Panicum dichotomiflorum</i>		
Panicum, Texas <i>Panicum texanum</i>		
Pigweed, redroot <i>Amaranthus retrofractus</i>		
Pigweed, smooth <i>Amaranthus hybridus</i>		
Witchgrass <i>Panicum capillare</i>		

*For control in no till systems, use 24 fluid ounces per acre.

PERENNIAL WEED SPECIES	GROWTH STAGE	RATE PER ACRE, PINTS
Fescue (tall) <i>Festuca arundinacea</i>	Fall applications	4 to 12"
	Spring applications	4 to 12"

Mully, crested <i>Muhlenbergia frondosa</i>	8 to 12"	3 to 5
Orchardgrass <i>Dactylis glomerata</i>	8 to 12"	3 to 5
Quackgrass <i>Alopecurus repens</i>	8 to 12"	3 to 5

APPLICATION EQUIPMENT

Apply this product with the following equipment: broadcast sprayer, backpack sprayer, or other equipment. For optimal results, use a spray nozzle which produces a fine spray. The spray should be applied during the growing period of the weeds. Do not apply this product to dormant weeds.

during both growing and dormant periods, since even small quantities of spray can cause severe plant injury.

Ground Application: Apply recommended rates of this product in 5 to 30 gallons of water per acre as a broadcast spray. For optimum spray distribution and coverage, use flat fan or low-volume flood nozzles. When using flood nozzles, space them no more than 40 inches apart and ensure double overlap of spray pattern. Refer to the manufacturer's recommendations for correct pressure and nozzle height above the target canopy. Avoid pressure and nozzle combinations which produce fine droplets or mist susceptible to drift.

Use appropriate marking devices to ensure uniform spray coverage and best results from this product.

Aerial Application: Apply the recommended rates of this product in 3 to 15 gallons of water per acre as a broadcast spray. DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR WHEN OTHER CONDITIONS WILL FAVOR DRIFT. DRIFT MAY CAUSE DAMAGE TO ANY VEGETATION CONTACTED TO WHICH TREATMENT WAS NOT INTENDED TO PREVENT INJURY TO ADJACENT DESIRABLE VEGETATION. APPROPRIATE BUFFER ZONES MUST BE MAINTAINED.

Coarse sprays are less likely to drift, therefore do not use nozzles or nozzle configurations which disperse spray as fine spray droplets. Do not angle nozzles forward into airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used when making aerial applications. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

MIXING INSTRUCTIONS

Fill the spray tank to about 3/4 of the desired volume with clean water. Add the recommended amount of this product, then complete the mixing process while maintaining agitation. Remove the hose from the mix tank immediately after filling to avoid siphoning back into the carrier source. During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, terminate by pass and return lines at the tank bottom and/or use an agriculturally approved anti-foam or defoaming agent.

Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Do not add additional surfactant to the spray solution when using this product.

NOTE: Reduced results may occur if water containing soil is used, such as water from ponds, streams, or ditches.

When mixing in water below 10°F with this product, this product may be used with the following water temperature range: 10°F to 60°F.

SPRAYER CLEANUP

After use, clean the sprayer thoroughly with water. Rinse the tank, hoses, and nozzles with water. Do not use the sprayer for other purposes until it has been thoroughly cleaned. Do not use the sprayer for other purposes until it has been thoroughly cleaned.



RANGER
Herbicide
For control of quackgrass and listed weeds in annual cropping systems, pastures and sods.


AVOID CONTACT WITH FOLIAGE OF CROP OR OTHER DESIRABLE VEGETATION, SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Complete Directions for Use

EPA Reg. No. 104-382

Ranger is a registered trademark of Monsanto Company.

1394-1 892-20 006/45/53



When mixing in water below 10°F with this product, this product may be used with the following water temperature range: 10°F to 60°F.

RANGER



Herbicide

For control of quackgrass and listed weeds in annual cropping systems, pastures and sods.

DO NOT CONTACT WITH FEEDABLE LIVESTOCK OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

Complete Directions for Use

EPA Reg. No. 504-382

*Ranger is a registered trademark of Monsanto Company.

1994 | 892 29 000 65-53

MONSANTO COMPANY 1994

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
500 NORTH MISSOURI AVENUE, ST. LOUIS, MO 63103

Read the entire label before using this product.

Use only according to label instructions.

Read LIMIT OF WARRANTY AND LIABILITY before using this product. If forms are not available, contact your distributor.

REFORMULATION IS PROHIBITED. SEE THE ORIGINAL CONTAINER LABEL FOR REPAKAGING INSTRUCTIONS.

LIMIT OF WARRANTY AND LIABILITY

Buyer warrants that the product is used in accordance with the label instructions. The Company does not warrant the product for any purpose not set forth in the label. The Company does not warrant the product when used in accordance with the label instructions for purposes not set forth in the label. The Company does not warrant the product when used in accordance with the label instructions for purposes not set forth in the label. **NO OTHER LIMITED WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE.** This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims, whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, application to or contact with desirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE DISCRETION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

Hazards to Humans
and Domestic Animals

Keep out of reach of children.

DANGER! PELIGRO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

CAUSES EYE BURNS

HARMFUL IF SWALLOWED OR INHALED

Irritates eyes, nose, skin or on clothing.

Avoid breathing vapors or spray mist.

FIRST AID - IF IN EYES: Immediately flush with plenty of water for at least 15 minutes. Get medical attention.

IF INHALED: Remove individual to fresh air. Get medical attention if breathing difficulty develops.

IF SWALLOWED: The product will cause gastric irritation. Do not induce vomiting. Drink plenty of water. Do not take any water while feeling ill.

Personal Protective Equipment

Applicators and other handlers (other than mixers and loaders) must wear: long-sleeved shirt and long pants, shoes plus socks, and protective eyewear. **Mixers and loaders must wear:** long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendation

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

In case of an emergency involving this product, call Collect day or night (314) 694-4000.

Environmental Hazards

Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

ACTIVE INGREDIENT

*Glyphosate N (phosphonomethyl) glycine
in the form of its isopropylamine salt 28.6%
INERT INGREDIENTS 71.4%
100.0%

*Contains 374 grams per liter of glyphosate in the form of its isopropylamine salt. Equivalent to 240 grams per liter of 2 pound per gallon of the acid phosphate.

Product protected by U.S. Pat. 4,711,411. Other patents pending. No known acute toxicity to humans.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will cause injury to other persons, either directly or through drift, or to valuable handlers may be in the area during application.

For any requirements specific to your State or to consult the agency responsible for pesticide registrations.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, waterproof gloves, shoes plus socks and protective eyewear.

For more product information call toll free 1-800-332-3111.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE

At temperatures below 50 F, the viscosity of this product may increase to a point that pouring and mixing could be difficult.

DISPOSAL

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Empty container retains vapor and product residue. Observe all labeled safeguards until container is destroyed.

Do not reuse container. Triple rinse container, then puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state or local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY