


FALLOW MASTER LABEL CHANGES AND ADDITIONS

1. Add:
corn
2. Add:
This product is recommended for control of emerged weeds prior to establishment of labeled crops. Large amounts of green or decaying vegetation left standing or incorporated into the seedbed may enhance the development of disease in newly planted crops. This may result in poor emergence and/or stands, especially under cool and/or wet conditions.
3. Add:
Spraying early to control young weeds before dense stands develop or light cultivation to assist weed decay will favor preparation of suitable seedbeds.
4. Add:
Lambsquarters
Chenopodium album
5. Add:
Pigweed
Amaranthus spp.
6. Add:
Stinkgrass
Eragrostis cilianensis
7. Add:
Witchgrass
Panicum capillare
8. Add:
Sandbur, field
Cenchrus spp.
9. Change Oat to Oats, wild
10. Change 42 oz. to 44 oz.
21 oz. to 22 oz.



11. Add:

FALLOW MASTER PLUS GLEAN™ TANK MIXTURE
(Fallow Applications)

This tank mixture provides initial postemergence control and extended soil residual control of labeled annual weeds in reduced tillage fallow areas to be planted only to wheat. Applications of this tank mixture should be made in the fall following grain harvest or in the early spring to summer fallow acres. Use the labeled rate of Glean with the recommended rate of this product and mix as directed in the "MIXING INSTRUCTIONS". Refer to the Glean label for mixing instructions, proper stages, rates, planting, cropping and other restrictions. Follow all precautionary statements on this and the Glean label.

12. Add:

FALLOW MASTER plus ALLY™ TANK MIXTURE
(Fallow Applications)

This tank mixture provides initial postemergence control of labeled annual weeds in reduced tillage fallow areas. Depending on environmental conditions, Ally may provide short-term residual control of certain annual broadleaf weed species. Applications of this tank mixture should be made following grain harvest or in the early spring-to-summer fallow acres. Use the labeled rate of Ally with the recommended rate of this product and mix as directed in the "MIXING INSTRUCTIONS" section. Refer to the Ally label for mixing instructions, proper stages, rates, planting, cropping and other restrictions. Follow all precautionary statements on this and the Ally label.

Ally and Glean are trademark of E. I. duPont de Nemours and Company

13. Add:

AID TO TILLAGE

This product used in combination with preplant and conventional fallow tillage practices will provide control of downy brome, cheat, volunteer wheat, tansy mustard and foxtails.

Apply 22 fluid ounces of this product in 3 to 10 gallons of water per acre to weeds that are actively growing. Treat when weeds are less than 6 inches in height. Application must be followed by conventional tillage practices before regrowth of the treated plant occurs. Allow at least 1 day after application before tillage.

14. Add:

POSTHARVEST APPLICATIONS

This product will provide control of weeds following grain harvest. Weeds should be allowed to regrow after damage incurred during harvesting operations, and to recover from environmental stress, before application of this product. Weeds should be treated prior to the heading stage of annual grasses and before broadleaf weeds exceed 24 inches in height. Ammonium sulfate will improve performance on annual weeds under stress conditions.

Weeds controlled at the 32 ounce-per-acre rate include: downy brome, green foxtail, stinkgrass and volunteer wheat.

Weeds controlled at the 44 ounce-per-acre rate include kochia, lambsquarters, mustard, pigweed and Russian thistle.

Weeds controlled at the 52 ounce per acre rate include: barnyardgrass, sandbur, witchgrass, yellow foxtail, and prickly lettuce.

15. UNDER MIXING INSTRUCTIONS SECTION, CHANGE SURFACTANT STATEMENT TO READ:

Additional surfactant is not recommended for this formulation.

16. Add:

FALLOW MASTER plus ATRAZINE or BLADEX™.

For use only in Colorado, Kansas, Nebraska, Oklahoma, Oregon, South Dakota and Washington.

16. Add: (cont.)

Tank mixtures of Fallow Master plus atrazine or Bladex will provide postemergence control of listed annual weeds in fallow and reduced tillage systems. In addition, these tank mixtures will provide soil residual control of weeds listed on the atrazine and Bladex labels.

ANNUAL WEEDS CONTROLLED

Barnyardgrass Echinochloa crus-galli	Pigweed, redroot Amaranthus retroflexus
Brome, downy Bromus tectorum	Sandbur, field Cenchrus spp.
Foxtail, green Setaria viridis	Stinkgrass Eragrostis cilianensis
Kochia Kochia scoparia	Thistle, Russian Salsola kali
Lambquarters Chenopodium album	Wheat Triticum aestivum
Lettuce, prickly Lactuca serriola	Witchgrass Panicum capillare
Mustard, tansy Descurainia pinnata	

Apply 32 ounces of this product plus 1 pound or less of the active ingredient atrazine per acre. Use 44 ounces of this product plus 2 pounds or less of the active ingredient atrazine per acre. Use 52 ounces of this product plus 1 to 3 pounds of the active ingredient atrazine per acre. Use 44 ounces of this product plus 2.4 to 4 pounds of the active ingredient cyanazine in Bladex per acre. In Oregon and Washington, do not exceed 1 pound of the active ingredient atrazine per acre. Barnyardgrass will be suppressed at the 52 ounce per acre rate and will require up to 70 ounces per acre for control. The addition of ammonium sulfate is recommended to increase the performance of Fallow Master herbicide plus atrazine or Bladex tank mixtures.

Consult the atrazine and Bladex labels (90 DF, 80W, 4L) for use rates, soil type, planting, cropping and other restrictions, as well as other precautionary statements and use according to the most restrictive label.

These tank mixtures may be applied with ground or aerial equipment. See the "APPLICATION EQUIPMENT" section for instructions.

For ground application, apply in 3 to 10 gallons of water per acre.

For aerial application, apply in 3 to 5 gallons of water per acre.

17. Delete:

remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Call a physician.

Add:

remove individual to fresh air. Seek medical attention if breathing difficulty develops.

FALLOW MASTER



For control or suppression of emerged weeds in fallow and reduced tillage small grain systems in the states of Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington, and Wyoming.

Complete Directions for Use
AVOID CONTACT WITH FOLIAGE OF CROP OR OTHER DESIRABLE VEGETATION SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT

EPH Reg. No. 526 990

*Fallow Master is a trademark of and used under license from Syntex, Ltd.

1987 1 000.05-000 01 /52

BEST AVAILABLE COPY

MONSANTO COMPANY 1987

In case of an emergency involving this product
 Call Collect, day or night (313) 694-6000

MONSANTO COMPANY
 800 NORTH MAIN STREET
 ST. LOUIS, MISSOURI 63101 U.S.A.

Read the entire label before using this product

Use only according to label instructions

Read **LIMIT OF WARRANTY AND LIABILITY** before buying or using. It forms an integral part of the contract and is not acceptable return of once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is, reasonable in for the purposes set forth in the Complete Directions for Use label headed (Directions) when used in accordance with those Directions under the conditions described therein. **NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSES OR MERCHANTABILITY IS MADE.** This warranty is the subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims, whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those listed in the Directions, application to or contact with undesirable vegetation, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. WE DO NOT EVEN BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

Hazards to Humans
 and Domestic Animals

Keep out of reach of children

DANGER!

CAUSES EYE BURNS

HARMFUL IF SWALLOWED OR INHALED
 MAY CAUSE ALLERGIC SKIN REACTION

Do not get in eyes, on skin or on clothing

Wear goggles or face shield and rubber gloves in transferring and mixing and when adjusting, repairing or cleaning equipment.

Avoid breathing vapors or spray mist

Wash thoroughly with soap and water after handling
FIRST AID: IF IN EYES: Immediately flush with plenty of water for at least 15 minutes. Call a physician.

IF ON SKIN: Immediately flush with plenty of water while removing contaminated clothing. As soon as possible, wash skin thoroughly with soap and water. If irritation persists, call a physician. Avoid clothing before reuse. **Do not get on face, neck, or hands.**

IF SWALLOWED: Do not induce vomiting. If not breathing, give artificial respiration. Preferable route is mouth-to-mouth. Call a physician.

IF INHALED: This product will cause permanent lung irritation. Immediately discontinue breathing vapors or mist. Call a physician. **NEVER USE BURNING OILS TO CLEAN EQUIPMENT.**

In case of an emergency involving this product

Call Collect, day or night (313) 694-6000

Environmental Hazards

Do not apply directly to water or to fish, shellfish, birds, mammals or pollinators. Do not contaminate water by cleaning of equipment or disposal of wastes.

Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied only in equipment that is clean, compatible plastic and glass based and vented.

DO NOT USE STEEL OR ALUMINUM THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN CARBONIZED STEEL OR UNFINISHED STEEL (OR STAINLESS STEEL CONTAINERS OR SPRAY TUBES. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode causing serious personal injury if ignited by open flame, spark, static electricity, or other ignition source.

- Do not use more concentrate to apply the product when covering crops and plants are growing at the vicinity of way to be treated
- Applications should be made only when there is no hazard from spray drift onto very small quantities of spray, which may not be visible. They severely injure susceptible crops or desirable vegetation
- The likelihood of injury occurring to adjacent crops from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions including heavy wind velocities will favor spray drift
- Movement of the product on the particles during windstorms may cause damage to susceptible plants that are contacted. This hazard is reduced if contact occurs shortly after application
- Do not use all crops recommended for all tests or through an obstruction such as the use of handling of the forms of this herbicide or other materials that are not expressly recommended in this labeling. Mixing this product with herbicides or other materials not recommended on this label may result in reduced performance. Use of the product in any manner not described with this label may result in injury to crops, animals or crops, or other undesirable consequences
- Fallow Master is subject to all state and county regulations for dicamba

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Do not store near foodstuffs, feeds, stockstuffs or fertilizer.

STORAGE
STORING ABOVE 70°F in some product in solution in crystalline form, plus in a month when (77°F) after the product to reach room temperature and will or should periodically until crystals have dissolved. Keep containers closed to prevent spills and contamination.

For bulk containers, use the container label for alternate storage information.

DISPOSAL

Wastes resulting from the use of this product that can not be used in chemically regenerated should be disposed of in a landfill approved for pesticide disposal or burned in a safe place away from water supplies. All disposal should be in accordance with applicable Federal, state or local regulations.

See the individual container label for disposal information.

TIMING OF APPLICATION

This product should be applied postemergence to vigorously growing weeds when they have reached the recommended rate given in the "RECOMMENDED RATES AND WEEDS CONTROLLED" section of this label. A solution should be diluted until maximum emergence of the target weeds, but before weeds exceed the maximum rate recommended for annual weeds, after 1 day after treatment before tillage.

Reduced control may result if applications are made too early into growing conditions such as drought stress, disease or insect damage or if weeds have been mowed, grazed or cut. Heavy dust or foliage on an ornamental canopy covering targeted weeds may also reduce control. Residual or regrowth occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or crop loss within 7 hours after application may wash the product off the foliage and a repeat application may be required.

RECOMMENDED RATES AND WEEDS CONTROLLED

For best results, apply this product after most weed seeds have germinated but before seedhead formation in grasses or before bud formation in broadleaves.

When applied as directed, Fallow Master herbicide will provide control or suppression of the grasses and broadleaf weed species listed below. Rates recommended are for maximum weed height at treatment time.

WEED SPECIES	FALLOW MASTER™	
	RATE PLR ACRE	MAXIMUM HEIGHT
10 Annual, grass	21 oz	12"
Setaria weeds		
Barley	32 oz	6"
Hairy wood sorrel		
Beard, dimity		
Dimity suctorial		
Chick*		
Bromus spretus		
Festool		
Setaria spp.		
Ryegrass*		
Alopecurus spretus		
Lactuca scariola		
Lactuca serriola		
11 Annual, broad	32 oz	12"
Dicentra pinnata		
Mustard, hairy		
Scythium affinis		
Blind		
12 Triennial, perennial		
13 Annual/perennial	42 oz	6"
14 Chickweed cross path		
Bullfinch		
Johnson grass stem		
Goat grass		
Agrostis cynosuroides		
Mustard, black		
Chenopodium hybridum		
Panicum, tall		
Panicum dichotomiflorum		
15 1st, 2nd	42 oz	12"
Annual broad		
Beard, dimity		
16 Setaria spp.		

*For control of heavy infestations of these nonannual weeds use 47 ounces.

AMMONIUM SULFATE

The addition of 2 percent dry ammonium sulfate by weight to the liquid concentrate at 17 pounds per 100 gallons of water may increase the performance of Fallow Master on annual weeds. The improvement in performance may be apparent when environmental stress is a concern. Low quality ammonium sulfate may contain material that will not readily dissolve which could result in nozzle tip plugging. To determine quality, you have a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for 1 minute. If undissolved sediment is observed, purchase the ammonium sulfate in water and filter prior to addition to the spray tank. If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may cause water loss. **ENSURE THAT AMMONIUM SULFATE IS COMPLETELY DISSOLVED IN THE SPRAY TANK BEFORE ADDING THIS PRODUCT. THOROUGHLY WASH THE SPRAY SYSTEM WITH CLEAN WATER AFTER USE TO REDUCE CORROSION.**

Note: When using ammonium sulfate, compatibility problems may occur at certain volumes below 5 GPA.

APPLICATION EQUIPMENT

This product may be applied using either ground or aerial spray equipment. Use extreme care to avoid mixing or drifting of herbicide solution onto foliage, grass stems or fruit of desirable crops, trees or plants during both ground and aerial spray since even very small quantities of spray can cause severe plant injury.

GROUND APPLICATION: Apply recommended rates of this product in 3 to 10 gallons of water per acre as a broadcast spray. For optimum spray distribution and coverage, use 40 psi or less volume flow nozzles. When using flow nozzles, space them no more than 40 inches apart and overlap double overlap of spray pattern. Refer to the manufacturer's recommendations for correct pressure and nozzle height above the target canopy. Road proximity and machine effect produce low drifters to avoid the appropriate machine for use.

AERIAL APPLICATION: Apply the recommended rates of this product in 3 to 5 gallons of water per acre as a "road cut" spray. DO NOT APPLY DURING INVERSION CONDITIONS. Wind direction and gusty or uneven ground conditions will favor drift. Drift may cause damage to any vegetation contacted to which treatment was not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

Extreme spray or low drift, therefore, do not use nozzles or other configurations which produce spray as low spray droplets. Do not angle nozzles toward into the airstream and do not increase spray volume by increasing nozzle pressure.

MIXING INSTRUCTIONS

Fill the spray tank to about 1/2 of desired volume with clean water. Add this amount of the product then complete the filling process while maintaining agitation. Before the tank has the desired level immediately after filling to avoid splashing back into the canvas source. During mixing and application, turning of the spray solution may occur. To prevent or minimize foam formation by foam and return loss at the tank bottom, use an agriculturally approved agitator or defoaming agent.

Additional surfactant is not necessary for this formulation. **NOTE:** Reduced results may occur if water containing surfactant is used such as water from ponds and untreated ditches.

SPRAYER CLEANUP

The steps listed below are suggested for thorough cleaning of spray equipment following applications of Fallow Master herbicide. Failure to clean the sprayer thoroughly may result in injury to desirable vegetation subsequently sprayed with the equipment.

1. Run down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank full with water. Flush by operating the sprayer until the solution is purged of the entire water.

2. Fill the tank with water while adding 1 quart of household ammonia for every 75 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.

3. Flush the solution out of the spray tank through the boom.

4. Reverse the nozzles and screens and flush the system with two full tanks of water.

Thoroughly wash parts off, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNPAINTED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PARTS. LANDING GEAR AND MOST SPRAY DEVICES.** The maintenance of an organic coating (Paint) which meets ammonia specification (MIL-C-30413) may prevent corrosion.

006 05 000 01 53

MONSANTO COMPANY 1987

In case of an emergency, washing this product. Call Contact: day or night (314) 650-4000

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI 63167 U.S.A.



For control or suppression of emerged weeds in fallow and reduced tillage small grain systems in the states of Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, North Dakota, Oklahoma, Oregon, South Dakota, Texas, U.S. Washington, and Wyoming.

Complete Directions for Use

Avoid Contact with Foliage of Crop or Other Desirable Vegetation since Severe Injury or Destruction may Result

(Fallow Master is a trademark of and used under license from Sandoz Ltd)

Fallow Master is a trademark of and used under license from Sandoz Ltd

1987 | 006 05 000 01 53

BEST AVAILABLE COPY

L2067