

1760

[illegible]

corn.

1389-2

00017

2760

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

CAUTION!

MAY CAUSE EYE IRRITATION

MAY CAUSE ALLERGIC SKIN REACTION.

Avoid contact with eyes, skin and clothing.

Wear goggles or face shield, rubber gloves, long trousers, long sleeved shirt or jacket of tightly woven material along with boots high enough to cover ankles when transferring and mixing, and when adjusting, repairing or cleaning equipment. Clothing which comes in contact with Lasso must be washed before reuse.

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

IF ON SKIN immediately flush with plenty of water while removing contaminated clothing. Get medical attention. Wash clothing before reuse. Sensitized persons should avoid further contact and reuse of contaminated clothing. Get medical attention.

Get medical attention.

As soon as soap is available, wash thoroughly with soap and water.

Get medical attention.

In case of an emergency involving this product.
Call Collect, day or night, (314) 694-4000.

Environmental Hazards

This chemical has been identified in limited ground water sampling and there is the possibility that it can leach through the soil to ground water, especially where soils are coarse and ground water is near the surface. Following application, and during rainfall events that cause runoff, this chemical may reach surface water bodies including streams, rivers and reservoirs.

Care must be taken when using this product to prevent back siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Check valves or antisiphoning devices must be used on all mixing equipment. Disposal of excess pesticide, spray mixtures or rinsate should be according to label use instructions or according to the state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office.

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

Physical or Chemical Hazards

In case of:
SPILL OR LEAK, remove and keep away from humans and domestic animals.

ACTIVE INGREDIENT:	
*Atrachlor	66.7%
INERT INGREDIENTS:	33.3%
	100.0%

*Contains 667 grams/kilogram or 10.67 ounces/pound of 2-chloro-2',6'-diethyl-N-(methoxymethyl) acetanilide.

Patent application pending.

GENERAL INFORMATION

Lasso® herbicide is recommended for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label. This product may be applied either as a surface application after planting or shallowly incorporated prior to planting to blend the herbicide treatment into the upper 1 to 2 inches of soil. Except for minimum or conservation tillage systems, the seedbed should be fine, firm and free of clods and trash.

Read and carefully observe precautionary statements and all other information appearing on the labeling of all products used in mixtures and sequential treatments.

~~NOTE THAT GRAMOXONE SUPER HERBICIDE IS A POISON WHICH MAY BE FATAL OR HARMFUL TO HUMANS AND DOMESTIC ANIMALS.~~

Do not flood irrigate following application of this product or tank mixtures. This product will not control emerged seedlings.

Do not apply when conditions favor drift.

Use of this product not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

Flush sprayer with clean water after use.

* Lasso is a registered trademark of Monsanto Company.

~~*Gramoxone is a trademark of ICI Americas Inc.~~

SOIL TEXTURE

Applicators should evaluate soil and weed conditions carefully to assure that they choose the lowest effective label rate.

The recommended use rates of this product and the other herbicides labeled for use in tank mixtures with

when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, seed or feed by storage or disposal.

STORAGE:
NOT SUBJECT TO FREEZING.

Product can be stored under either heated or unheated conditions.

DISPOSAL:
Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or burned on site in a safe place away from water supplies. All disposal should be in accordance with applicable Federal, state or local procedures.

Emptied container retains product residue. Observe all labeled safeguards until container is destroyed.

Completely empty bag into application equipment. Then dispose of empty bag and box in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(omit)

(omit)

4760

this product generally vary with soil texture. Unless soil texture is specifically named, rate tables throughout this label refer to only three soil texture groups: coarse, medium and fine. The following is a complete listing of soil textures included in each of these three soil texture groups:

TEXTURAL	
SOIL TEXTURE GROUP	SOIL TEXTURE
COARSE:	sand, loamy sand, sandy loam
MEDIUM:	loam, silt loam, silt, sandy clay loam
FINE:	silty clay loam, clay loam, sandy clay, silty clay, clay

Refer to the above table to determine the correct soil texture group for the soil to be treated.

MIXING, SPRAYING AND HANDLING INSTRUCTIONS

NOTE

Direct contact or exposure to Lasso or spray mixtures of Lasso should be minimized. The following instructions for location and repairing equipment should be in order to minimize this exposure. Review the protective clothing requirements as listed in the Precautionary Statements section of this label (front panel) and do not use this product until you have the necessary protective clothing.

Equipment Cleaning & Repair

Cleaning and repair of transfer systems and application equipment is a source of exposure to this product. Care should be taken to minimize exposure during cleaning and repair of application equipment. Whenever possible, these systems or equipment should be rinsed before being cleaned or repaired. When repairs must be made during transfer or application, the equipment should be shutdown, and special care taken to avoid contact with the pesticide.

Always predetermine the compatibility of Lasso or labeled tank mixtures of Lasso with water carrier or sprayable fluid fertilizer carrier by mixing small proportional quantities in advance. See the "STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST" section in this label to determine the compatibility of this product and the labeled tank mixtures recommended for use with sprayable fluid fertilizer carrier. *Note: For soybeans only Lasso and the tank mixtures of Lasso plus Amiben, Lasso plus Canopy, Lasso plus Command, Lasso plus Gemma, Lasso plus Luron, Lasso plus Luron Plus, Lasso plus Preview, Lasso plus Sencor, or Lasso and carbo plus Trellan may be applied in sprayable fluid fertilizers. DO NOT apply Lasso plus Sceptor in sprayable fluid fertilizer. DO NOT apply Lasso or labeled tank mixtures of Lasso in sprayable fluid fertilizer on any other crops.*

Mix Lasso or labeled tank mixture of Lasso with the appropriate carrier as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with the appropriate carrier.

texture?

NOTE: Lasso and all tank mixtures listed for corn and soybeans in the "DIRECTION FOR USE" section may be applied in sprayable fluid fertilizers. DO NOT apply Lasso or labeled tank mixtures of Lasso in sprayable fluid fertilizers on any other crops.

3. If a compatibility agent is necessary, add it to the carrier while maintaining agitation to improve mixing or to prevent the formation of undesirable and unsprayable gels or precipitates. Use only compatibility agents cleared by FDA for this use. Read and follow all directions for use, cautionary statements and all other information appearing on the selected compatibility agent label. Check for adequate agitation.
4. Remove the screen prior to adding this product to the spray tank. Mixing of water dispersible granule formulations in sprayable fluid fertilizer carriers may be enhanced by prewetting with water and adding slowly through the screen into the tank. Continue agitation.
5. If a wettable powder or dry flowable formulation is used, make a slurry with water and add it slowly through the screen into the tank. Continue agitation.
6. If a flowable formulation is used, add slowly through the screen into the tank. Mixing and compatibility may be improved when the flowable is premixed one part flowable with one part water and added to the tank in diluted form.
7. Make a slurry of Lasso with water. Add it slowly through the screen into the tank.
8. Complete filling the sprayer tank with carrier. If Gramoxone Super is used, add the required amount near the end of the filling process. Remove hose from tank immediately after filling to avoid siphoning back into the carrier source.

TM

Maintain good agitation at all times until the contents of the tank are sprayed. NOTE: If spray mixture is allowed to settle at any time thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. Check for even distribution of spray droplets. To reduce loss of the chemical due to drift of a fine mist, apply at nozzle pressures below 40 psi.

*Amidon is a trademark of Rhône-Poulenc Inc.

*Conquest is a trademark of ICI Americas Inc.

*Diquat is a trademark of Monsanto Company.

*EPA is a trademark of the U.S. Environmental Protection Agency.

*EPA is a trademark of the U.S. Environmental Protection Agency.

*EPA is a trademark of the U.S. Environmental Protection Agency.

*EPA is a trademark of the U.S. Environmental Protection Agency.

STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST

Herbicides may not always mix evenly throughout a sprayable fluid fertilizer or the components may separate too quickly to make their combined use of practical value. This may be due to certain characteristics of the different fluid fertilizers. A simple test using small quantities of the components is suggested to provide compatibility potential.

The test follows:

A. Materials Required For A Compatibility Test

1. Two one-quart jars with lid or stopper (marked "with" and "without").
2. TEA spoons (for a more exacting test, a five to ten milliliter (ml.) pipette or graduated cylinder is desirable).
3. Sprayable fluid fertilizer to be tested.
4. The herbicide chemicals to be mixed.
5. A compatibility agent (the purpose of the adjunct is to help keep the fertilizer and crop protection chemical in suspension if this assistance is needed)

TM Gramoxone is a trademark of ICI Americas Inc.


TM Treflan is a trademark of Elanco Products Company.

00021


6760

B. Procedure


1. Add one pint of the sprayable fluid fertilizer that will be used or other herbicide carrier to each jar marked "with" and "without"



2. To the jar marked "with", add $\frac{1}{4}$ TEAspoon or 1.2 milliliters of a suitable compatibility agent; shake gently for five to ten seconds to mix ($\frac{1}{4}$ TEAspoon in one pint is the equivalent of two pints per 100 gallons of liquid fertilizer).



3. To each jar add the appropriate amount of herbicides. If more than one is used, add them separately with the wettable powders or dry flowables added first, flowables second and liquid last. Shake gently five to ten seconds after each addition.



listed in the following tables.

Amount to Be Added Per Pint Of Sprayable Fluid Fertilizer Assuming Volume is 25 gal/A)

HERBICIDE	RATE/A	Level	TEAspoons
WDG'S	1 lb. =		1.5
or	2 lb. =		3
	3 lb. =		4.5
Dry Flowables	4 lb. =		6
	5 lb. =		7.5

HERBICIDE	RATE/A	Level	TEAspoons	Milliliters
Emulsifiable	1 pint =		0.5	or 2.4
Concentrates or	1 quart =		1	or 4.7
Flowables or	2 quarts =		2	or 9.5
Liquids or	3 quarts =		3	or 14.2
Solutions	1 gallon =		4	or 19.0
	5 quarts =		5	or 23.8

This compatibility test is designed for 25 gallons of spray per acre, with the maximum labeled rate of herbicide. For changes in spray volume or herbicide rates, make appropriate changes in the ingredients of the test. Regardless of spray volume, the amount of compatibility agent should be equal to two or three pints (two pints = $\frac{1}{4}$ TEAspoon or 1.2 milliliters, three pints = $\frac{1}{4}$ TEAspoon or 1.8 milliliters per pint of sprayable fluid fertilizer) per 100 gallons of liquid fertilizer. (omit)

C. Observations and Decisions

1. If the herbicide(s) and the sprayable fluid fertilizer are compatible
2. If a compatibility agent is necessary

00022

Five minutes after the final addition and mixing, observe both jars for the formation of large flakes, sludge, gels or other precipitates. Observe if the herbicide(s) cannot be physically mixed with the liquid fertilizer but remains as small oily particles in the solution.

If incompatibility in any form described above occurs in the jar "with" the compatibility agent added, the liquid fertilizer and the herbicide(s) should not be used together in the same spray tank.

If incompatibility as described above occurs in the jar "without" the adjuvant but not in the jar "with" adjuvant, the use of a compatibility adjuvant is recommended.

Both jars should be allowed to stand and be observed periodically for one-half hour. If the separate layers of liquid fertilizer and additives can be resuspended by shaking, commercial application is possible. An emulsifiable concentrate normally will go to the top after standing; WDG's or dry flowables will either settle to the bottom of the tank or jar, or float to the top, depending upon the density of the fertilizers.

If the herbicide(s) is compatible with fluid fertilizer in the foregoing test without having to use a compatibility agent, fluid fertilizer may be used for the premixing. If it is not compatible without the compatibility agent, the herbicide(s) should be premixed with water before adding to the spray tank.

APPLICATION INFORMATION

APPLICATION SYSTEMS

Do not apply this product through any type of irrigation system.

Ground Broadcast Treatment — Apply this product and the labeled tank mixtures in 15 or more gallons of solution per acre using broadcast boom equipment. The carrier may be either water or sprayable fluid fertilizer as specified for the crop to be treated in the "DIRECTIONS FOR USE" section of this label. Do not apply during periods of gusty winds, when winds are in excess of 15 mph or when other conditions favoring drift exist.

Band Treatment — Apply a broadcast equivalent rate and volume per acre. To determine these:

Band width in inches	x	Broadcast RATE per acre	=	Band RATE per acre
Row width in inches				
Band width in inches	x	Broadcast VOLUME per acre	=	Band VOLUME per acre
Row width in inches				

Aerial — Fixed wing, helicopter — Apply the recommended rates of this product or approved tank mixtures in 3 to 10 gallons of water per acre. AVOID DRIFT — DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray or fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Ensure uniform application. To avoid streaked, uneven or overlapped application use mechanical marking systems only. HUMAN FLAGGERS PROHIBITED!

APPLICATION TIMING AND METHODS

NOTE: Unless specifically recommended, only one treatment of Lasso may be applied to a labeled crop.

Preemergence Surface Application — This product and all labeled tank mixtures except Lasso plus EptamTM Lasso plus TreflanTM and Lasso plus Command may be applied to the soil surface after planting and prior to either crop or weed emergence. Apply within 5 days of last preplant tillage. If weeds emerge after treatment, or if treatment is applied more than 5 days after last preplant tillage, rotary hoe or shallowly cultivate immediately to improve performance.

Precipitation or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone. The amount of precipitation or overhead sprinkler irrigation required depends on existing soil moisture, soil type and percent organic matter content, but 1/3 to 3/4 inches is normally adequate. Performance is improved when moisture is received within 7 days after application and prior to weed emergence. High intensity or excessive rainfall or excessive irrigation after application may reduce control.

Preplant Incorporation — This product and many of the labeled tank mixtures may be mixed into the soil using shallow incorporation equipment any time within 7 days prior to planting. Refer to the crop to be treated in the "DIRECTIONS FOR USE" section of this label to determine if preplant incorporation is recommended. Apply the recommended treatment rate to the soil surface as a broadcast application. ~~Either surface or soil-moisture~~ subsequent precipitation or irrigation is required to bring incorporated herbicide treatments into contact with germinating weed seedlings. If weeds emerge after treatment, rotary hoe or shallowly cultivate immediately to improve performance.

NOTE: Do not incorporate unless specifically recommended in this label for the crop and herbicide treatment selected. Incorporation of this product and its labeled tank mixtures on coarse soil types may reduce length of weed control due to leaching with rainfall or irrigation.

1-Pass Incorporation (Surface Blend) — Shallowly incorporate (Surface Blend) the treatment into the upper 1 to 2 inches of the soil. Equipment should be operated at manufacturer's designed speed for incorporation to ensure adequate mixing and distribution of the herbicide treatment in the soil.

Equipment design including any drag attachments must be adequate to avoid soil ridging which may result in streaked or reduced weed control. Soil conditions, including moisture content and crop residue levels, must be suitable to allow thorough and uniform mixing with the equipment used for 1-pass incorporation.

Contact Monsanto Company for additional information regarding recommended equipment and methods for Surface Blend application.

2-Pass Incorporation: When 2-pass incorporation is used, shallowly incorporate the herbicide treatment into the upper 1 to 2 inches of the soil with equipment set to work the soil NO DEEPER THAN 4 INCHES. The second pass must be made at an angle to and no deeper than the first pass to ensure proper distribution of the herbicide treatment in the soil.

Cultivation Information — Delay cultivation after application for as long as possible unless weeds or grasses emerge. Shallowly cultivate or rotary hoe immediately if weeds or grasses emerge. If cultivation is necessary because of soil crusting or compaction, set equipment shallow and minimize lateral soil movement to avoid dilution or displacement of the herbicide treatment. Only after

TM

The amount of precipitation or irrigation required depends on existing soil moisture, soil type and percent organic matter content, but 1/3 to 3/4 inch is normally adequate.

96024

weeds emerge in the treated band, set cultivator to throw soil into the row covering the band.

*Surface Blend is a trademark of Monsanto Company

*Eptam is a trademark of ICI Americas Inc.

WEEDS CONTROLLED

Applicators should evaluate soil and weed conditions carefully to assure that they choose the lowest effective label rate.

When applied as directed under conditions described, yellow nutsedge and the annual grasses and broadleaf weeds listed are CONTROLLED by Lasso.

ANNUAL GRASSES	ANNUAL BROADLEAVES
ryegrass	Carnation
Chenopodium crus-galli	Mollugo verticillata
Crabgrass	Galinsoga
Digitaria pruriens	Galinsoga spp.
Digitaria sanguinalis	Groundcherry, cutleaf*
Foxtail, giant	Physalis angulata
Setaria faberii	Nightshade, eastern black*
Foxtail, green, robust	Solanum elaeagnifolium
purple, robust white	Nightshade, hairy*
Setaria viridis	Solanum sarothamoides
Foxtail, yellow	Pigweed, Canada-weed
Setaria lutescens	Amaranthus spp.***
Goosegrass	Portulaca
Elymus indica	Portulaca oleracea
Panicum, browneyed	Pasley, Florida
Panicum fasciculatum	Richardia scabra
Panicum, tall	
Panicum dichotomiflorum	SEDGE
Rice, red**	Nutsedge, yellow**
Oryza sativa	Cyperus esculentus
Signalgrass, broadleaf	
Brachiaria platyphylla	
Sprangletop, red	
Leptochloa filiformis	
Witchgrass	
Panicum capillare	

*Use a minimum of 3.75 pounds of this product per acre on coarse soils, and 4.5 to 6 pounds alone or in tank-mixture combinations on medium and fine-textured soils for control of these weeds.

**Use 4.5 to 6 pounds of this product per acre applied alone or in tank-mixture combinations listed on this label. Preplant incorporate only for control of these weeds.

When tank mixed with Scepter, use 3 to 4.5 pounds of this product per acre and preplant incorporate only for control of these weeds.

***Does not include waterhemp (*Amaranthus tuberculatus*).

When applied preplant incorporated, or preemergence immediately after planting and within 5 days of last tillage, Lasso at a rate of 3.75 to 6 pounds per acre on a broadcast basis applied alone or in a labeled tank mixture REDUCES COMPETITION from the following hard-to-control annual grasses and broadleaf weeds.

Tri-Command is a trademark of the FMC Corporation.

When applied as directed under conditions described, this product and tank mixtures of this product will control or reduce competition from the weeds listed. Refer to the crop on which treatment is intended to determine the rates and tank mixtures recommended.

ANNUAL GRASSES

Johnsongrass, seedling
Sorghum halepense
Millet, proso***
Panicum miliaceum
Panicum, Texas*
Panicum texanum
Sawtooth, Gumbo
Cenchrus incertus
Shattercane, wild cane**
Sorghum bicolor

SEDGE

Nutsedge, yellow***
Cyperus esculentus

*For reduced competition from this weed, use a minimum of 4.5 pounds of this product per acre and apply under sprinkler irrigation systems.

**Use 6 pounds per acre of this product applied alone or in tank mixture combinations listed in this label. Preplant incorporate only to reduce competition from this weed.

***Use 4.5 to 6 pounds of this product per acre as a preemergence surface application on coarse textured soils with less than 3% organic matter to reduce competition from this weed.

****For reduced competition from this weed, use 6 pounds per acre of this product applied preplant incorporated. Improved control may be obtained by applying subsequent labeled treatments of registered postemergence herbicides.

When applied as directed under conditions described, tank mixtures of Lasso with products listed in the following table will control or reduce competition from the additional broadleaf weeds as indicated. Refer to the crop on which treatment is intended in the "DIRECTIONS FOR USE" section of this label to determine the tank mixtures and rates recommended for use in each crop.

• = Not Applicable
NOTE: C = Control R = Reduced Competition

	Amber	Canopy	Command	Gemini	Inturon
Beggartweed, Florida					
<i>Desmodium tortuosum</i>	•	C	•	C	•
Cocklebur					
<i>Xanthium strumarium</i>	•	C	•	C	•
Dandelion					
<i>Hemp sesbania</i>					
<i>Sesbania exaltata</i>	•	C	•	C	•
Copperleaf, Hophornbeam					
<i>Acalypha</i> spp.	•	C	•	•	•
Cupress, woolly					
<i>Eriochloa villosa</i>	•	•	R	•	•
Groundcherry, annual					
<i>Physalis</i> spp.	•	C	•	C	C
Jimsonweed					
<i>Datura stramonium</i>	•	C	C	C	•
Johnsongrass, seedling*					
<i>Sorghum halepense</i>	•	•	•	•	•
Kochia					
<i>Kochia scoparia</i>	•	•	•	•	•
Lambsquarters					
<i>Chenopodium album</i>	C	C	C	C	C
Morningglory, Tall*					
<i>Ipomea purpurea</i>	•	C	•	C	•

ANNUAL BROADLEAVES

Beggartweed, Florida
Desmodium tortuosum
Lambsquarters
Chenopodium album
Ragweed, common
Ambrosia artemisiifolia
Sida, prickly, Teawood
Sida spinosa
Smartweed
Polygonum pensylvanicum
Polygonum persicaria

	Amber	Canopy	Command	Gemini	Inturon
Pitted					
<i>Ipomea lacunosa</i>	•	C	•	C	•
Ivyleaf*					
<i>Ipomea hederacea</i>	•	C	•	C	•
Entireleaf*					
<i>Ipomea hederacea</i>	•	C	•	C	•
<i>var. integruscula</i>	•	C	•	C	•
Santhol					
<i>Jacquemontia tamnifolia</i>	•	C	•	C	•
Mustard					
<i>Brassica</i> spp.	R	C	•	C	C
Nightshade, eastern black					
<i>Solanum ptycanthum</i>	R	C	R	C	R
Nightshade, hairy					
<i>Solanum sarachoides</i>	•	C	•	C	•
Ragweed, common					
<i>Ambrosia artemisiifolia</i>	C	C	C	C	C
Sicklepod					
<i>Cassia obtusifolia</i>	•	C	•	R	•
Sida, prickly, Teawood					
<i>Sida spinosa</i>	•	C	C	C	C
Smartweed					
<i>Polygonum pensylvanicum</i>					
<i>Polygonum persicaria</i>	C	C	R	C	C
Starbur, bristly					
<i>Acanthospermum hispidum</i>	•	•	•	•	•
Velvetleaf, Buttonweed*					
<i>Abutilon theophrasti</i>	R	C	C	C	R
Waterhemp					
<i>Amaranthus tuberculatus</i>	•	•	•	•	•

• = Not Applicable
NOTE: C = Control R = Reduced Competition

	Lorox Plus	Prevail	Scepter	Sencor / Leone	Trellan
Beggartweed, Florida					
<i>Desmodium tortuosum</i>	•	R	R	•	•
Cocklebur					
<i>Xanthium strumarium</i>	C	C	C	R	•
Coffaweed					
<i>Hemp sesbania</i>	•	C	•	•	•
<i>Sesbania exaltata</i>	•	C	•	•	•
Copperleaf, Hophornbeam					
<i>Acalypha</i> spp.	•	C	•	•	•
Cupress, woolly					
<i>Eriochloa villosa</i>	•	•	•	•	R
Groundcherry, annual					
<i>Physalis</i> spp.	C	C	C	C	•
Jimsonweed					
<i>Datura stramonium</i>	C	C	C	R	•
Johnsongrass, seedling*					
<i>Sorghum halepense</i>	•	•	C	•	•
Kochia					
<i>Kochia scoparia</i>	•	•	•	•	C
Lambsquarters					
<i>Chenopodium album</i>	C	C	C	C	C
Morningglory, Tall*					
<i>Ipomea purpurea</i>	R	R	R	•	•
Pitted					
<i>Ipomea lacunosa</i>	R	R	C	•	•

	Lorox Plus	Prevail	Scepter	Sencor / Leone	Trellan
Ivyleaf*					
<i>Ipomea hederacea</i>	R	R	R	•	•
Entireleaf*					
<i>Ipomea hederacea</i>	R	R	R	•	•
<i>var. integruscula</i>	R	R	R	•	•
Santhol					
<i>Jacquemontia tamnifolia</i>	•	•	C	•	•
Mustard					
<i>Brassica</i> spp.	C	C	C	•	•
Nightshade, eastern black					
<i>Solanum ptycanthum</i>	C	R	C	R	R
Nightshade, hairy					
<i>Solanum sarachoides</i>	•	•	•	•	•
Ragweed, common					
<i>Ambrosia artemisiifolia</i>	C	C	C	C	•
Sicklepod					
<i>Cassia obtusifolia</i>	•	•	R	R	•
Sida, prickly, Teawood					
<i>Sida spinosa</i>	C	C	C	•	•
Smartweed					
<i>Polygonum pensylvanicum</i>					
<i>Polygonum persicaria</i>	C	C	C	C	•
Starbur, bristly					
<i>Acanthospermum hispidum</i>	•	•	•	•	•
Velvetleaf, Buttonweed*					
<i>Abutilon theophrasti</i>	C	C	C	R	•
Waterhemp					
<i>Amaranthus tuberculatus</i>	C	C	C	•	•

For the scientific names to the weeds listed below, refer to the appendix at the end of this booklet.

NOTE: C = Control R = Reduced Competition

ANNUAL GRASSES

	Lasso [®]	Lasso [®] plus					
		Amben [®]	Atrazine	Banvel [®]	Bladex [®]	Canopy [®]	Comm [®]
Barnyardgrass	C	C	C	C	C	C	C
Crabgrass	C	C	C	C	C	C	C
Cudgrass woolly	R	R	R	R	R	R	R
Foxtail giant green robust							
Purple robust white							
Yellow	C	C	C	C	C	C	C
Goosegrass	C	C	C	C	C	C	C
Johnsongrass seedling ¹	R	R	R	R	R	R	R
Allet. proso ²	C	C	C	C	C	C	C
Panicum browntop, tall	C	C	C	C	C	C	C
Panicum Texas ³	R	R	R	R	R	R	R
Rice, red ⁴	C	C	C	C	C	C	C
Sandbar Grassbur ⁵	R	C	C	C	C	C	C
Shattercane wild cane ⁶	R	R	R	R	R	R	R
Signalgrass broadleaf	C	C	C	C	C	C	C
Sprangletop, red	C	C	C	C	C	C	C
Witchgrass	C	C	C	C	C	C	C

ANNUAL BROADLEAVES

	Lasso [®]	Lasso [®] plus					
		Amben [®]	Atrazine	Banvel [®]	Bladex [®]	Canopy [®]	Comm [®]
Beggarweed, Florida ¹	R	C	C	C	C	C	C
Carpetweed	C	C	C	C	C	C	C
Cocklebur	C	C	R	R	R	C	C
Coffeeweed, Hemp sesbania	C	C	C	C	C	C	C
Copperleaf, hophornbeam	C	C	C	C	C	C	C
Galinsoga	C	C	C	C	C	C	C
Groundcherry annual	C	C	C	C	C	C	C
Groundcherry cutleaf ²	C	C	C	C	C	C	C
Jimsonweed	C	C	C	C	C	C	C
Kochia	C	C	C	C	C	C	C
Lambsquarters ³	R	C	C	C	C	C	C
Morningglory Tail ⁴	C	C	R	R	R	C	C
Pitted, ivyleaf ⁵ entire leaf ⁶	C	C	C	C	C	C	C
Smallflower	R	R	C	C	C	C	C
Mustard	C	C	C	C	C	C	C
Nightshade black ⁷ , hairy ⁸	C	C	C	C	C	C	C
Pigweed, Carelessweed	C	C	C	C	C	C	C
Purslane	C	C	C	C	C	C	C
Posley, Florida	C	C	C	C	C	C	C
Ragweed common ⁹	R	C	C	C	C	C	C
Ragweed, giant	C	C	C	C	C	C	C
Sicklepod	C	C	C	C	C	C	C
Sida, prickly, Teaweed ¹⁰	R	C	C	C	C	C	C
Smartweed ¹¹	R	C	C	C	C	C	C
Starbur, bristly	R	R	C	C	C	C	C
Sunflower, common	C	C	C	C	C	C	C
Velvetleaf Buttonweed ¹²	C	R	R	R	R	C	C
Waterhemp	C	C	C	C	C	C	C
SEDGE							
Nutsedge, yellow ¹³	C	C	C	C	C	C	C

NOTE: C = Control R = Reduced Competition

ANNUAL GRASSES

	Lasso [®] plus				
	Inuron	Lasso [®] Plus	Preempt [®]	Scout [®]	TriAllin [®]
Barnyardgrass	C	C	C	C	C
Crabgrass	C	C	C	C	C
Cupgrass, woolly ¹	•	•	R	R	R
Foxtail giant: green robust					
purple, robust white:					
yellow	C	C	C	C	C
Goosegrass	C	C	C	C	C
Johnsongrass, seedling ²	R	R	R	C	R
Millet, proso ³	•	•	•	•	•
Panicum, browntop: fall	C	C	C	C	C
Panicum, Texas ⁴	R	R	R	R	R
Rice, red ⁵	•	•	C	C	C
Sandbur Grassbur ^{2,6}	•	•	•	•	•
Shattercane, wild cane ¹	•	•	R	R	R
Signalgrass, broadleaf	C	C	C	C	C
Sprangletop, red	C	C	C	C	C
Witchgrass	C	C	C	C	C

ANNUAL BROADLEAVES

	Lasso [®] plus				
	Inuron	Lasso [®] Plus	Preempt [®]	Scout [®]	TriAllin [®]
Beggarweed Florida ²	•	•	R	•	R
Carpetweed	C	C	C	C	C
Cocklebur	•	C	C	C	R
Coffeeweed	•	•	C	•	•
Copperleaf hophornbeam	•	•	C	•	•
Galinsoga	C	C	C	C	C
Groundcherry, annual	C	C	C	C	•
Groundcherry, cutleaf ²	C	C	C	C	C
Jimsonweed	•	C	C	C	R
Kochia	•	•	•	•	C
Lambsquarters ¹	C	C	C	C	•
Morningglory tall ³	•	R	R	R	•
Pitted	•	R	R	C	•
Ivyleaf ⁴ entireleaf ⁵	•	R	R	R	•
Smallflower	•	•	•	C	•
Mustard	C	C	C	C	•
Nightshade, black ⁷ , hairy ⁷	C	C	C	C	C
Pigweed, Carelessweed	C	C	C	C	C
Purslane	C	C	C	C	C
Pusley, Florida	C	C	C	C	C
Ragweed, common ²	C	C	C	C	•
Ragweed, giant	•	R	R	R	•
Sicklepod	•	•	•	R	R
Sida, prickly, Tearweed ²	C	C	C	C	•
Smartweed ¹	C	C	C	C	•
Starbur, bristly	•	•	•	C	C
Sunflower common	•	•	R	C	•
Velvetleaf Buttonweed ¹	R	C	C	C	R
Waterhemp	C	C	C	C	C

SEDGE

Nutsedge yellow ¹	•	•	C	C	C
------------------------------	---	---	---	---	---

- 13760
- ¹Use 4 quarts per acre of this product applied alone or in tank mix combinations. Preplant incorporate only to reduce competition from this weed.
 - ²For reduced competition from this weed with Lasso alone, use 2.5 to 4 quarts of Lasso per acre.
 - ³Plant corn into freshly prepared seedbed. Apply 4 quarts of Lasso per acre preemergence surface immediately after corn planting for reduced competition from this weed. Improved control may be obtained by applying a labeled postemergence herbicide.
 - ⁴Use a minimum of 3 quarts of Lasso per acre and apply through center pivot irrigation or under sprinkler irrigation systems. Only Lasso and Lasso plus atrazine tank mixture can be applied through center pivot irrigation systems.
 - ⁵Use 3 to 4 quarts of Lasso per acre applied alone or in tank mixture and apply preplant incorporated only for control on medium and fine textured soils. When tank mixed with Scepter, use 2 to 3 quarts of Lasso per acre and preplant incorporate only. On coarse textured soils with less than 3% organic matter, use 3 to 4 quarts of Lasso per acre as a preemergence surface application to reduce competition from yellow nutsedge.
 - ⁶Apply 2.5 to 3 quarts of Lasso plus 1 to 2 quarts of atrazine 4L per acre. Preemergence surface applications should be applied to a freshly prepared seedbed. For best control, rotary hoe or initiate overhead irrigation (0.33 to 0.5 inch) within 24 hours of application.
 - ⁷Use a minimum of 2.5 quarts of Lasso per acre on coarse soils, or 3 to 4 quarts of Lasso per acre on medium and fine soils when used alone or in tank mixture with Amiben, Banvel, Canopy, Command, Preview, Sencor/Lexone or Treflan for control of this weed.
 - ⁸For best results on these species when using Lasso plus Scepter tank mixture, use preplant incorporated treatments.

NOTE: The 3-way tank mixture of Lasso plus atrazine plus Bladex controls or reduces competition as specified for the respective weeds listed under the Lasso plus atrazine and Lasso plus Bladex recommendations.

The 3-way tank mixture of Lasso plus Amiben plus Lexone or Sencor controls or reduces competition as specified for the respective weeds listed under the Lasso plus Amiben and Lasso plus Lexone or Lasso plus Sencor recommendations.

~~*For best results on these species when using Lasso plus Scepter tank mixture, use preplant incorporated treatments.~~

~~The 3-way tank mixture of Lasso plus Amiben plus Lexone or Sencor controls or reduces competition as specified for the respective weeds listed under the Lasso plus Amiben and Lasso plus Lexone or Sencor recommendations.~~

~~Basagran™ may be applied sequentially to Lasso in soybeans for control of additional emerged weeds listed on the Basagran label, such as cocklebur and velvetleaf.~~

Certain tank mixtures of Lasso and broadleaf herbicides may be combined with applications of Gramoxone Super for control of many emerged weeds prior to soybean emergence.

Refer to the Gramoxone Super label for a list of emerged weeds controlled by tank mixtures with these products.

~~Basagran is a trademark of BASF Aktiengesellschaft~~

™ Amiben is a trademark of Rhône-Poulenc.

™ Panvel is a trademark of Sandoz, Inc.

™ Bladex, Canopy, Gemini, Lexone, Lorox and Preview are trademarks of E. I. duPont de Nemours and Company.

™ Scepter is a trademark of American Cyanamid Corporation.

™ Sencor is a trademark of the parent company of Farben Fabricken Bayer GmbH.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, seed or feed by storage or disposal.

STORAGE:

NOT SUITABLE TO FREEZING.

Product can be stored under either heated or unheated conditions.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or buried on site in a safe place away from water supplies. All disposal should be in accordance with applicable Federal, state or local procedures.

Emptied container retains product residue. Observe all labeled safeguards until container is destroyed.

Completely empty bag into application equipment. Then dispose of empty bag and box in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

← (Moved to area following Physical and Chemical Hazards section)

← (Insert corn and grain sorghum (milo) sections)

CORN (All Types)

LASSO[®]

Apply this product in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground—Broadcast boom banded

Aerial—Fixed wing, helicopter

Center Pivot Irrigation

Dry Bulk Fertilizer Impregnation

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated—Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. ~~This product must be incorporated when treating furrow irrigated corn. Performance may be increased by packing the soil following incorporation. If furrow irrigation is to be used, irrigation within 10 days following planting may improve weed control.~~

Preemergence Surface—Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

DO NOT apply this product postemergence, sequential or lay-by.

DO NOT apply any alachlor containing product sequential to this product.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse***	3 to 3.4	3 to 3.4
Medium	3 to 4.1	3 to 4.1
Fine	3 to 4.1	3.8 to 4.9

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~soil~~ group for the soil to be treated (soil texture).

**Use the higher rate in the recommended range in areas of heavy weed infestations. Use a minimum of 3.8 to 6 pounds quarts of this product per acre on coarse soils and 4.5 to 6 pounds ~~quarts~~ on medium or fine textured soils to control black or hairy nightshade ~~to reduce competition from hard to control weeds listed in this label.~~

***When applied through center pivot irrigation or under sprinkler irrigation systems on coarse soils, use a minimum of ~~3.8~~ pounds per acre of this product.

← 4.5 pounds

BEST AVAILABLE COPY

00032

17 7 65

corn insert

Preplant Incorporated Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse***	3.8	3.8
Medium	3.8 to 4.5	3.8 to 4.5
Fine	3.8 to 4.5	4.5 to 5.3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~soil~~ group for the soil to be treated. soil textural

**Use the higher rate in the recommended range in areas of heavy weed infestations. Use 3.8 to 5.3 pounds per acre where furrow irrigation is used. Use a minimum of 2 quarts on coarse textured soils for control of yellow nutsedge or red rice. Rates may be increased to a maximum of 4.5 pounds on any soil type when heavy infestations of yellow nutsedge or red rice are present, or to reduce competition from the hard-to-control weeds listed in this label. 6 pounds

**LESSO® PLUS ATRAZINE
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

- Ground—Broadcast boom, banded
- Aerial—Fixed wing; helicopter
- Center Pivot Irrigation
- Dry Bulk Fertilizer Impregnation

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated—Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface—Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

~~DO NOT apply to sweet corn after crop emergence.~~

~~DO NOT apply as a coverable fluid fertilizer after crop emergence since crop injury may result.~~

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: PLANT ONLY CORN, PEANUTS, SORGHUM OR SOYBEANS the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn. DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

BEST AVAILABLE COPY

00033

12760

corn insert

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE **			
	Less than 3% organic matter		3% or more organic matter	
	Lasso [®] + atrazine 4L***		Lasso + atrazine 4L***	
	(pounds)	(quarts)	(pounds)	(quarts)
Coarse	2.3 to 3	0.75 to 2	2.3 to 3	1 to 2
Medium	2.3 to 3	1 to 2	3 to 3.4	1.25 to 2
Fine	3 to 3.4	1.25 to 2	3 to 4.1	1.25 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~texture~~ group for the soil to be treated. soil textural

**Use the higher rates in the recommended ranges in areas of heavy weed infestations.

***Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

NOTE: When applied through center pivot irrigation or under sprinkler irrigation systems on coarse soils, use 1 quart per acre of atrazine 4L and a minimum of 3.8 pounds per acre of Lasso in this tank mixture.

Preplant Incorporated Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE **			
	Less than 3% organic matter		3% or more organic matter	
	Lasso [®] + atrazine 4L***		Lasso + atrazine 4L***	
	(pounds)	(quarts)	(pounds)	(quarts)
Coarse	2.6 to 3.4	1.25 to 2	2.6 to 3.4	1.25 to 2
Medium	2.6 to 3.4	1.25 to 2	3.4 to 3.8	1.5 to 2
Fine	3.4 to 3.8	1.5 to 2	3.4 to 4.1	1.5 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~texture~~ group for the soil to be treated. soil textural

**Use the higher rates in the recommended ranges in areas of heavy weed infestations. Use ~~3.8 pounds of Lasso per acre for control of yellow nutsedge or to reduce competition from the hard to control weeds listed in this label.~~

***Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

LASSO[®] plus BANVEL (TANK MIXTURE)

For use in ILLINOIS, IOWA, MINNESOTA and WISCONSIN on level or flat-planted field corn on fine textured (silty clay loam, clay loam, sandy clay, silty clay or clay) soils with more than 4 percent organic matter.

Apply in water or sprayable fluid fertilizer solutions for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label

NOTE: Use on coarse or medium textured soils or on fine textured soils with 4 percent or less organic matter may result in crop injury and/or destruction

96034

APPROVED APPLICATION SYSTEMS

Ground—Broadcast boom banded

NOTE: DO NOT apply by air or injection through center pivot irrigation systems

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using ground spray equipment.

APPROVED APPLICATION METHODS

Preemergence Surface—Apply this tank mixture after planting, before crop and weeds emerge and within 5 days of last preplant tillage operation. Corn seeds must be planted 1-1/2 inches or deeper beneath the soil surface. Direct chemical contact with corn seed must be avoided since crop injury may result. Apply far enough behind planter equipment to avoid any incorporation by the planter wheel or other covering device. If corn seeds are planted less than 1-1/2 inches beneath the soil surface, delay application until corn has spiked.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: PREVENT DRIFT TO SOYBEANS OR OTHER DESIRABLE PLANTS. Do not use on furrow irrigated corn, or when corn is planted at the bottom of a furrow, utilizing lister, till or other similar planting methods. DO NOT incorporate prior to planting or corn emergence. If it is necessary to drag for leveling or rotary hoe to break soil crust, DO NOT disturb the soil more than 1/2 inch deep.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury and/or destruction.

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE	
	More than 4 % organic matter Lasso® (pounds)	Below 4 % Bladex (quarts)
Fine-silty clay loam through clay	3.8	0.5

**LASSO® plus BLADEX
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge, and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground—Broadcast boom, banded

Aerial—Fixed wing, helicopter

Dry Bulk Fertilizer Impregnation

Reference: The "APPLICATION SYSTEMS" section of this label provides information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated—Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil

Preemergence Surface—Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation

2 7 6
corn insert

Refer to the APPLICATION TIMING AND METHODS section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

BROADCAST RATE PER ACRE				
SOIL TEXTURAL GROUP*	1% to 3% organic matter		3% or more organic matter	
	Bladex Lasso [®] + 4L***		Bladex Lasso + 4L***	
	(pounds)	(quarts)	(pounds)	(quarts)
Coarse**	2.3 to 3	1 to 1.2	2.3 to 3	1 to 1.6
Medium	2.3 to 3	2 to 1.4	3 to 3.4	1.5 to 2
Fine	3 to 3.4	1.1 to 1.3	3 to 4.1	1.3 to 2.2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~soil~~ group for the soil to be treated. soil texture

**On coarse textured soils containing less than 2% organic matter, use Lasso plus Bladex only on sandy loam. Do not use on sand and loamy sand with less than 2% organic matter.

***Use rates listed in this label using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 lbs of Bladex 80W.

Preplant Incorporated Application Rates

BROADCAST RATE PER ACRE				
SOIL TEXTURAL GROUP*	1% to 3% organic matter		3% or more organic matter	
	Bladex Lasso [®] + 4L***		Bladex Lasso + 4L***	
	(pounds)	(quarts)	(pounds)	(quarts)
Coarse**	2.6 to 3.4	1.2 to 1.4	2.6 to 3.4	1.4 to 1.8
Medium	2.6 to 3.4	1.4 to 1.6	3.4 to 3.8	1.5 to 2.2
Fine	3.4 to 3.8	1.5 to 2	3.4 to 4.1	1.5 to 2.4

NOTE: See Incorporation instructions.

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~soil~~ group for the soil to be treated. soil texture

**On coarse textured soils containing less than 2% organic matter, use Lasso plus Bladex only on sandy loam. Do not use on sand and loamy sand with less than 2% organic matter.

***Use rates listed in this label when using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.

← (insert Lasso plus Princep tank mixture).

**LASSO[®] plus ATRAZINE PLUS
BLADEX (TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

PROVED APPLICATION SYSTEMS

Ground—Broadcast boom, banded

Aerial—Fixed wing, helicopter

00036

LASSO® plus PRINCEP™
(TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom, banded
Aerial - Fixed wing, helicopter

Reference: "THE APPLICATIONS SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: the "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**					
	Less than 3% organic matter			3% or more organic matter		
	Lasso® (pounds)	+	Princep 4L*** (quarts)	Lasso (pounds)	+	Princep 4L*** (quarts)
Coarse - sandy loam only	2.3 to 3		0.75 to 2	2.3 to 3		1 to 2
Medium	2.3 to 3		1 to 2	3 to 3.4		1.25 to 2
Fine	3 to 3.4		1.25 to 2	3 to 4.1		1.25 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rates in the recommended ranges in acres of heavy weed infestations.

***Use rates listed in this label when using Princep 4L. Use equivalent rates when using Princep 80W or 90% dry flowable formulations. One quart of the 4L formulation equals 1.25 pounds of the 80W formulation and 1.1 pounds of the 90% dry flowable formulations.

Preplant Incorporated Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**					
	Less than 3%			3% or more		
	organic matter			organic matter		
	Lasso@	+	Princep	Lasso	+	Princep
	(pounds)		(4L*** (quarts)			4L*** (quarts)
Coarse - sandv						
loam only	2.6 to 3.4		1.25 to 2	2.6 to 3.4		1.25 to 2
Medium	2.6 to 3.4		1.25 to 2	3.4 to 3.8		1.5 to 2
Fine	3.4 to 3.8		1.5 to 2	3.4 to 4.1		1.5 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rates in the recommended ranges in areas of heavy weed infestations.

***Use rates listed in this label when using Princep 4L. Use equivalent rates when using Princep 80W or 90% dry flowable formulations. One quart of 4L formulation equals 1.25 pounds of 80W and 1.1 pounds of the 90% dry flowable formulations.

corn insert

Refer to the APPLICATION SYSTEMS section of this label for detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface—Apply this tank mixture after planting, before crop and weed emergence and within 7 days after last preplant tillage operation.

Preplant Incorporated—Apply this tank mixture within 7 days prior to planting and shallowly incorporate or surface blend into the upper 1 to 2 inches of soil.

Reference the APPLICATION TIMING AND METHODS section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: PLANT ONLY CORN, PEANUTS, SORGHUM OR SOYBEANS* the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans are planted the following year. **DO NOT** plant soybeans the year following use of this tank mixture on narrow-ridged corn. **DO NOT** graze treated area or feed treated forage to livestock for 30 days following application of this tank mixture.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	Less than 3% organic matter		
	Lasso [®] + (pounds)	atrazine + 4L** (quarts)	Bladex 4L*** (quarts)
Coarse****	2.3 to 3	0.75	0.5 to 1
Medium	2.3 to 3	0.75 to 1	1 to 1.25
Fine	3 to 3.4	1 to 1.25	1 to 1.5

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	3% or more organic matter		
	Lasso [®] + (pounds)	atrazine + 4L** (quarts)	Bladex 4L*** (quarts)
Coarse****	2.3 to 3	0.75 to 1	1 to 1.25
Medium	3 to 3.4	1 to 1.25	1 to 2
Fine	3 to 4.1	1.25 to 1.5	1 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~soil~~ group for the soil to be treated (soil texture).

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

***Use rates listed in this label using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.

****Do not use this mixture on sand or loamy sand soils with less than 2% organic matter.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

24

corn insert

Preplant Incorporated Application Rates

BROADCAST RATE PER ACRE					
Less than 3% organic matter					
SOIL TEXTURAL GROUP*	Lasso [†]	atrazine 4L**	atrazine 4L**	Bladex 4L***	Bladex 4L***
	(pounds)	(quarts)	(quarts)	(quarts)	(quarts)
Coarse****	2.6 to 3.4	1		0.5 to 1	
Medium	2.6 to 3.4	1	to 1.25	1	to 1.25
Fine	3.4 to 3.8	1.25	to 1.5	1	to 1.5

BROADCAST RATE PER ACRE					
3% or more organic matter					
SOIL TEXTURAL GROUP*	Lasso [†]	atrazine 4L**	atrazine 4L**	Bladex 4L***	Bladex 4L***
	(pounds)	(quarts)	(quarts)	(quarts)	(quarts)
Coarse****	2.6 to 3.4	1	to 1.25	1	to 1.25
Medium	3.4 to 3.8	1.25	to 1.5	1	to 2
Fine	3.4 to 4.1	1.5	to 1.75	1	to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~soil~~ group for the soil to be treated. **soil texture**

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

***Use rates listed in this label using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.

****Do not use this mixture on sand or loamy sand soils with less than 2% organic matter.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

LASSO® plus ATRAZINE plus PROWL™
(TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of triazine resistant lambsquarter and pigweed in addition to yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded
Aerial - Fixed wing; helicopter.

Reference: The "APPLICATION SYSTEMS" section of this label provides information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHOD

Preemergence Surface: Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

DO NOT preplant incorporate this tank mixture as serious crop injury can result.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: PLANT ONLY CORN, PEANUTS, SORGHUM OR SOYBEANS* the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn. DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

Recommended Rates: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	Less Than 3% Organic Matter		
	Lasso® (Pounds)	+ Atrazine 4L** (Quarts)	+ Prowl (Quarts)
Coarse	2.3 to 3	0.75 to 2	0.75
Medium	2.3 to 3	1 to 2	0.75 to 1
Fine	3 to 3.4	1.25 to 2	1

20 4 60
corn insert

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	3% Or More Organic Matter		
	Lasso® (Pounds)	Atrazine 4L** (Quarts)	Prowl (Quarts)
Coarse	2.3 to 3	1 to 2	0.75 to 1
Medium	3 to 3.4	1.25 to 2	1
Fine	3 to 4.1	1.25 to 2	1

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

™Prowl is a trademark of American Cyanamid Corporation.

LASSO® plus BLADEx plus PROWL
(TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of triazine resistant lambsquarter and pigweed in addition to yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded
Aerial - Fixed wing; helicopter

Reference: The "APPLICATION SYSTEMS" section of this label provides information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHOD

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

DO NOT preplant incorporate this tank mixture as serious crop injury can result.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

Recommended Rates: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	1% to 3% Organic Matter		
	Lasso® (pounds)	+ Bladex 4L(quarts)**	+ Prowl (quarts)
Coarse**	2.3 to 3	1 to 1.2	0.75
Medium	2.3 to 3	1.2 to 1.4	0.75 to 1
Fine	3 to 3.4	1.4 to 1.8	1

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	3% or More Organic Matter		
	Lasso (pounds)	+ Bladex 4L(quarts)**	+ Prowl (quarts)
Coarse**	2.3 to 3	1.2 to 1.6	0.75 to 1
Medium	3 to 3.4	1.6 to 2	1
Fine	3 to 4.1	1.8 to 2.2	1

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Do not use this tank mixture on sand or loamy sand soils with less than 2% organic matter.

***Use rates listed in this label when using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 lbs of Bladex 80W.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

LASSO® plus ATRAZINE plus LINURON
(TANK MIXTURE)

FOR USE ONLY IN DELAWARE, MARYLAND, NEW JERSEY, NEW YORK, PENNSYLVANIA,
VIRGINIA AND WEST VIRGINIA.

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of triazine resistant lambsquarter and pigweed in addition to yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Reference: the "APPLICATION SYSTEMS" section of this label provides information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHOD

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: PLANT ONLY CORN, PEANUTS, SORGHUM, OR SOYBEANS* the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn. DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

RECOMMENDED RATES. Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	Less Than 3% Organic Matter		
	Lasso® (Pounds)	Atrazine 4L** (Quarts)	Linuron 4L*** (Quarts)
Coarse - sandy			
loam only	2.3 to 3	0.75 to 2	0.33
Medium	2.3 to 3	1 to 2	0.5
Fine	3 to 3.4	1.25 to 2	0.5

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE			
	3% Or More Organic Matter			
	Lasso® (Pounds)	+	Atrazine 4L** (Quarts)	+ Linuron 4L*** (Quarts)
Coarse - sandy loam only	2.3	to 3	1 to 2	0.5
Medium	3	to 3.4	1.25 to 2	0.5
Fine	3	to 4.1	1.25 to 2	0.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

***Use the rates listed in this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50% dry flowable formulations. One quart of linuron 4L equals 2 pounds of linuron 50WP or 2 pounds of linuron 50% dry flowable.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

90046

**Conservation or Minimum
Tillage Systems
TANK MIXTURES**

**FIELD CORN OR
SILAGE CORN ONLY**

(omit)

AT PLANTING APPLICATIONS

When applied as directed under the conditions described, these tank mixtures control many emerged annual weeds, suppress many emerged perennial weeds and give preemergence control of many annual grasses and weeds when corn will be planted directly into a cover crop, established sod or in previous crop residues. These tank mixtures will not control regrowth from perennial weeds.

DO NOT APPLY BY AIR

For mixing instructions, see the "MIXING AND SPRAYING INSTRUCTIONS" section of this label

Refer to specific product labels for crop rotation and cautionary statements of all products used in the tank mixture.

■ LASSO[®] plus GRAMOXONE SUPER ■

or

■ LASSO plus ATRAZINE plus GRAMOXONE SUPER ■

or

■ LASSO plus BLADEX plus GRAMOXONE SUPER ■

or

■ LASSO plus PRINCEP[™] plus GRAMOXONE SUPER ■

Apply these tank mixtures with Gramoxone Super in 20 to 60 gallons of water spray solution per acre immediately before, during or after planting, but BEFORE CROP EMERGENCE. As density of stubble, crop residue or weeds increases, spray gallonage and rate should be increased within the recommended ranges to insure complete coverage. In the absence of emerged vegetation, the Gramoxone Super portion of these tank mixtures

PRINCEP is a trademark of Ciba-Geigy Corporation.

or
LASSO plus ATRAZINE plus
BLADEX plus GRAMOXONE SUPER

or
LASSO plus ATRAZINE plus
PRINCEP plus GRAMOXONE SUPER

CONTROL OR SUPPRESSION OF EMERGED WEEDS

GRAMOXONE SUPER

When used as directed, Gramoxone Super in tank mixture controls many emerged annual weeds and suppresses many emerged perennial weeds.

Broadcast Treatment—Apply 1.5 to 2.5 pints of Gramoxone Super ~~4.25 to 0.5 lb. active ingredient~~ in these tank mixtures immediately before, during or after planting but **BEFORE CROP EMERGENCE**. Use the 2.5 pints rate if any weeds are 4 to 6 inches tall. This mixture may not control weeds taller than 6 inches. As density of stubble crop residue or weeds increases, spray gallonage should

be increased within the recommended range for complete coverage. Add a nonionic spreader-surfactant (approved for use on crops) containing at least 75 percent surfactant active agent at 8 ounces per 100 gallons of diluted spray. Refer to the Gramoxone Super label for precautionary statements.

(all caps)

PREEMERGENCE WEED CONTROL

LASSO*

For weeds controlled preemergence, see the sections of this label on Lasso. See the following table for recommended rates of Lasso in this tank mixture on various soil types:

BROADCAST RATE PER ACRE (POUNDS)	
SOIL TEXTURAL GROUP*	Lasso®
Coarse	3.8
Medium	4.5
Fine	4.5 to 5.3

*Refer to the "SOIL TEXTURE" section of the label to determine the corresponding ~~soil~~ group for the soil to be treated. **soil textural**

Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestation or when organic matter content is 3 percent or more. ~~The rate of Lasso may be increased to a maximum of 10.0 lbs. per acre on any soil type to reduce competition with the crop to control weeds listed on this label.~~

LASSO* plus ATRAZINE

For weeds controlled preemergence, see the sections of this label for Lasso and Lasso plus atrazine.

Refer to the following table for recommended rates of Lasso plus Bladex in this tank mixture on various soil types

BROADCAST RATE PER ACRE		
SOIL TEXTURAL GROUP*	Lasso+ (pounds)	+ Bladex 4L** (quarts)
Coarse—sandy		
loam only	2.6 to 3	1 to 2
Medium	3 to 3.8	1.2 to 2
Fine	3.8 to 4.5	1.6 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~soil texture~~ group for the soil to be treated. **soil texture**

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present. ~~The rate of Lasso may be increased to the maximum rate on any soil type to reduce competition from the weeds to control weeds listed in this label.~~

Use the higher rate of atrazine in the recommended ranges on soils with greater than 3% organic matter.

NOTE: PLANT ONLY CORN, PEANUTS, SORGHUM OR SOYBEANS the year following use of this mixture.

There is a possibility of crop injury due to carryover of atrazine if soybeans are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

■ LASSO® plus BLADEX ■

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of the label for Lasso and this tank mixture.

See the following table for recommended rates of Lasso plus Bladex in this tank mixture with Roundup on various soil types.

BROADCAST RATE PER ACRE		
SOIL TEXTURAL GROUP*	Lasso+ (pounds)	+ Bladex 4L** (quarts)
Coarse—sandy		
loam only	2.6 to 3	1 to 1.6
Medium	3 to 3.8	1.2 to 1.6
Fine	3.8 to 4.5	1.6 to 2.2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~soil texture~~ group for the soil to be treated. **soil texture**

**When using Bladex 80W use equivalent rates. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.

Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present. ~~The rate of Lasso may be increased to the maximum rate on any soil type to reduce competition from the weeds to control weeds listed in this label.~~

Use the higher rate of Bladex in the recommended ranges on soils with greater than 3% organic matter.

NOTE: Do not use this mixture on sand or loamy sand soils with less than 2% organic matter.

corn insert

■ LASSO[®] plus PRINCEP ■

For weeds controlled preemergence see the "WEEDS CONTROLLED" section of the labels for Lasso and Princep herbicides.

See the following table for recommended rates of Lasso plus Princep in this tank mixture on various soil types.

BROADCAST RATE PER ACRE

SOIL TEXTURAL GROUP*	Lasso [®] (pounds)	+ Princep 4L** (quarts)
Coarse—sandy		
Sandy only	2.6 to 3	1 to 1.2
Medium	3 to 3.8	1.2 to 1.6
Fine	3.8 to 4.5	1.6 to 2.2

Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~texture~~ group for the soil to be treated **soil textural**

**When using Princep 80W use equivalent rates. One quart of Princep 4L equals 1.25 pounds of Princep 80W

Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present.

Use the higher rate of Princep in the recommended ranges on soils with greater than 3% organic matter

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR AFTER HARVEST OF TREATED CROP. PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS

BEST AVAILABLE COPY

LASSO® plus ATRAZINE plus BLADEX

This tank mixture maybe applied for additional preemergence control of heavy weed infestation and/or hard-to-control weeds listed in the "WEEDS CONTROLLED" section of this label. Refer to the following rate table to determine the recommended rates for this tank mixtures on various soil types.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	Lasso® (Pounds)	+ Atrazine 4L** (Quarts)	+ Bladex 4L*** (Quarts)
Coarse****	2.6 to 3	0.75 to 1	1 to 1.25
Medium	3 to 3.8	1 to 1.25	1 to 2
Fine	3.4 to 4.5	1 to 1.25	1 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

***Use rates listed in this label when using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.

****DO NOT use this mixture on sand or loamy sand soils with less than 2% organic matter.

Use the higher rates in the recommended ranges in area of heavy weed infestations.

NOTE: PLANT ONLY corn, peanuts, sorghum or soybeans* the year following use of this mixture.

*There is a possibility of crop injury due to carryover of atrazine if soybeans are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

■ LASSO[®] plus ATRAZINE plus PRINCEP ■

For additional preemergence control of heavy weed infestation and/or hard-to-control weeds, a three-way tank mixture of Lasso herbicide plus atrazine plus Princep may be applied. Refer to the following rate table to determine the recommended rates for ~~these~~ tank mixtures on various soil types.

this

SOIL TEXTURAL GROUP ^a	BROADCAST RATE PER ACRE		
	Lasso ^b (pounds)	atrazine 4L ^c (quarts)	Princep 4L ^d (quarts)
Coarse	2.6 to 3	0.75 to 1	0.75 to 1
Medium	3 to 3.8	1 to 1.25	1 to 1.5
Fine	3.8 to 4.5	1 to 1.25	1 to 1.5

Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil ~~texture~~ group for the soil to be treated.

^aWhen using 4L formulations use the rates listed. Use equivalent rates when using EW or 90% dry flowable formulations. One quart of the 4L formulation of these products equals 1.25 pounds of the 80W formulation. One quart of atrazine 4L or Princep 4L equals 1.1 pounds of the 90% dry flowable formulations of these products.

Use the higher rates of atrazine and Princep in the recommended ranges on soils with greater than 3% organic matter.

In the areas of heavy grass infestation, use the higher rate of Lasso in the recommended ranges when tall panicum, crabgrass, or soils with greater than 3% organic matter will be present. The rate of Lasso may be increased to ~~6~~

~~to control weeds on any soil type to reduce competition from the hard-to-control weeds listed in this label.~~

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR AFTER HARVEST OF TREATED CROP. PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING-PLANTED CROPS. PLANT ONLY CORN, PEANUTS, SORGHUM OR SOYBEANS THE YEAR FOLLOWING USE OF THIS MIXTURE.

There is a possibility of crop injury due to carryover of atrazine if soybeans are planted the following year.

DO NOT plant soybeans the year following use of this tank mixture on land not registered corn.

DO NOT graze treated areas or feed treated forage to livestock for 21 days following application of this tank mixture.

BEST AVAILABLE COPY

EARLY PREPLANT APPLICATION

LASSO® PLUS ATRAZINE (TANK MIXTURE)

3726
corn insert

For use in no-till and other conservation tillage systems in DELAWARE, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MARYLAND, MINNESOTA, MISSOURI, NEBRASKA, NORTH CAROLINA, OHIO, SOUTH DAKOTA, VIRGINIA, AND WISCONSIN.

This tank mixture, when applied in a single application (alone or in a 3 way combination with Bladex or Princep), split application or as a sequential application to Bladex or Princep in early preplant programs, will provide pre-emergence control or reduced competition of listed annual grasses and broadleaf weeds.

If emerged weeds are present at the time of treatment Gramoxone Super or 2,4-D should be added to the product according to the directions for use on their respective product labels. If unsatisfactory weed control occurs (due to excessively dry or excessively wet conditions) following the earlier application, a postemergence application of an appropriate labelled grass and/or broadleaf weed herbicide may be used. If a postemergence treatment includes the herbicide used early preplant, do not exceed the labelled rate for corn on a given soil texture. Observe all precautions and limitations on the labels for Lasso, atrazine, Bladex, Princep, Gramoxone Super, 2,4,-D and other post-emergence herbicides before use of these products.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application - Application of this tank mixture should be made less than 30 days before planting but prior to weed emergence. This application should be restricted to areas north of Interstate 70 when west of the Mississippi River and north of Interstate 64 when east of the Mississippi River where cool spring weather minimizes or slows biodegradation of herbicides.

See the following table for recommended rates.

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE	
	Lasso® (pounds)**	atrazine 4L (quarts)**
Coarse*	3.4 to 3.8	1.25 to 1.5
Medium	3.8 to 4.5	1.5 to 2
Fine	4.5 to 5.6	1.75 to 2.25

This tank mixture may also be mixed in a 3-way combination with 1 to 1.25 quarts per acre of Princep or 1.5 to 2 quarts per acre of Bladex to provide improved control of fall panicum and crabgrass.

*Applications on coarse soils should not be made more than 2 weeks prior to planting.

**Use the lower rate where light weed infestation is expected and the higher rate where heavy weed infestation is expected.

Split application - Apply 60 percent of the recommended rate as a split treatment prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting.

FOR CONTROL OF FALL PANICUM, CRABGRASS OR BROADLEAF SIGNALGRASS USE THE
SEQUENTIAL APPLICATION OF THIS TANK MIXTURE FOLLOWING BLADEX OR PRINCEP. corn insert

See the following table for recommended rates.

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE	
	Lasso®* (pounds)*	+ atrazine 4L (quarts)*
Coarse	3.4 to 4.1	1.25 to 1.75
Medium	3.8 to 4.5	1.5 to 2
Fine	4.5 to 5.6	1.75 to 2.25

*Use the lower rate where light weed infestation is expected and the higher rate where heavy weed infestation is expected.

Sequential application - Application of this tank mixture following Bladex or Princep should be utilized for the control of fall panicum, crabgrass or broadleaf signalgrass. Apply 1.5 to 2 quarts per acre of Bladex or 1 to 1.25 quarts per acre of Princep prior to weed emergence and no more than 45 days prior to planting. At or immediately following planting apply the recommended rate of this tank mixture.

Following applications of Bladex or Princep see the following table for recommended rates.

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE	
	Lasso® (pounds)*	+ atrazine 4L (quarts)*
Coarse	2.6 to 3.8	1 to 1.5
Medium	3.4 to 4.1	1.25 to 1.75
Fine	3.8 to 4.5	1.5 to 2

*Use the lower rate where light weed infestation is expected and the higher rate where heavy weed infestation is expected.

Use rates listed in this label when using the 4L formulation of atrazine, Bladex or Princep. Use equivalent rates of the 30W or 90% dry flowable formulation of atrazine, Bladex or Princep. One quart of the 4L formulation of atrazine, Bladex or Princep equals 1.25 pounds of the 80W and 1.1 pounds of the 90% dry flowable formulations of atrazine, Bladex or Princep.

DO NOT use Bladex on coarse textured soils or soils of less than 1 percent organic matter.

Plant only corn, peanuts, sorghum (milo) or soybeans the year following the use of this program. There is a possibility of crop injury due to carryover of atrazine if soybeans are planted the following year.

NOTE: Soil treated with Princep should not be planted to any crop except corn for one year following treatment as crop injury may occur. After harvest of the treated crop, plow and thoroughly till the soil in the fall or spring to minimize possible injury to spring seeded rotational crops.

SOYBEANS

LASSO®

Apply this product in water or sprayable fluid fertilizer solution for control of yellow nutsedge, and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Aerial — Fixed wing; helicopter

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface — Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE (POUNDS)**		TEXTURAL		
	Less than 3% organic matter	3% or more organic matter			
Coarse***	3 to 3.4	3 to 3.4	3	to	3.4
Medium	3 to 4.1	3 to 4.1	3	to	4.1
Fine	3 to 4.1	3 to 4.1	3	to	4.1

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

soil
textural

**Use the higher rate in the recommended range in areas of heavy weed infestations. Use a minimum of 3.75 pounds of this product per acre on coarse soils and 4.5 to 6 pounds on medium or fine textured soils to control eastern black or hairy nightshade or to reduce competition from hard-to-control weeds listed in this label.

***When applied under sprinkler irrigation systems on coarse soils, use a minimum of 4.5 pounds per acre of this product.

On coarse soils with 10% or more organic matter use 6 pounds of Lasso per acre.

Preplant Incorporated Application Rates

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE (POUNDS)**		TEXTURAL
	Less than 3% organic matter	3% or more organic matter	
Coarse	3.75	4.75	3.8 3.8 to 4.5 3.8 to 4.5
Medium	3.75 to 4.5	3.75 to 5.25	
Fine	3.75 to 4.5	4.5 to 6	

soil textural

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the higher rate in the recommended range in areas of heavy weed infestations. Use a minimum of 4.5 pounds per acre on coarse textured soils for control of yellow nutsedge or red rice. Rates may be increased to a maximum of 6 pounds on any soil type when heavy infestations of yellow nutsedge or red rice are present ~~to reduce competition from the hard to control weeds listed in this label.~~

On coarse soils with 10% or more organic matter use 6 pounds of Lasso per acre.

**LASSO® plus AMIBEN
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid for trigger solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

DO NOT use this tank mixture on sand or loamy sand soils as crop injury may result.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom, banded

Aerial — Fixed wing, helicopter

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface — Apply after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES:

Preplant Incorporated — Apply 3.75 to 4.5 pounds of Lasso plus 4 quarts of Amiben per acre.

NOTE Use the 4.5 pound per acre rate of Lasso in areas of heavy weed infestation. Use 4.5 to 6 pounds of Lasso when eastern black or hairy nightshade, yel-

3.8

41 7 60

low nutsedge or red rice is present ~~go to reduce competition from the hard-to-control weeds listed in this label.~~

Preemergence Surface — Apply 3 pounds Lasso plus 4 quarts Amben per acre.

3.8

NOTE: Use a minimum of 3.75 pounds of Lasso on coarse soils and 4.5 to 6 pounds in this tank mixture on medium or fine textured soils to control black or hairy nightshade ~~go to reduce competition from hard-to-control weeds listed in this label.~~

LASSO® plus CANOPY (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Lasso and Canopy labels including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

DO NOT plant rice or peanuts within 18 months after application of this tank mixture.

APPROVED APPLICATION SYSTEM

Ground — Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of the label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight soybean injury early, but soybeans should recover with no subsequent effect on growth or yield.

Preemergence Surface — Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

BROADCAST RATES PER ACRE**

SOIL TEXTURE GROUP*	SOIL ORGANIC MATTER		Lasso® (pounds)	Canopy (quarts)
	1/2 to 3%	3% or more		
Coarse	2.25 to 3	2.25 to 3.5	6 to 10	
Medium	2.25 to 3.75	3 to 4.5	8 to 13	
Fine	3 to 4.5	3 to 6.25	10 to 14	

TEXTURAL

Coarse- sandy loam only	2.6 to 3.4	2.6 to 3.4	6
Medium	2.6 to 3.4	3 to 3.8	8
Fine	3 to 3.8	3.4 to 4.1	10

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

**Use the higher rate of Lasso and Canopy in areas of heavy weed infestation.

** Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present.

Preplant Incorporated Application Rate

BROADCAST RATES PER ACRE**

SOIL TEXTURE GROUP*	organic matter		TEXTURAL	
	1/4 to 3%	3% or more		
	Lasso [®] (pounds)	Command (ounces)		
Coarse	2.25 to 3.75	3 to 5	Coarse - sandy loam	
Medium	2.25 to 3.75	3 to 5	only	3 to 3.8
Fine	3 to 4.5	5 to 10	Medium	3 to 3.8
			Fine	3.4 to 4.1
				3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group of the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/4% organic matter.

**Use the higher rate of Lasso and Command in areas of heavy weed infestation.

LASSO[®] plus COMMAND (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Lasso and Command labels before use of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 3 to 3.75 pounds of Lasso plus 1 pint of Command 4E per acre. In areas of heavy pigweed and lambsquarter infestations, use 3.75 to 4.5 pounds of Lasso per acre in the tank mixture.

LASSO[®] plus GEMINI (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Lasso and Gemini labels including precautions on soil pH, minimum recropping interval and rotational guidelines.

DO NOT plant rice or peanuts within 18 months after application of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface — Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

** Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. See "WEEDS CONTROLLED" section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.

00058

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURE GROUP*	BROADCAST RATES PER ACRE**			
	1/2 to 3% organic matter		3% or more organic matter	
	Lasso (pounds)	Gemini (ounces)	Lasso (pounds)	Gemini (ounces)
Coarse - sandy loam only	2.25 to 3	12 to 16	2.25 to 3.5	12 to 16
Medium	2.25 to 3.75	16 to 20	3 to 4.5	16 to 20
Fine	3 to 4.5	20 to 24	3 to 5.25	20 to 24

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture for the soil to be treated.

**Use the higher rate of Lasso and Gemini in areas of heavy weed infestation.

LASSO® plus LEXONE

or

**LASSO plus SENCOR
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: DO NOT use on much soils or on alkaline soils with a pH greater than 7.4. Crop injury may result if atrazine was applied the year preceding use of this tank mixture.

DO NOT REPLANT CROPS OTHER THAN SOYBEANS FOR 120 DAYS AFTER APPLICATION OF THESE TANK MIXTURES.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; hand

Aerial — Fixed wing; helicopter

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight soybean injury early, but soybeans should recover with no subsequent effect on growth or yield.

Preemergence Surface — Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*

1/2 to 3% organic matter

3% or more organic matter

Lasso (pounds)

Lasso (pounds)

Gemini (ounces)

Coarse - sandy loam

only

2.6 to 3.4

2.6 to 3.4

12 to 16

Medium

2.6 to 3.4

3 to 3.8

16 to 20

Fine

3 to 3.8

3.4 to 4.1

20 to 24

soil textural

** Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. Use the higher rate of Gemini in the recommended ranges on soils with higher organic matter or heavy pressure from large deep germinating weed seeds.

Surface Application Rates

BROADCAST RATES PER ACRE

SOIL TEXTURE GROUP*	Less than 3% organic matter		3% or more organic matter	
	Lasso [®] 4L** or Sencor 4F** (pounds)	Lasso [®] 4L** or Sencor 4F** (pint)	Lasso [®] 4L** or Sencor 4F** (pounds)	Lasso [®] 4L** or Sencor 4F** (pint)
Coarse - sandy loam only	2.6 to 3.4	0.5	2.6 to 3.4	0.75
Medium	2.6 to 3.4	0.5 to 0.75	3 to 3.8	0.75
Fine	3 to 3.8	0.75 to 1	3.4 to 4.1	1

TEXTURAL

soil
textural

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the rates listed in this label when using Lasso 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lasso 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

NOTE: DO NOT use on sand or loamy sand soil. Use 1.5 ~~to 2~~ pints of Lasso 4L or Sencor 4F per acre in this tank mixture when applying on ~~heavy clay~~ heavy clay soils of the Mississippi Delta. Use a minimum of 2.75 pounds of Lasso on coarse soils and 4.5 to 6 pounds in this tank mixture on medium or fine textured soils to control eastern black or hairy nightshade or to ~~reduce competition from heavy infestations of the~~ hard-to-control weeds listed in this label.

3.8

Preplant Incorporated Application Rates

BROADCAST RATES PER ACRE

SOIL TEXTURE GROUP*	Less than 3% organic matter		3% or more organic matter	
	Lasso [®] 4L** or Sencor 4F** (pounds)	Lasso [®] 4L** or Sencor 4F** (pint)	Lasso [®] 4L** or Sencor 4F** (pounds)	Lasso [®] 4L** or Sencor 4F** (pint)
Coarse - sandy loam only	3 to 3.8	0.5	3 to 3.8	0.75
Medium	3 to 3.8	0.5 to 0.75	3.4 to 4.1	0.75
Fine	3.4 to 4.1	0.75 to 1	3.8 to 4.5	1

TEXTURAL

Coarse - sandy loam

only

Medium

Fine

soil
textural

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the rates listed in this label when using Lasso 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lasso 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

***Use the 4.5 pounds per acre rate of Lasso in areas of heavy weed infestation. Use 4.5 to 6 pounds of Lasso when eastern black or hairy nightshade, yellow nutsedge or red rice is present ~~to reduce competition from the~~ hard-to-control weeds listed in this label.

NOTE: Do not use on coarse textured soils. Do not use on medium or fine textured soils with less than 1% organic matter. These mixtures, preplant incorporated, may cause slight soybean injury early, but the soybeans should recover and subsequent growth or yield should not be affected.

90060

45 60

LASSO® plus AMIBEN plus LEXONE or SENCOR (TANK MIXTURES)

Apply this tank mixture in water solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: DO NOT use on sand, loamy sand, or on soils with less than 1% organic matter. DO NOT use the foliage from soybeans treated with Lasso plus Amiben plus Lexone or Sencor for feed or forage.

NOTE: DO NOT use on much soils or on alkaline soils with greater than 7.4 pH. Crop injury may result if atrazine was applied the year preceding use of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom

Aerial — Fixed wing, helicopter

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporation — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight soybean injury early but soybeans should recover with no subsequent effect on growth or yield.

Preemergence Surface — Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing methods selected.

RECOMMENDED RATES: Refer to the following tables for recommended broadcast treatment rates. Applications which are not constant with recommendations in this label may result in unsatisfactory weed control or crop injury.

BROADCAST RATE PER ACRE
Less than 3% organic matter

SOIL TEXTURE GROUP*	Lexone 4L**			TEXTURAL
	Lasso® (pounds)	Amiben (quarts)	SenCor 4F (pint)	
Coarse — sandy loam	2.5 to 3	3	0.5	{ 2.6 to 3.4 2.6 to 3.4 3 to 3.8
only	2.5 to 3	3	0.5	
Medium	2.5 to 3.75	3 to 4	0.5	
Fine	3 to 3.75	4 to 5	0.75	

BROADCAST RATE PER ACRE
3% or more organic matter

SOIL TEXTURE GROUP*	Lexone 4L**			TEXTURAL
	Lasso® (pounds)	Amiben (quarts)	SenCor 4F (pint)	
Coarse — sandy loam	2.5 to 3	3 to 4	to 0.75	{ 2.6 to 3.4 3 to 3.8 3.4 to 4.1
only	2.5 to 3	3 to 4	to 0.75	
Medium	3 to 4.5	4 to 5	0.5 to 0.75	
Fine	3 to 4.5	4 to 6	0.75 to 1	

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated. On alkaline (calcareous) soils with a pH of 7.4 or greater, apply Lexone 4L or SenCor 4F at 0.5 pint/A on medium soils and 0.5 to 0.75 pint or corresponding equivalent of WP or DF formulations on fine soils. Use the higher rates of Lasso in the recommended ranges in areas of heavy weed infestations and for preplant incorpora-

soil
textural

46860

tion applications. Use a minimum of 3.75 pounds of Lasso on coarse soils and 4.5 to 6 pounds in this tank mixture on medium or fine soils, when hard-to-control weeds or eastern black or hairy nightshade are present. Use 4.5 to 6 pounds of Lasso per acre preplant incorporated for control of yellow nutsedge.

3.8

**Use rates listed in this label when using Lasso 4L or Sencor 4F. Use equivalent rates when using the 50WP or DF formulations of these products. One pint of Lasso 4L or Sencor 4F equals 1 pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

**LASSO® plus LINURON
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Plant seed at least 1-3/4 inches deep and DO NOT use on sand or loamy sand or on soils with less than 1% organic matter as crop injury may occur. DO NOT spray over the top of emerged soybeans.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

~~Aerial — Fixed-wing; helicopter~~

(omit)

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface — Apply after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

BROADCAST RATES PER ACRE				
SOIL TEXTURE GROUP*	1% to 3% organic matter		3% or more organic matter	
	Lasso® + Imazapyr® 4L (pounds)	4L (quarts)	Lasso + Imazapyr® 4L (pounds)	4L (quarts)
Coarse— sandy loam				
only	2.25 to 3	0.5	2.25 to 3	0.5 to 0.75
Medium	2.25 to 3.4	0.5 to 0.75	2.25 to 3.4	0.75 to 1
Fine	2.25 to 3.4	1	2.25 to 3.4	1.25 to 1.5

TEXTURAL

SOIL
textural

2.6 to 3.4	2.6 to 3.4
2.6 to 3.4	3 to 3.8
3 to 3.8	3.4 to 4.1

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the rates listed on this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50% dry flowable formulations. One quart of linuron 4L equals 2 pounds of linuron 50WP or 2 pounds of linuron 50% dry flowable.

NOTE: Use a minimum of 3.75 pounds per acre of Lasso on coarse textured soils and use 4.5 to 6 pounds on medium and fine textured soils to control eastern black or hairy nightshade or to reduce competition from the hard-to-control weeds listed in this label.

LASSO® plus LOROX plus (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Lasso and Lorox Plus labels including precautions on soil pH, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface — Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label contains detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**			
	1/2 to 3% organic matter		3% or more organic matter	
	Lasso (pounds)	Lorox Plus (ounces)	Lasso (pounds)	Lorox Plus (ounces)
Coarse - sandy loam only	2.6 to 3.4	12 to 14	2.6 to 3.4	12 to 14
Medium	2.6 to 3.4	14 to 16	3 to 3.8	14 to 16
Fine	3 to 3.8	16 to 18	3.4 to 4.1	16 to 18

SOIL TEXTURAL GROUP*

Coarse - sandy loam

only

Medium

Fine

BROADCAST RATE PER ACRE**

1/2 to 3% organic matter

3% or more organic matter

Lasso (pounds)

Lasso (pounds)

Lorox Plus (ounces)

soil textural

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the higher rate of Lasso and Lorox Plus in areas of heavy weed infestation.

LASSO® plus PREVIEW (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Lasso and Preview label including precautions of soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of the label provides detailed information on the proper application of this tank mixture using the system selected.

** Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. Use the higher rate of Lorox Plus in the recommended ranges on soils with higher organic matter or heavy pressure from large deep germinating weed seeds.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight or severe injury early, but soybeans should recover with no subsequent effect on growth or yield.

Preemergence Surface — Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

BROADCAST RATES PER ACRE**

SOIL TEXTURE GROUP*	1/4 to 3% organic matter		3% or more organic matter	
	Lasso® (pounds)	Prevex (ounces)	Lasso (pounds)	Prevex (ounces)
Coarse	2.25 to 3	6	2.25 to 3	7
Medium	2.25 to 2.75	7	2.25 to 2.5	8
Fine	2.25 to 2.5	8	2.25 to 2.5	9

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated. DO NOT apply to soil with less than 1/4% organic matter.

**Use the higher rate of Lasso and Prevex in the rate range in areas of heavy weed infestation.

Preplant Incorporated Application Rates

BROADCAST RATES PER ACRE**

SOIL TEXTURE GROUP*	1/4 to 3% organic matter		3% or more organic matter	
	Lasso® (pounds)	Prevex (ounces)	Lasso (pounds)	Prevex (ounces)
Medium	2.25 to 2.5	7	2.25 to 2.5	8
Fine	2.25 to 2.5	8	2.25 to 2.5	9

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group of the soil to be treated. DO NOT apply to soil with less than 1/4% organic matter.

**Use the higher rate of Lasso and Prevex in the rate range in areas of heavy weed infestation.

LASSO® plus SCEPTER (TANK MIXTURE)

Apply this tank mixture in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

Note: Observe all precautions and limitations on the Lasso and Scepter labels before use of this tank mixture including precautions on minimum mowing interval and rotational guidelines.

DO NOT apply this tank mixture in sprayable fluid form.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Aerial — Fixed wing helicopter

TEXTURAL

Coarse — sandy loam

only 2.6 to 3.4 6 2.6 to 3.4 7

Medium 2.6 to 3.4 7 3 to 3.8 8

Fine 3 to 3.8 8 3.4 to 4.1 9

soil
textural

** Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present.

TEXTURAL

Coarse — sandy loam

only 3 to 3.8 6 3 to 3.8 7

Medium 3 to 3.8 7 3.4 to 4.1 8

Fine 3.4 to 4.1 8 3.8 to 4.5 9

soil
textural

** Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. See "WEEDS CONTROLLED" section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.

or sprayable fluid fertilizer

(omit)

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface — Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 2.6 to 3.75 pounds of Lasso plus 2/3 pint of Scepter per acre. Use the higher rate of Lasso in the recommended high organic matter, fine-textured soils. See the "WEEDS CONTROLLED" section of this label for recommended rates and application methods for control of certain weeds.

LASO® plus TREFLAN (TANK MIXTURE)

Apply this tank mixture in water solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: See The "MIXING, SPRAYING AND HANDLING INSTRUCTION" section of this label for special mixing directions for Lasso plus Treflan.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Aerial — Fixed wing; helicopter

Reference: The "APPLICATION SYSTEMS" section of the label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil within 8 hours after application.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES: Preplant Incorporated — Apply 3.75 to 4.5 pounds of Lasso plus 0.5 quart of Treflan per acre. Applications which are not consistent with the recommendation in this label may result in unsatisfactory weed control or crop injury.

~~BASAGRAN SEQUENTIAL TREATMENT~~

~~Goybeans Only~~

~~Basagran may be applied following preemergence applications of Lasso or tank mixtures of Lasso on this label for control of emerged weeds such as crabgrass, velvet leaf and other weed species listed on the label for Basagran. Read and follow the cautionary statements, directions for use and all other information appearing on the label for Basagran.~~

~~**RECOMMENDED RATES:** Apply 0.75 to 1.5 quarts of Basagran as directed on the label for Basagran.~~

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	Less than 3% organic matter	3% or more organic matter	Scepter*** (pints)
	Lasso (pounds)	Lasso (pounds)	
Coarse	2.3 to 3	2.3 to 3	1/2 to 2/3
Medium	2.3 to 3	2.6 to 3.4	1/2 to 2/3
Fine	2.6 to 3.4	3 to 3.8	1/2 to 2/3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when the following weeds are present: fall panicum, barnyardgrass, broadleaf signalgrass, seedling johnsongrass, red rice or yellow nutsedge.

*** Use the higher rate of Scepter in the recommended range when red rice, sicklepod, beggarweed or velvetleaf are present.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command 4E herbicide per acre to this tank mixture and preplant incorporate. Observe all precautions and limitations on the Scepter and Command labels before use of this tank mixture.

3.8

Conservation or Minimum
Tillage Systems
TANK MIXTURES

SOYBEANS ONLY

When applied as directed under the conditions described, these tank mixtures control many emerged annual weeds, suppress many emerged perennial weeds and give preemergence control of many annual grasses and weeds when soybeans will be planted directly into a cover crop, stale seedbed, or in previous crop residues such as wheat stubble.

These tank mixtures will not control regrowth from perennial weeds.

DO NOT APPLY BY AIR.

For mixing instructions, see the "MIXING, SPRAYING AND HANDLING INSTRUCTIONS" section of this label.

- LASSO® plus GRAMOXONE SUPER ■
- or
- LASSO plus CANOPY plus GRAMOXONE SUPER ■
- or
- LASSO plus GEMINI plus GRAMOXONE SUPER ■
- or
- LASSO plus LEXONE plus GRAMOXONE SUPER ■
- or
- LASSO plus LINURON plus GRAMOXONE SUPER ■
- or
- LASSO plus LOROX plus GRAMOXONE SUPER ■
- or
- LASSO plus PREVIEW plus GRAMOXONE SUPER ■
- or
- LASSO plus SCEPTER plus GRAMOXONE SUPER ■
- or
- LASSO plus SENCOR plus GRAMOXONE SUPER ■

Apply these tank mixtures in 20 to 60 gallons of water per acre immediately before, during or after planting, but BEFORE CROP EMERGENCE. As density of stubble, crop residue or weeds increases, spray gallonage and rate should be increased within the recommended ranges to insure complete coverage. In the absence of emerged vegetation delete the Gramoxone Super portion of these mixes.

CONTROL OR SUPPRESSION OF EMERGED WEEDS

- GRAMOXONE SUPER ■

When used as directed, Gramoxone Super tank-mixed with labeled Lasso tank mixtures controls many emerged annual weeds and suppresses many emerged perennial weeds.

Broadcast Treatment — Apply 1.5 to 2.5 pints of Gramoxone Super in this tank mixture immediately before, during or after planting but BEFORE CROP EMERGENCE. Use the 2.5 pint rate if any weeds are 4 to 6 inches tall. This mixture may not control weeds taller than 6 inches. As density of stubble, crop residue or weeds increases, spray volume should be increased within the recommended range for complete coverage. Add a nonionic spreader surfactant (approved for use in growing crops) containing at least 75 percent surface active agent at 8 ounces per 100 gallons of diluted spray. Refer to the Gramoxone Super label for precautionary statements.

00066

PREEMERGENCE WEED CONTROL

■ LASSO® ■

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of this label.

See the following table for recommended rates of Lasso on various soil types.

BROADCAST RATE PER ACRE (POUNDS)

SOIL TEXTURE GROUP*	Lasso®	TEXTURAL
Coarse	3.75 to 4.5	3.8
Medium	4.5 to 5.25	4.5
Fine	4.5 to 6	4.5 to 5.3

*Refer to the "SOIL TEXTURE" section of the label to determine the corresponding texture group for the soil to be treated.

soil
textural

Use the higher rate in the recommended ranges in areas of heavy grass infestation, or when organic matter content is 3 percent or more. The rate of Lasso may be increased to a maximum of 6 pounds per acre on any soil type to reduce competition from hard-to-control weeds listed in this label.

■ LASSO® plus CANOPY ■

For weeds controlled preemergence see the "WEEDS CONTROLLED" section of this label for Lasso and Lasso plus Canopy.

NOTE: Observe all precautions and limitations on the Lasso and Canopy labels including precautions on soil pH, sensitive soybean varieties and rotational crop restrictions.

See the following table for recommended rates of Lasso and Canopy in this tank mixture.

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE**		TEXTURAL
	Lasso® (pounds)	+ Canopy (ounces)	
Coarse —			
sandy loam only	3 to 3.75	6 to 10	2.6 to 3.4
Medium	3.75 to 4.5	8 to 10	3 to 3.8
Fine	3.75 to 4.5	10 to 14	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

soil
textural

**Use the higher rate of Lasso and Canopy in areas of heavy weed infestation. The rate of Lasso may be increased to a maximum of 6 pounds per acre on any soil type to reduce competition from the hard-to-control weeds listed in this label.

■ LASSO® plus GEMINI ■

For weed controlled preemergence see the "WEEDS CONTROLLED" section of this label for Lasso plus Gemini.

Note: Observe all precautions and limitations on the Lasso and Gemini labels including precautions on soil pH and rotational crop restrictions.

See the following table for recommended rates of Lasso and Gemini in this tank mixture.

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE**		TEXTURAL
	Lasso® (pounds)	+ Gemini (ounces)	
Coarse —			
sandy loam only	2 to 2.75	12 to 16	2.6 to 3.4
Medium	2.75 to 4.5	16 to 20	3 to 3.8
Fine	2.75 to 4.5	20 to 24	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the higher rate of Lasso and Gemini in areas of heavy weed infestation.

~~The rate of Lasso may be increased to a maximum of 6 pounds per acre on any soil type to reduce competition from the hard-to-control weeds listed in this label.~~

LASSO® plus LEXONE

or

LASSO plus SENCOR

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of this label for Lasso and Lasso plus Lexone 4L or Sencor 4F.

See the following table for recommended rates of Lasso plus Lexone 4L or Lasso plus Sencor 4F in this tank mixture on various soil types.

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE		TEXTURAL
	Lasso® (pounds)	+ Lexone 4L** or Sencor 4F** (pints)	
Coarse — sandy			
loam only	2 to 2.75	0.5 to 0.75	2.6 to 3.4
Medium	2.75 to 4.5	0.75 to 1	3 to 3.8
Fine	2.75 to 4.5	1 to 1.5***	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the rates listed in this label when using Lexone 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lexone 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

***On the silty clay or heavy clay soils of the Mississippi Delta, use 1.5 to 2 pints of Lexone 4L or Sencor 4F per acre.

Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestations or when fall panicum or crabgrass will be present. ~~The rate of Lasso may be increased to a maximum of 6 pounds per acre on any soil type to reduce competition from the hard-to-control weeds listed in this label.~~

Use the higher rate of Lexone or Sencor in the recommended ranges on soils with greater than 3% organic matter.

DO NOT use this mixture on sand or loamy sand soils as crop injury from Lexone or Sencor may occur.

DO NOT use on much soils.

53 7 60

DO NOT apply on alkaline soils with a pH greater than 7.4.

Crop injury may occur if atrazine was applied the year before use of this tank mixture.

DO NOT REPLANT CROPS OTHER THAN SOYBEANS FOR 120 DAYS AFTER APPLICATION OF THESE TANK MIXTURES.

■ **LASSO® plus LINURON** ■

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of this label for Lasso and Lasso plus linuron.

See the following table for recommended rates of Lasso and linuron in this tank mixture.

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE (QUARTS)		TEXTURAL
	Lasso® (pounds)	+ Linuron 4L** (quarts)	
Coarse — sandy loam only	3 to 3.75	0.5 to 0.75	2.6 to 3.4
Medium	3.75 to 4.5	0.75 to 1	3 to 3.8
Fine	3.75 to 4.5	1 to 1.5	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of the label to determine the corresponding texture group for the soil to be treated.

soil
textural

**Use the rates listed on this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50 percent dry flowable formulations. One quart of linuron 4L equals 2 pounds of linuron 50WP or 50 percent dry flowable formulation.

Use the higher rate of Lasso in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present. The rate of Lasso may be increased to a maximum of 6 pounds per acre on any soil type to reduce competition from the hard-to-control weeds listed in this label.

Use the higher rate of linuron in the recommended ranges on soils with greater than 3% organic matter.

DO NOT use this tank mixture on sand or loamy sand or on soil with less than 1% organic matter as crop injury from linuron may occur.

■ **LASSO® plus LOROX PLUS** ■

For weeds controlled preemergence see the "WEEDS CONTROLLED" section of this label for Lasso and Lasso plus Lorox Plus.

NOTE: Observe all precautions and limitations on the Lasso and Lorox Plus labels including precautions on soil pH and rotational crop restrictions.

See the following table for recommended rates of Lasso and Lorox Plus in this tank mixture.

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE**		TEXTURAL
	Lasso® (pounds)	+ Lorox Plus (ounces)	
Coarse — sandy loam only	3 to 3.75	12 to 14	2.6 to 3.4
Medium	3.75 to 4.5	14 to 16	3 to 3.8
Fine	3.75 to 4.5	16 to 18	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

soil
textural

**Use the higher rate of Lasso and Lorox Plus in areas of heavy weed infestation. The rate of Lasso may be increased to a maximum of 6 pounds per acre on any soil type to reduce competition from the hard-to-control weeds listed in this label.

511960

■ LASSO plus PREVIEW ■

For weeds controlled preemergence see the "WEEDS CONTROLLED" section of this label for Lasso and Lasso plus Preview.

NOTE: Observe all precautions and limitations on the Lasso and Preview labels including precautions on soil pH, sensitive soybean varieties and rotation crop restrictions. See the following table for recommended rates of Lasso and Preview in this tank mixture.

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE**		TEXTURAL
	Lasso® (pounds)	+ Preview (ounces)	
Coarse — sandy loam only	3 to 3.75	6 to 7	2.6 to 3.4
Medium	3.75 to 4.5	7 to 8	3 to 3.8
Fine	3.75 to 4.5	8 to 10	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

soil textural

**Use the higher rate of Lasso and Preview in areas of heavy weed infestation. The rate of Lasso may be increased to a maximum of 5 pounds per acre on any soil type to reduce competition from the hard-to-control weeds listed in this label.

■ LASSO plus SCEPTER ■

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of this label for Lasso and Lasso plus Scepter.

Note: Observe all precautions and limitations on the Lasso and Scepter labels before use of this tank mixture including minimum recropping interval and rotational guidelines.

~~RECOMMENDED RATES: Apply 2.75 pounds of Lasso plus 2/3 pint of Scepter per acre. The rate of Lasso may be increased to a maximum of 5 pounds per acre on any soil type to reduce competition from the hard-to-control weeds listed in this label.~~

~~DO NOT apply this tank mixture in sprayable fluid fertilizer.~~

See the following table for recommended rates of Lasso and Scepter in this tank mixture.

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE**	
	Lasso (quarts)	Scepter (pints)
Coarse	2.6 to 3.4	2/3
Medium	3 to 3.8	2/3
Fine	3.8 to 4.5	2/3

* Refer to the "SOIL TEXTURE" section of this label to determine the corresponding textural group for the soil to be treated.

**Use the higher rate of Lasso in areas of heavy weed infestations.

DRY BEANS

Dry Beans (All Types)

FOR USE ONLY WEST OF THE MISSISSIPPI RIVER, EXCEPT IN CALIFORNIA.

LASSO®

Apply this product in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: DO NOT APPLY ON DRY BEANS AFTER PLANTING AS CROP INJURY MAY OCCUR. This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

55 60

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 3.75 to 4.5 pounds of this product per acre and shallowly incorporate prior to planting. Use the higher rate in areas of heavy weed infestations or when ~~hard-to-control weeds~~, eastern black or hairy nightshade or yellow nutsedge are present. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

3.8

LASSO®

Lima Bean (Green) ←

FOR USE IN ALL STATES,
EXCEPT IN CALIFORNIA

(Highlight as separate crop section and move this section immediately before the the Mung Bean crop section)

Apply this product in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

APPROVED APPLICAT. SYSTEMS

Ground — Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the ~~proper~~ application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface — Apply this product after planting and before crop and weed emergence and within 5 days after the last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following table for recommended broadcast treatment rates.

Surface Application Rates

SOIL TEXTURE GROUP*	BROADCAST RATE PER ACRE (POUNDS)		TEXTURAL
	LASSO®		
Coarse	2.25 to 3		2.3 to 3
Medium	2.75 to 4.5	←	3.8 to 4.5
Fine**	3.75 to 4.5		3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding ~~texture~~ group for the soil to be treated.

soil
textural

**Use the higher rate in the recommended range in areas of heavy weed infestation or when ~~hard-to-control weeds~~, eastern black or hairy nightshade are present.

NOTE: Preemergence applications are not recommended for use in Illinois or Wisconsin as crop injury may occur.

00071

**LASSO® plus EPTAM
(TANK MIXTURE)
Dry Beans (All Types)**

FOR USE ONLY IN MICHIGAN AND
WEST OF THE MISSISSIPPI RIVER,
EXCEPT IN CALIFORNIA

Apply this tank mixture in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Read the label for Eptam thoroughly for rotational crop restrictions. This tank mixture may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate into the upper 2 inches of soil immediately.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 3 to 4.5 pounds of Lasso plus 2 to 3 pints of Eptam 7-E per acre. Use the higher recommended rate of Lasso for this tank mixture when heavy weed infestations, hard-to-control weeds, eastern

black or hairy nightshade or yellow nutsedge are present. USE ONLY THE 3-POUND RATE OF LASSO IN MICHIGAN. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

**LASSO® plus TREFLAN
(TANK MIXTURE)
Dry Beans (All Types)**

FOR USE ONLY WEST OF THE
MISSISSIPPI RIVER, EXCEPT IN CALIFORNIA

Apply this tank mixture in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Read the label for Treflan thoroughly for rotational crop restrictions. This tank mixture may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this tank mixture within 7 days prior to planting and shallowly incorporate into the upper 2 inches of soil within 8 hours after application.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

00072

57760

Preplant Incorporated Application Rates

BROADCAST RATE PER ACRE (POUNDS)		
SOIL TEXTURE GROUP*	LASSO®	TEXTURAL
Coarse	3	3
Medium	3.75 to 4.5	3.8 to 4.5
Fine**	3.75 to 4.5	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding texture group for the soil to be treated.

**Use the higher rate in the recommended range in areas of heavy weed infestation or when hard-to-control weeds — eastern black or hairy nightshade or yellow nutsedge are present.

soil textural

LASSO® FOLLOWED BY AMIBEN

Dry Beans (All Types)

FOR USE IN MICHIGAN ONLY

Apply Amiben as a sequential preemergence application to Lasso preplant incorporated.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of these products using the system selected.

APPROVED APPLICATION METHODS

LASSO®

Preplant Incorporated — Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

AMIBEN

Preemergence Surface — Apply Amiben after planting and before crop and weed emergence. Read and carefully follow all use instructions and precautionary statements listed on the Amiben label.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: LASSO — Apply 3 pounds of this product per acre and shallowly incorporate or Surface Blend prior to planting.

NOTE: DO NOT USE ON SAND OR LOAMY SANDS. DO NOT APPLY ON DRY BEANS AFTER PLANTING AS CROP INJURY MAY OCCUR. This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

AMIBEN — Apply 4 quarts of this product per acre.

NOTE: When Lasso followed by Amiben is used on soils with 6 percent or more organic matter, weed control may be reduced.

RECOMMENDED RATES: Apply 3.75 to 4.5 pounds of this product plus 1/2 quart of Treflan per acre. Use the higher rate of this product for this tank mixture in areas of heavy weed infestation or when ~~hard-to-control weeds~~, eastern black or hairy nightshade or yellow nutsedge are present. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

3.8

LASSO®

Reg. Kidney Bean Types Only

FOR USE IN ILLINOIS AND WISCONSIN ONLY

Apply this product in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

NOTE: Only one application of Lasso may be made per growing season.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated — Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface — Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

3.8

RECOMMENDED RATES: Apply 3.75 to 4.5 pounds of this product per acre. Use the higher recommended rate when heavy weed infestations, ~~hard-to-control weeds~~, eastern black or hairy nightshade or yellow nutsedge are present. Preplant incorporate only for control of yellow nutsedge.

← (Move Lima Bean crop section here)

LASSO®

Mung Beans ←

← (Highlight this as separate crop section)

FOR USE IN OKLAHOMA ONLY

Apply this product in water solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of product using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface — Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following table for recommended broadcast rates.

Surface Application Rates

SOIL TEXTURE GROUP	BROADCAST RATE PER ACRE (POUNDS)		TEXTURAL
	LASSO		
Coarse*	2.25	← 2.3	
Medium	3		
Fine	3		

*DO NOT USE ON SAND AS CROP INJURY MAY OCCUR.

WOODY ORNAMENTALS

LASSO®

FOR USE ON JUNIPER
(JUNIPERUS SPP.)
AND YEW (TAXUS SPP.)

Apply this product in water solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Topical application or contact of this product with green foliage of the woody ornamental may cause injury and is not recommended. In case green foliage is accidentally sprayed, overhead irrigation within 1 hour after application will minimize injury symptoms (i.e., leaf tip burn, discoloration, distortion, etc.). This product is recommended for this use for application by certified, professional applicators only.

DO NOT USE THIS PRODUCT ON SEEDBEDS, UNROOTED CUTTINGS OR PRIOR TO TRANSPLANTING. DO NOT USE THIS PRODUCT IN GREENHOUSES OR OTHER ENCLOSED STRUCTURES APPLIED THROUGH IRRIGATION SYSTEMS OR MIXED WITH SPRAYABLE FLUID FERTILIZERS. APPLICATIONS DURING PERIODS WHEN TEMPERATURE EXCEEDS 90°F MAY CAUSE INJURY TO ORNAMENTALS.

APPROVED APPLICATION SYSTEMS

Ground — Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the proper application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Post-Transplant Directed — Apply this product as a directed spray after transplanting or to established plantings prior to weed emergence.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 6 pounds of this product per acre as a broadcast treatment. Applications which are not consistent with the recommendations in this label may result in unsatisfactory weed control or injury to ornamental plantings.

NOTE: Repeat applications of this product may be required for sustained weed control. DO NOT exceed 3 applications per year or retreat within 21 days.

← (Insert the following appendix here)

EPA Reg. No. 524-403

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000.

©MONSANTO COMPANY 1988

843.12-000.01/53 — 843.12-000.02/53

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI, 63167 U.S.A.



APPENDIX
LIST OF COMMON AND SCIENTIFIC NAME OF ANNUAL GRASSES
AND BROADLEAF WEEDS

ANNUAL GRASSES

Barnyardgrass	Panicum brevipes
Echinochloa crus-galli	Panicum fasciculatum
Cenchrus	Panicum tall
Digitaria ischaemum	Panicum dichotomiflorum
Digitaria sanguinalis	Panicum, Talls
Cyperus, weedy	Panicum rezarum
Echinochloa villosa	Rice, red
Furcata, giant	Oryza sativa
Setaria faberii	Sandbar, Greenbar
Furcata, green, robust	Cenchrus incertus
perilla, robust white	Shallwgrass, wild cane
Setaria viridis	Sorghum bicolor
Furcata, yellow	Sorghum, broadleaf
Setaria leucocoma	Bracharia platyphylla
Cenchrus	Sorghum, red
Eleusine indica	Lepidochloa filiformis
Johnsongrass, smooth	Whicgrass
Sorghum halepense	Panicum capillare
Millet, green	
Panicum miliaceum	

ANNUAL BROADLEAVES

Beggarweed, Florida	Mustard
Desmodium illinoense	Brassica spp.
Carpetweed	Nightshade, black
Mollugo verticillata	Solanum nigrum
Cocklebur	Nightshade, heavy
Xanthium strumarium	Solanum strachanoides
Colewort	Pigweed, Cornflower
Homo sessilis	Anagallis spp.
Setaria exaltata	Portulaca
Caperweed, baphuracanth	Portulaca oleraceae
Acalypha spp.	Poke, Florida
Gallberry	Pichardia scabra
Gonolobus spp.	Ragweed, common
Greencherry, smooth	Andros artemisiifolia
Physalis spp.	Ragweed, giant
Greencherry, corded	Andros trifida
Physalis angulata	Schlepp
Immortelle	Cassia obtusifolia
Datura stramonium	Sida, prickly, Tootweed
Rudbeckia	Sida spinosa
Kochia scoparia	Smartweed
Lambquarters	Polygonum pennsylvanicum
Chenopodium album	Polygonum persicaria
Morningglory	Starbur, broad
Tall	Acanthaceae
Ipomoea purpurea	Hyssopus
Pitted	Sonchus, common
Ipomoea lacunosa	Helianthus annuus
Hybrid	Polypodium, Blackswamp
Ipomoea hederacea	Asteraceae
Euphorbia	Waterhemp
Ipomoea hederacea	Amaranthus retrofractus
var. integrifolia	
Smallflower	
Jacquemontia caroliniana	

SEDGE

Yellowish, yellow
Cyperus escentus