

Reg # 524-403

PM-25

1777

PARTNER®
WDG Herbicide by Monsanto

Herbicide for weed control in corn, soybeans, and crops listed.

RESTRICTED USE PESTICIDE

due to oncogenicity. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

The use of this product may be hazardous to your health. This product contains alachlor, which has been determined to cause tumors in laboratory animals.

In case of an emergency involving this product or for user safety information on this product, Call Collect, day or night, (314) 694-4000.

Complete Directions For Use

Partner® is a trademark of Monsanto Company.

EPA Reg. No. 524-403

~~1990-2 1992-1~~

~~C843.12-000.32/53 843.12-000.21/53~~

Read each of these sections of this label for essential product performance information.

PRECAUTIONARY STATEMENTS

GENERAL INFORMATION

MIXING, SPRAYING AND
HANDLING INSTRUCTIONS



FLUID FERTILIZER
COMPATIBILITY TEST

APPLICATION INFORMATION

CULTIVATION INFORMATION

WEED CONTROL WITH PARTNER®

CROPS



Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OF MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

(INDEX PAGES)

**Hazards to Humans
and Domestic Animals**

Keep out of reach of children.

CAUTION!

MAY CAUSE EYE IRRITATION.

MAY CAUSE ALLERGIC SKIN REACTION.

Avoid contact with eyes, skin and clothing.

Wear goggles or face shield, rubber gloves, long trousers, long sleeved shirt or jacket of tightly woven material along with boots high enough to cover ankles when transferring and mixing, and when adjusting, repairing or cleaning equipment. Clothing which comes in contact with Partner must be washed before reuse.

Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

IF ON SKIN, immediately flush with plenty of water while removing contaminated clothing. As soon as soap is available, wash skin thoroughly with soap and water. Wash clothing before reuse. Sensitized persons should avoid further contact and reuse of contaminated clothing. Get medical attention.

**In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000.**

Environmental Hazards

This chemical has been identified in limited ground water sampling and there is the possibility that it can leach through the soil to ground water, especially where soils are coarse and ground water is near the surface. Following application, and during rainfall events that cause runoff, this chemical may reach surface water bodies including streams, rivers and reservoirs.

Care must be taken when using this product to prevent back siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Check valves or antisiphoning devices must be used on all mixing equipment. Disposal of excess pesticide, spray mixtures or rinsate should be according to label use instructions or according to the state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. or wetlands. Do not contaminate water when disposing of equipment washwaters.

In case of:

SPILL OR LEAK, remove and keep away from humans and domestic animals.

ACTIVE INGREDIENT:

*Alachlor 65.0%

INERT INGREDIENTS:..... 35.0%
100.0%

*Contains 650 - gms/kilogram or 10.39 ounces/pound of 2-chloro-2'-6'-diethyl-N-(methoxymethyl) acetanilide.

Protected by U.S. Patent No. 4,280,833 and No. 4,417,916. Patent application pending.

~~*Partner is a trademark of Monsanto Company.~~

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

NOT SUBJECT TO FREEZING.

Product can be stored under either heated or unheated conditions.

DISPOSAL:

Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

Emptied container retains product residue. Observe all labeled safeguards until container is destroyed.

Completely empty bag into application equipment. Then dispose of empty bag and box in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Partner herbicide is recommended for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label. This product may be applied either as a surface application before or after planting or shallowly incorporated prior to planting to blend the herbicide

5897

treatment into the upper 1 to 2 inches of soil. ~~Except for minimum or conservation tillage systems,~~ The seedbed should be fine, firm and free of clods and trash when tillage is performed.

Read and carefully observe precautionary statements and all other information appearing on the labeling of all products used in mixtures and sequential treatments.

Do not flood irrigate following application of this product or tank mixtures. This product will not control emerged seedlings.

Do not apply when conditions favor drift.

Use of this product not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

Flush sprayer with clean water after use.

~~Partner® is a trademark of Monsanto Company.~~

SOIL TEXTURE

Applicators should evaluate soil and weed conditions carefully to assure that they choose the lowest effective label rate.

The recommended use rates of this product and the other herbicides labeled for use in tank mixtures with this product generally vary with soil texture. Unless soil texture is specifically named, rate tables throughout this label refer to only three soil texture groups: coarse, medium and fine. The following is a complete listing of soil textures included in each of these three soil texture groups:

<u>SOIL TEXTURAL GROUP</u>	<u>SOIL TEXTURE</u>
COARSE:	sand, loamy sand, sandy loam
MEDIUM:	loam, silt loam, silt, sandy clay loam
FINE:	silty clay loam, clay loam, sandy clay, silty clay, clay

Refer to the above table to determine the corresponding soil textural group for the soil to be treated.

MIXING, SPRAYING AND HANDLING INSTRUCTIONS

Direct contact or exposure to Partner or spray mixtures of Partner should be minimized. The following instructions for location and repairing equipment should be in order to minimize this exposure. Review the protective clothing requirements as listed in the

"PRECAUTIONARY STATEMENTS" section of this label (front panel) and do not use this product until you have the necessary protective clothing.

Equipment Cleaning & Repair

Cleaning and repair of transfer systems and application equipment is a source of exposure to this product. Care should be taken to minimize exposure during cleaning and repair of application equipment. Whenever possible, these systems or equipment should be rinsed before being cleaned or repaired. When repairs must be made during transfer or application, the equipment should be shutdown, and special care taken to avoid contact with the pesticide.

Always predetermine the compatibility of Partner or labeled tank mixtures of Partner with water carrier or sprayable fluid fertilizer carrier by mixing small proportional quantities in advance. See the **"STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST"** section in this label to determine the compatibility of this product and the labeled tank mixtures recommended for use with sprayable fluid fertilizer carrier.

NOTE: Partner and all tank mixtures listed for corn and soybeans in the **"DIRECTION FOR USE"** section may be applied in sprayable fluid fertilizers. **DO NOT** apply Partner or labeled tank mixtures of Partner in sprayable fluid fertilizers on any other crops.

Mix Partner or labeled tank mixture of Partner with the appropriate carrier as follows:

1. Place a ~~20 to 35~~ 50 mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with the appropriate carrier.
3. If a compatibility agent is necessary, add it to the carrier while maintaining agitation to improve mixing or to prevent the formation of undesirable and unsprayable gels or precipitates. Use only compatibility agents cleared by FDA for this use. Read and follow all directions for use, cautionary statements and all other information appearing on the selected compatibility agent label. Check for adequate agitation.
4. Remove the screen prior to adding this product to the spray tank. Mixing of water dispersible granule formulations in sprayable fluid fertilizer carriers may be enhanced by pre-slurrying with water and adding slowly through the screen into the tank. Continue agitation.
5. If a wettable powder or dry flowable formulation is used, make a slurry with water and add it slowly through the screen into the tank. Continue agitation.
6. If a flowable formulation is used, add slowly through the screen into the tank. Mixing and compatibility may be improved

when the flowable is premixed one part flowable with one part water and added to the tank in diluted form.

7. Make a slurry of Partner with water. Add it slowly through the screen into the tank.
8. Complete filling the sprayer tank with carrier. If GranoxoneTM Super Extra is used, add the required amount near the end of the filling process. Remove hose from tank immediately after filling to avoid siphoning back into the carrier source.

Maintain good agitation at all times until the contents of the tank are sprayed. ~~NOTE: If spray mixture is allowed to settle at any time thorough agitation is required to resuspend the mixture before spraying is resumed.~~ If the spray mixture is expected to set unagitated for 8 hours or more (overnight), add a resuspendability agent prior to shutdown (i.e., Hold-UpTM at 8 ounces per 100 gallons of spray solution) or nitrogen liquid fertilizer solution (i.e., 28 percent) at 4 gallons per 100 gallons of spray solution. Allow resuspendability agent to mix with spray mixture before shutting off agitation. Agitate the spray mixture thoroughly to resuspend settled product before resuming spraying.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. Check for even distribution of spray droplets. To reduce loss of the chemical due to drift of a fine mist, apply at nozzle pressures below 40 psi.

TMGranoxone is a trademark of ICI Americas Inc.

TMHold-Up is a trademark of Rigo Company.

STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST

Herbicides may not always mix evenly throughout a sprayable fluid fertilizer or the components may separate too quickly to make their combined use of practical value. This may be due to certain characteristics of the different fluid fertilizers. A simple test using small quantities of the components is suggested to provide compatibility potential.

The test follows:

A. Materials Required For A Compatibility Test

1. Two one-quart jars with lid or stopper (marked "with" and "without").
2. TEAspoons (for a more exacting test, a five to ten milliliter (ml.) pipette or graduated cylinder is desirable).
3. Sprayable fluid fertilizer to be tested.
4. The herbicide chemicals to be mixed.

- 5. A compatibility agent (the purpose of the adjuvant is to help keep the fertilizer and crop protection chemical in suspension, if this assistance is needed.)

B. Procedure

- 1. Add one pint of the sprayable fluid fertilizer that will be used or other herbicide carrier to each jar marked "with" and "without".

**Add One Pint Liquid Fertilizer
To Two Quart Jars.**

WITH

WITHOUT

- 2. To the jar marked "with", add 1/4 TEAspoon or 1.2 milliliters of a suitable compatibility agent; shake gently for five to ten seconds to mix (1/4 TEAspoon in one pint is the equivalent of two pints per 100 gallons of liquid fertilizer).

**To Jar Marked "With"
Add Compatibility Agent
And Shake to Mix.**

WITH

WITHOUT

- 3. To each jar add the appropriate amount of herbicide(s) listed in the following tables. If more than one is used, add them separately with the wetttable powders or dry flowables added first, flowables second and liquid last. Shake gently five to ten seconds after each addition.

**Add Herbicide(s) To Both Jars
And Shake to Mix.**

WITH

WITHOUT

**Amount to Be
Added Per Pint Of
Sprayable Fluid Fertilizer
(Assuming Volume is 25 gal/A)
Level**

HERBICIDE	RATE/A	TEAspoons
WDG's	1 lb.	= 1.5
or	2 lb.	= 3
	3 lb.	= 4.5
Dry Flowables	4 lb.	= 6
	5 lb	= 7.5

HERBICIDE	RATE/A		Level TEAspoons		Milliliters
Emulsifiable	1 pint	=	0.5	or	2.4
Concentrates or	1 quart	=	1	or	4.7
Flowables or	2 quarts	=	2	or	9.5
Liquids or	3 quarts	=	3	or	14.2
Solutions	1 gallon	=	4	or	19.0
	5 quarts	=	5	or	23.8

This compatibility test is designed for 25 gallons of spray per acre. For changes in spray volume or herbicide rates, make appropriate changes in the ingredients of the test. Regardless of spray volume, the amount of compatibility agent should be equal to two or three pints (two pints = 1/4 TEAspoon or 1.2 milliliters, three pints = 3/8 TEAspoon or 1.8 milliliters per pint of sprayable fluid fertilizer) per 100 gallons of liquid fertilizer.

C. Observations and Decisions

1. If the herbicide(s) and the sprayable fluid fertilizer are compatible.
2. If a compatibility agent is necessary.

Five minutes after the final addition and mixing, observe both jars for the formation of large flakes, sludge, gels or other precipitates. Observe if the herbicide(s) cannot be physically mixed with the liquid fertilizer but remains as small oily particles in the solution.

If incompatibility in any form described above occurs in the jar 'with' the compatibility agent added, the liquid fertilizer and the herbicide(s) should not be used together in the same spray tank.

If incompatibility as described above occurs in the jar 'without' the adjuvant but not in the jar 'with' adjuvant, the use of a compatibility adjuvant is recommended.

Both jars should be allowed to stand and be observed periodically for one-half hour. If the separate layers of liquid fertilizer and additives can be resuspended by shaking, commercial application is possible. An emulsifiable concentrate normally will go to the top after standing; WDG's or dry flowables will either settle to the bottom of the tank or jar, or float to the top, depending upon the density of the fertilizers.

If the herbicide(s) is compatible with fluid fertilizer in the foregoing test without having to use a compatibility agent, fluid fertilizer may be used for the premixing. If it is not compatible without the compatibility agent, the herbicide(s) should be premixed with water before adding to the spray tank.

APPLICATION INFORMATION APPLICATION SYSTEMS

Do not apply this product through any type of irrigation system.

Ground Broadcast Treatment - Apply this product and the labeled tank mixtures in 15 or more gallons of solution per acre using broadcast boom equipment. The carrier may be either water or sprayable fluid fertilizer as specified for the crop to be treated in the "DIRECTIONS FOR USE" section of this label. Do not apply during periods of gusty winds, when winds are in excess of 15 mph or when other conditions favoring drift exist.

Band Treatment - Apply a broadcast equivalent rate and volume per acre. To determine these:

Band width <u>in inches</u>	X	Broadcast RATE per acre	=	Band RATE per acre
Row width in inches				

Band width <u>in inches</u>	X	Broadcast VOLUME per acre	=	Band VOLUME per acre
Row width in inches				

Aerial - Fixed wing; helicopter - Apply the recommended rates of this product or approved tank mixtures in 3 to 10 gallons of water per acre. **AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT.** Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray or fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Ensure uniform application. To avoid streaked, uneven or overlapped application use mechanical marking systems only. **HUMAN FLAGGERS PROHIBITED!**

APPLICATION TIMING AND METHODS

~~NOTE: Unless specifically recommended, only one treatment of Partner may be applied to a labeled crop.~~

Early Preplant Surface - This product and some labeled tank mixtures of this product may be applied in no-till and other conservation tillage systems before weeds emerge and up to 45 days before planting certain crops. Split applications can be made 30 to 45 days prior to planting, with 60 percent of the recommended broadcast rate applied initially and the remaining 40 percent applied at planting. Applications made less than 30 days prior to planting can be made either as a split or as a single application. Refer to the individual crop to determine if early preplant application is recommended. If weeds are present at the time of application, apply this product in tank mixture with an appropriate

contact herbicide. Observe direction for use, precautions and restrictions on the label of the contact herbicide. During the planting operation, be careful not to move untreated soil to the surface or move treated soil out of the row, as weed control may be reduced.

[[Preplant Incorporation - This product and many of the labeled tank mixtures may be mixed into the soil using shallow incorporation equipment any time within 7 days prior to planting. Refer to the crop to be treated in the "DIRECTIONS FOR USE" section of this label to determine if preplant incorporation is recommended. Apply the recommended treatment rate to the soil surface as a broadcast application. Precipitation or irrigation is required to bring incorporated herbicide treatments into contact with germinating weed seedlings. The amount of precipitation or irrigation required depends on existing soil moisture, soil type and percent organic matter content, but 1/3 to 3/4 inch is normally adequate. If weeds emerge after treatment, rotary hoe or shallowly cultivate immediately to improve performance.

NOTE: Do not incorporate unless specifically recommended in this label for the crop and herbicide treatment selected. Incorporation of this product and its labeled tank mixtures on coarse soil types may reduce length of weed control due to leaching with rainfall or irrigation.

1-Pass Incorporation (Surface Blend™): Shallowly incorporate (Surface Blend) the treatment into the upper 1 to 2 inches of the soil. Equipment should be operated at manufacturer's designed speed for incorporation to ensure adequate mixing and distribution of the herbicide treatment in the soil.

Equipment design including any drag attachments must be adequate to avoid soil ridging which may result in streaked or reduced weed control. Soil conditions, including moisture content and crop residue levels, must be suitable to allow thorough and uniform mixing with the equipment used for 1-pass incorporation.

Contact Monsanto Company for additional information regarding recommended equipment and methods for Surface Blend application.

2-Pass Incorporation: When 2-pass incorporation is used, shallowly incorporate the herbicide treatment into the upper 1 to 2 inches of the soil with equipment set to work the soil NO DEEPER THAN 4 INCHES. The second pass must be made at an angle to and no deeper than the first pass to ensure proper distribution of the herbicide treatment in the soil.]]

Preemergence Surface Application - This product and all labeled tank mixtures except Partner plus Eptam™, Partner plus Treflan™ and Partner plus Command™ may be applied to the soil surface after planting and prior to either crop or weed emergence. Apply within 5 days of last preplant tillage. If weeds emerge after treatment, or if treatment is applied more than 5 days after last preplant tillage, rotary hoe or shallowly cultivate immediately to improve performance.

12377

Precipitation or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone. The amount of precipitation or overhead sprinkler irrigation required depends on existing soil moisture, soil type and percent organic matter content, but 1/3 to 3/4 inches is normally adequate. Performance is improved when moisture is received within 7 days after application and prior to weed emergence. High intensity or excessive rainfall or excessive irrigation after application may reduce control.

Cultivation Information - Delay cultivation after application for as long as possible unless weeds or grasses emerge. Shallowly cultivate or rotary hoe immediately if weeds or grasses emerge. If cultivation is necessary because of soil crusting or compaction, set equipment shallow and minimize lateral soil movement to avoid dilution or displacement of the herbicide treatment. Only after weeds emerge in the treated band, set cultivator to throw soil into the row covering the band.

TMSurface Blend is a trademark of Monsanto Company.

TMCommand is a trademark of the FMC Corporation.

TMEptam is a trademark of ICI Americas Inc.

TMTreflan is a trademark of DowElanco Products Company.

WEEDS CONTROLLED

Applicators should evaluate soil and weed conditions carefully to assure that they choose the lowest effective label rate.

When applied as directed under conditions described, this product and tank mixtures of this product will control or reduce competition from the weeds listed. Refer to the crop on which treatment is intended to determine the rates and tank mixtures recommended.

For the scientific names to the weeds listed below, refer to the appendix at the end of this booklet.

The tank mixtures of Partner plus atrazine, Partner plus BladexTM, Partner plus ExtrazineTM and Partner plus PrincepTM may not control triazine-resistant weeds as indicated in the tables below for these tank mixtures.

NOTE: C = Control R = Reduced Competition

- 1 Partner®
- 2 Amiben™
- 3 atrazine
- 4 Banvel™
- 5 Bladex™
- 6 Canopy™
- 7 Command
- 8 Gemini™

ANNUAL GRASSES

	Partner® plus							
	1	2	3	4	5	6	7	8
Barnyardgrass	C	E	C	C	C	C	C	C
Crabgrass	C	E	C	C	C	C	C	C
Cupgrass, woolly ¹	R	R	R	•	R	R	R	•
Foxtail: giant, green, robust purple, robust white, yellow	C	E	C	C	C	C	C	C
Goosegrass	C	E	C	C	C	C	C	C
Johnsongrass, seedling ^{2,8}	R	R	R	R	R	R	R	R
Millet, proso ³	Re	•	Re	•	Re	•	•	•
Panicum: browntop, fall	C	E	C	C	C	C	C	C
Panicum, Texas ⁴	R	R	R	R	R	R	R	R
Rice, red ⁵	C	E	C	•	C	C	C	•
Sandbur, Grassbur ^{2,6}	R	•	C	•	•	•	•	•
Shattercane, wild cane ¹	R	R	R	•	R	R	R	•
Signalgrass, broadleaf	C	E	C	C	C	C	C	C
Sprangletop, red	C	E	C	C	C	C	C	C
Witchgrass	C	E	C	C	C	C	C	C

ANNUAL BROADLEAVES

	Partner® plus							
	1	2	3	4	5	6	7	8
Beggarweed, Florida ²	R	E	C	•	C	C	•	C
Carpetweed	C	E	C	C	C	C	C	C
Cocklebur	•	•	R	R	R	C	•	C
Coffeeweed, Hemp sesbania	•	•	•	•	•	C	•	C
Copperleaf, hophornbeam	•	•	•	•	•	C	•	•
Galinsoga	C	E	C	C	C	C	C	C
Groundcherry, annual	•	•	C	•	•	C	•	C
Groundcherry, cutleaf ⁷	C	E	C	C	C	C	C	C
Jimsonweed	•	•	C	•	•	C	C	C
Kochia	•	•	C	•	•	•	•	•
Lambsquarters ²	R	E	C	C	C	C	C	C
Morningglory: tall ⁸	•	•	R	R	R	C	•	C
pitted, ivyleaf ⁸ , entireleaf ⁸	•	•	•	•	•	C	•	C
smallflower	R	R	C	•	C	C	•	C
Mustard	•	R	C	C	C	C	•	C
Nightshade: black ⁷ , hairy ⁷	C	E	C	C	C	C	C	C

Pigweed, Carelessweed	C	E	C	C	C	C	C	C
Purslane	C	E	C	C	C	C	C	C
Pusley, Florida	C	E	C	C	C	C	C	C
Ragweed, common ²	R	E	C	C	C	C	C	C
Ragweed, giant	•	•	•	•	•	R	•	R
Sicklepod	•	•	C	•	C	C	•	R
Sida, prickly; Teaweed ²	R	•	•	•	•	C	C	C
Smartweed ²	R	E	C	C	C	C	R	C
Starbur, bristly	R	R	C	•	C	C	•	•
Sunflower, common	•	•	•	•	•	R	•	•
Velvetleaf, Buttonweed ⁸	•	R	R	R	R	C	C	C
Waterhemp	C	E	C	C	C	C	C	C

SEDGE

Nutsedge, yellow ⁵	C	E	C	•	C	C	C	•
-------------------------------	---	---	---	---	---	---	---	---

NOTE: C = Control R = Reduced Competition

- 1 linuron
- 2 LoroxTM Plus
- 3 PreviewTM
- 4 PursuitTM
- 5 SceptorTM
- 6 SencorTM
- 6 TreflanTM

ANNUAL GRASSES

	Partner TM plus						
	1	2	3	4	4	5	6
Barnyardgrass	C	C	C	C	C	C	C
Crabgrass	C	C	C	C	C	C	C
Cupgrass, woolly ¹	•	•	R	R	R	R	R
Foxtail: giant, green, robust purple, robust white; yellow	C	C	C	C	C	C	C
Goosegrass	C	C	C	C	C	C	C
Johnsongrass, seedling ²	R	R	R	C	C	R	R
Millet, proso ³	•	•	•	•	•	•	•
Panicum: browntop, fall	C	C	C	C	C	C	C
Panicum, Texas ⁴	R	R	R	R	R	R	R
Rice, red ⁵	•	•	C	C	C	C	C
Sandbur, Grassbur ^{2,6}	•	•	•	•	•	•	•
Shattercane, wild cane ¹	•	•	R	R	R	R	R
Signalgrass, broadleaf	C	C	C	C	C	C	C
Sprangletop, red	C	C	C	C	C	C	C
Witchgrass	C	C	C	C	C	C	C

ANNUAL BROADLEAVES

	Partner® plus					
	1	2	3	4	5	6
Beggarweed, Florida ²	•	•	R	•	•	•
Carpetweed	C	C	C	C	C	C
Cocklebur	•	C	C	C	C	•
Coffeeweed	•	•	C	•	•	•
Copperleaf, hophornbeam	•	•	C	•	•	•
Galinsoga	C	C	C	C	C	C
Groundcherry, annual	C	C	C	C	C	C
Groundcherry, cutleaf ⁷	C	C	C	C	C	C
Jimsonweed	•	C	C	C	R	•
Kochia	•	•	•	•	•	•
Lambsquarters ²	C	C	C	C	C	•
Morningglory: tall ⁸	•	R	R	R	•	•
pitted	•	R	R	C	•	•
ivyleaf ⁸ ; entireleaf ⁸	•	R	R	R	•	•
smallflower	•	•	•	C	•	•
Mustard	C	C	C	C	•	•
Nightshade: black ⁷ ; hairy ⁷	C	C	C	C	C	C
Pigweed, Carelessweed	C	C	C	C	C	C
Purslane	C	C	C	C	C	C
Pusley, Florida	C	C	C	C	C	C
Ragweed, common ²	C	C	C	C	C	•
Ragweed, giant	•	R	R	R	R	•
Sicklepod	•	•	•	R	•	•
Sida, prickly; Teaweed ²	C	C	C	C	C	•
Smartweed ²	C	C	C	C	C	•
Starbur, bristly	•	•	•	C	C	•
Sunflower, common	•	•	R	C	•	•
Velvetleaf, Buttonweed ⁸	R	C	C	C	R	•
Waterhemp	C	C	C	C	C	C

SEDGE

Nutsedge, yellow ⁵	•	•	C	C	C	C
-------------------------------	---	---	---	---	---	---

¹Use 6 pounds per acre of this product applied alone or in tank-mix combinations. Preplant incorporate only to reduce competition from this weed.

²For reduced competition from this weed with Partner alone, use 3.8 to 6 pounds of Partner per acre.

³Plant corn into freshly prepared seedbed. Apply 6 pounds of Partner per acre preemergence surface immediately after corn planting for reduced competition from this weed. Improved control may be obtained by applying a labeled postemergence herbicide.

⁴Use a minimum of 4.5 pounds of Partner per acre and apply through center pivot irrigation or under sprinkler irrigation systems. DO NOT apply Partner plus atrazine tank mixtures through center pivot irrigation systems.

⁵Use 4.5 to 6 pounds of Partner per acre applied alone or in tank mixture and apply preplant incorporated only for control on medium and fine-textured soils. When tank-mixed with Pursuit or Scepter, use 3 to 4.5 pounds Partner per acre and preplant incorporate only. On coarse-textured soils with less than 3% organic matter, use 4.5 to 6 pounds of Partner per acre as a preemergence surface application to reduce competition from yellow nutsedge.

⁶Apply 3.8 to 4.5 pounds of Partner plus 1 to 2 quarts of atrazine 4L per acre. Preemergence surface applications should be applied to a freshly prepared seedbed. For best control, rotary hoe or initiate overhead irrigation (0.33 to 0.5 inch) within 24 hours of application.

⁷Use a minimum of 3.8 pounds of Partner per acre on coarse soils, or 4.5 pounds of Partner per acre on medium and fine soils when used alone or in tank mixture with Amiben, Banvel, Canopy, Command, Preview, Sencor/Lexone or Treflan for control of this weed.

⁸For best results on these species when using Partner plus Scepter tank mixture, use preplant incorporated treatments.

NOTE: The 3-way tank mixture of Partner plus atrazine plus Bladex controls or reduces competition as specified for the respective weeds listed under the Partner plus atrazine and Partner plus Bladex recommendations.

~~The 3-way tank mixture of Partner plus Amiben plus Lexone or Sencor controls or reduces competition as specified for the respective weeds listed under the Partner plus Amiben and Partner plus Lexone or Partner plus Sencor recommendations.~~

Certain tank mixtures of Partner and broadleaf herbicides may be combined with applications of Roundup® or Gramoxone Super for control of many emerged weeds prior to soybean emergence. Refer to the Roundup or Gramoxone Super label for a list of emerged weeds controlled by tank mixtures with these products.

~~™Amiben is a trademark of Rhone-Poulenc.~~

™Banvel is a trademark of Sandoz, Inc.

™Bladex, Canopy, Extrazine, Gemini, Lexone, Lorox and Preview are trademarks of E. I. du Pont de Nemours and Company.

™Pursuit and Scepter ~~is a~~ are trademarks of American Cyanamid Corporation.

™Sencor is a trademark of the parent company of Farben Fabricken Bayer GmbH.

™Princep is a trademark of Ciba Geigy Corporation.

®Roundup is a registered trademark of Monsanto Company.

Read each of these sections of this label for essential product performance information.

USE THESE CONVENIENT SECTION MARKERS.

CROPS

CORN

SOYBEANS

BEANS

WOODY ORNAMENTALS

**CORN
(All Types)**

When making applications on corn, do not make more than one two applications per year or exceed a total of 6 pounds of alachlor Partner per acre per year.

PARTNER®

Apply this product in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

Aerial-Fixed wing; helicopter

~~Center-Pivot-Irrigation~~

Dry Bulk Fertilizer Impregnation

Reference: The 'APPLICATION SYSTEMS' section of this label provides detailed information on the application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Irrigation within 10 days following planting may improve weed control.

Preemergence Surface - Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

DO NOT apply this product postemergence, sequential or lay-by.

DO NOT apply any alachlor containing product sequential to this product.

Reference: The 'APPLICATION TIMING AND METHODS' section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

[[Preplant Incorporated Application Rates

12 3 97

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse***	3.8	3.8
Medium	3.8 to 4.5	3.8 to 4.5
Fine	3.8 to 4.5	4.5 to 5.3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate in the recommended range in areas of heavy weed infestations. Use 3.8 to 5.3 pounds per acre where furrow irrigation is used. Use a minimum of 4.5 pounds on coarse textured soils for control of yellow nutsedge or red rice. Rates may be increased to a maximum of 6 pounds on any soil type when heavy infestations of yellow nutsedge or red rice are present.]]

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse***	3 to 3.4	3 to 3.4
Medium	3 to 4.1	3 to 4.1
Fine	3 to 4.1	3.8 to 4.9

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate in the recommended range in areas of heavy weed infestations. Use a minimum of 3.8 to 6 pounds of this product per acre on coarse soils and 4.5 to 6 pounds on medium or fine textured soils to control black or hairy nightshade.

***When applied through center pivot irrigation or under sprinkler irrigation systems on coarse soils, use a minimum of 4.5 pounds per acre of this product.

**PARTNER® plus ATRAZINE
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

[[DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this mixture.]]

NOTE: The total amount of atrazine 4L must not exceed 3 quarts per acre per year.

APPROVE APPLICATION SYSTEMS

Ground-Broadcast boom; banded

Aerial-Fixed wing; helicopter

Dry Bulk Fertilizer Impregnation

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporated or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: ~~PLANT ONLY~~ CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

* There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**				
	Less than 3% organic matter		3% or more organic matter		
Partner® (pounds)	+	atrazine 4L*** (quarts)	Partner (pounds)	+	atrazine 4L*** (quarts)
Coarse	2.3 to 3		0.75 to 2	2.3 to 3	1 to 2
Medium	2.3 to 3		1 to 2	3 to 3.4	1.25 to 2
Fine	3 to 3.4		1.25 to 2	3 to 4.1	1.25 to 2

20 4 71

*Refer to the SOIL TEXTURE section of this label to determine the corresponding soil cextural group for the soil to be treated.

**Use the higher rates in the recommended ranges in areas of heavy weed infestations.

***Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

NOTE: When applied under sprinkler irrigation systems on coarse soils, use 1 quart per acre of atrazine 4L and a minimum of 3.8 pounds per acre of Partner in this tank mixture.

Preplant Incorporated Application Rates

BROADCAST RATE PER ACRE**					
Less than 3% organic matter			3% or more organic matter		
SOIL TEXTURAL GROUP*	Partner® (pounds)	+ atrazine 4L*** (quarts)	Partner (pounds)	+ atrazine 4L*** (quarts)	
Coarse	2.6 to 3.4	1.25 to 2	2.6 to 3.4	1.25 to 2	
Medium	2.6 to 3.4	1.25 to 2	3.4 to 3.8	1.5 to 2	
Fine	3.4 to 3.8	1.5 to 2	3.4 to 4.1	1.5 to 2	

*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rates in the recommended ranges in areas of heavy weed infestations.

***Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

PARTNER® plus BANVEL (TANK MIXTURE)

For use in ILLINOIS, IOWA, MINNESOTA and WISCONSIN on level or flat-planted field corn on fine textured (silty clay loam, clay loam, sandy clay, silty clay or clay) soils with more than 4 percent organic matter.

2. 8 97

Apply in water or sprayable fluid fertilizer solutions for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Use on coarse or medium textured soils or on fine textured soils with 4 percent or less organic matter may result in crop injury and/or destruction.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded.

NOTE: DO NOT apply by air or injection through center pivot irrigation systems.

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using ground spray equipment.

APPROVED APPLICATION METHODS

Preemergence Surface - Apply this tank mixture after planting, before crop and weeds emerge and within 5 days of last preplant tillage operation. Corn seeds must be planted 1-1/2 inches or deeper beneath the soil surface. Direct chemical contact with corn seed must be avoided since crop injury may result. Apply far enough behind planter equipment to avoid any incorporation by the planter wheel or other covering device. If corn seeds are planted less than 1-1/2 inches beneath the soil surface, delay application until corn has spiked.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: PREVENT DRIFT TO SOYBEANS OR OTHER DESIRABLE PLANTS. Do not use on furrow irrigated corn, or when corn is planted at the bottom of a furrow, utilizing lister, till or other similar planting methods. DO NOT incorporate prior to planting or corn emergence if it is necessary to drag for leveling or rotary hoe to break soil crust. DO NOT disturb the soil more than 1/2 inch deep.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury and/or destruction.

BROADCAST RATE PER ACRE

SOIL TEXTURAL GROUP	More than 4% organic matter Partner® (pounds)	Banvel (quarts)
Fine-silty clay loam through clay	3.8 to 4.6	0.5

**PARTNER® plus BLADEX
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge, and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

~~Aerial Fixed wing; helicopter~~

Dry Bulk Fertilizer Impregnation

Reference; The "APPLICATION SYSTEMS" section of this label provides information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

23 7 97

[[Preplant Incorporated Application Rates

BROADCAST RATE PER ACRE

SOIL TEXTURAL GROUP*	1½ to 3½ organic matter		3½ or more organic matter	
	Partner® (pounds)	+ Bladex 4L*** (quarts)	Partner (pounds)	+ Bladex 4L*** (quarts)
Coarse**	2.6 to 3.4	1.2 to 1.4	2.6 to 3.4	1.4 to 1.8
Medium	2.6 to 3.4	1.4 to 1.6	3.4 to 3.8	1.8 to 2.2
Fine	3.4 to 3.8	1.6 to 2	3.4 to 4.1	2 to 2.4

NOTE: See "Incorporation" instructions.

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**On coarse textured soils containing less than 2% organic matter, use Partner plus Bladex only on sandy loam. Do not use on sand and loamy sand with less than 2% organic matter.

~~***Use rates listed in this label when using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equal 1.25 pounds of Bladex 80W.~~

~~***Use equivalent rates listed in this label when using Bladex 80W or Bladex 90DF. One quart of Bladex 4L equal 1.25 pounds of Bladex 80W or 1.1 pounds of Bladex 90DF.~~

Surface Application Rates

BROADCAST RATE PER ACRE

SOIL TEXTURAL GROUP*	1½ to 3½ organic matter		3½ or more organic matter	
	Partner® (pounds)	+ Bladex 4L*** (quarts)	Partner (pounds)	+ Bladex 4L*** (quarts)
Coarse**	2.3 to 3	1 to 1.2	2.3 to 3	1.2 to 1.6
Medium	2.3 to 3	1.2 to 1.4	3 to 3.4	1.6 to 2
Fine	3 to 3.4	1.4 to 1.8	3 to 4.1	1.8 to 2.2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

****On coarse textured soils containing less than 2% organic matter, use Partner plus Bladex only on sandy loam. Do not use on sand and loamy sand with less than 2% organic matter.**

~~*Use rates listed in this label using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 lbs of Bladex 80W.~~**

*****Use equivalent rates listed in this label when using Bladex 80W or Bladex 90DF. One quart of Bladex 4L equal 1.25 pounds of Bladex 80W or 1.1 pounds of Bladex 90DF.**

**PARTNER[®] plus EXTRAZINE II
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge and the annual grasses and broadleaf weeds listed for Micro-Tech plus Bladex in the "WEEDS CONTROLLED" section of this label.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface—Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Preplant Incorporated—Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

***There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.**

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Preplant Incorporated Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE			
	1% to 3% organic matter		3% or more organic matter	
	Partner® (pounds)	Extrazine II + 4L*** (quarts)	Partner (pounds)	Extrazine II + 4L*** (quarts)
Coarse**	2.7 to 3.5	1.5 to 2	2.7 to 3.5	2 to 2.5
Medium	2.7 to 3.5	2 to 2.25	3.5 to 3.8	2.5 to 3
Fine	3.5 to 3.8	2.25 to 2.75	3.5 to 4.2	2.75 to 3.25

NOTE: See "Incorporation" instructions.

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**On coarse textured soils containing less than 2% organic matter, use Partner plus Extrazine II only on sandy loam. Do not use on sand and loamy sand with less than 2% organic matter.

***Use equivalent rates when using Extrazine II DF. One quart of Extrazine II 4L equal 1.1 pounds of Extrazine II DF.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (QUARTS)			
	1% to 3% organic matter		3% or more organic matter	
	Partner® (pounds)	Extrazine II + 4L*** (quarts)	Partner (pounds)	Extrazine II + 4L*** (quarts)
Coarse**	2.3 to 3	1.25 to 1.5	2.3 to 3	1.5 to 2
Medium	2.3 to 3	1.5 to 2	3 to 3.5	2 to 2.75
Fine	3 to 3.5	2 to 2.5	3 to 4.2	2.5 to 3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

****On coarse textured soils containing less than 2% organic matter, use Partner plus Extrazine VI only on sandy loam. Do not use on sand and loamy sand with less than 2% organic matter.**

*****Use equivalent rates when using Extrazine II DF. One quart of Extrazine II 4L equals 1.1 pounds of Extrazine II DF.**

***Extrazine is a trademark of E.I. Du Pont de Nemours and Company.**

**PARTNER® plus MARKSMAN™
(TANK MIXTURE)**

Field and Silage Corn Only

For use in ILLINOIS, IOWA, MINNESOTA and WISCONSIN on level or flat-planted field corn on fine textured (silty clay loam, clay loam, sandy clay, silty clay or clay) soils with more than 4 percent organic matter.

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.

Apply in water or sprayable fluid fertilizer solutions for control of the annual grasses and broadleaf weeds for Partner plus atrazine listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Use on coarse or medium textured soils or on fine textured soils with 4 percent or less organic matter may result in crop injury and/or destruction.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

NOTE: DO NOT apply by air or injection through center pivot irrigation systems.

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using ground spray equipment.

APPROVED APPLICATION METHODS

Preemergence Surface—Apply this tank mixture after planting, before crop and weeds emerge and within 5 days of last preplant tillage operation. Corn seeds must be planted 1-1/2 inches or deeper beneath the soil surface. Direct chemical contact with corn seed must be avoided since crop injury may result. Apply far enough behind planter equipment to avoid any incorporation by the planter wheel or other covering device. If corn seeds are planted less than 1-1/2 inches beneath the soil surface, delay application until corn has spiked.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: CORN, PEANUTS, SORGHUM OR SOYBEANS- can be planted the year following use of this mixture.

* There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

NOTE: PREVENT DRIFT TO SOYBEANS OR OTHER DESIRABLE PLANTS. Do not use on furrow irrigated corn, or when corn is planted at the bottom of a furrow, utilizing lister, till or other similar planting methods. DO NOT incorporate prior to planting or corn emergence. If it is necessary to drag for leveling or rotary hoe to break soil crust, DO NOT disturb the soil more than 1/2 inch deep.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury and/or destruction.

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE	
	Partner® (pounds)	+ Marksman (pints)
Fine-silty clay loam through clay	3.8 to 4.6	3.5

**PARTNER® plus PRINCEP 4L
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

Aerial-Fixed wing; helicopter

Reference: 'THE APPLICATION SYSTEMS' section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

2-791

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

[[Preplant Incorporated Application Rates

BROADCAST RATE PER ACRE**				
Less than 3% organic matter			3% or more organic matter	
SOIL TEXTURAL GROUP*	Partner® + (pounds)	Princep 4L*** (quarts)	Partner + (pounds)	Princep 4L*** (quarts)
Coarse-sandy loam only	2.6 to 3.4	1.25 to 2	2.6 to 3.4	1.25 to 2
Medium	2.6 to 3.4	1.25 to 2	3.4 to 3.8	1.5 to 2
Fine	3.4 to 3.8	1.5 to 2	3.4 to 4.1	1.5 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rates in the recommended ranges in areas of heavy weed infestations.

***Use rates listed in this label when using Princep 4L. Use equivalent rates when using Princep 80W or 90% dry flowable formulations. One quart of 4L formulation equals 1.25 pounds of 80W and 1.1 pounds of the 90% dry flowable formulations.]]

Surface Application Rates

BROADCAST RATE PER ACRE**				
Less than 3% organic matter			3% or more organic matter	
SOIL TEXTURAL GROUP*	Partner® + (pounds)	Princep 4L*** (quarts)	Partner + (pounds)	Princep 4L*** (quarts)

27 9 77

Coarse-sandy loam only	2.3 to 3	0.75 to 2	2.3 to 3	1 to 2
Medium	2.3 to 3	1 to 2	3 to 3.4	1.25 to 2
Fine	3 to 3.4	1.25 to 2	3 to 4.1	1.25 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rates in the recommended ranges in acres of heavy weed infestations.

***Use rates listed in this label when using Princep 4L. Use equivalent rates when using Princep 80W or 90% dry flowable formulations. One quart of the 4L formulation equals 1.25 pounds of the 80W formulation and 1.1 pounds of the 90% dry flowable formulations.

~~TM Princep is a trademark of Ciba-Geigy corporation.~~

**PARTNER® plus ATRAZINE plus
BLADEX (TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

[[DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.]]

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

~~Aerial-Fixed-wing; helicopter~~

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: ~~PLANT-ONLY~~ CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

* There is possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

[[Preplant Incorporated Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	Less than 3% organic matter		
	Partner® (pounds)	+ atrazine 4L** (quarts)	+ Bladex 4L*** (quarts)
Coarse****	2.6 to 3.4	1	0.5 to 1
Medium	2.6 to 3.4	1 to 1.25	1 to 1.25
Fine	3.4 to 3.8	1.25 to 1.5	1 to 1.5

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	3% or more organic matter		
	Partner® (pounds)	+ atrazine 4L** (quarts)	+ Bladex 4L*** (quarts)
Coarse****	2.6 to 3.4	1 to 1.25	1 to 1.25
Medium	3.4 to 3.8	1.25 to 1.5	1 to 2
Fine	3.4 to 4.1	1.5 to 1.75	1 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

~~***Use rates listed in this label using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.~~

***Use equivalent rates listed in this label when using Bladex 80W or Bladex 90DF. One quart of Bladex 4L equal 1.25 pounds of Bladex 80W or 1.1 pounds of Bladex 90DF.

****Do not use this mixture on sand or loamy sand soils with less than 2% organic matter.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.]]

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	Less than 3% organic matter		
	Partner® (pounds)	+ atrazine 4L** (quarts)	+ Bladex 4L*** (quarts)
Course****	2.3 to 3	0.75 to 1.25	0.5 to 1
Medium	2.3 to 3	0.75 to 1	1 to 1.25
Fine	3 to 3.4	1 to 1.25	1 to 1.5

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	3% or more organic matter		
	Partner® (pounds)	+ atrazine 4L** (quarts)	+ Bladex 4L*** (quarts)
Course****	2.3 to 3	0.75 to 1	1 to 1.25
Medium	3 to 3.4	1 to 1.25	1 to 2
Fine	3 to 4.1	1.25 to 1.5	1 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

~~***Use rates listed in this label using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.~~

***Use equivalent rates listed in this label when using Bladex 80W or Bladex 90DF. One quart of Bladex 4L equal 1.25 pounds of Bladex 80W or 1.1 pounds of Bladex 90DF.

****Do not use this mixture on sand or loamy sand soils with less than 2% organic matter.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

**PARTNER® plus ATRAZINE plus PROXL™
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of triazine resistant lambsquarter and pigweed in addition to yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

Ground-Broadcast boom; banded.

Aerial-Fixed wing; helicopter.

Reference: The "APPLICATION SYSTEMS" section of this label provides information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHOD

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

[[DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.]]

DO NOT preplant incorporate this tank mixture as serious crop injury can result.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: ~~PLANT ONLY~~ CORN, PEANUTS, SORGHUM OR SOYBEANS* can be planted the year following use of this mixture.

* There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	Less than 3% organic matter		
	Partner® (pounds)	+ atrazine 4L** (quarts)	+ Prowl (quarts)
Coarse	2.3 to 3	0.75 to 2	0.75
Medium	2.3 to 3	1 to 2	0.75 to 1
Fine	3 to 3.4	1.25 to 2	1

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	3% or more organic matter		
	Partner® (pounds)	+ atrazine 4L** (quarts)	+ Prowl (quarts)
Coarse	2.3 to 3	1 to 2	0.75 to 1
Medium	3 to 3.4	1.25 to 2	1
Fine	3 to 4.1	1.25 to 2	1

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

TMProwl is a trademark of American Cyanamid Corporation.

**PARTNER® plus BLADEX plus PROWL
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of triazine resistant lambsquarter and rigweed in addition to yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

~~Aerial Fixed wing; helicopter~~

Reference: The "APPLICATION SYSTEMS" section of this label provides information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHOD

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

DO NOT preplant incorporate this tank mixture as serious crop injury can result.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

34 7 97

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	1% to 3% organic matter		
	Partner® (pounds)	+ Bladex 4L*** (quarts)	+ Prowl (quarts)
Coarse	2.3 to 3	1 to 1.2	0.75
Medium	2.3 to 3	1.2 to 1.4	0.75 to 1
Fine	3 to 3.4	1.4 to 1.8	1

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	3% or more organic matter		
	Partner® (pounds)	+ Bladex 4L*** (quarts)	+ Prowl (quarts)
Coarse	2.3 to 3	1.2 to 1.6	0.75 to 1
Medium	3 to 3.4	1.6 to 2	1
Fine	3 to 4.1	1.8 to 2.2	1

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Do not use this tank mixture on sand or loamy sand soils with less than 2% organic matter.

***Use rates listed in this label when using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 lbs of Bladex 80W.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

**PARTNER® plus ATRAZINE plus LINURON
(TANK MIXTURE)**

~~FOR USE ONLY IN DELAWARE, MARYLAND, NEW JERSEY, NEW YORK,
PENNSYLVANIA, VIRGINIA AND WEST VIRGINIA.~~
FOR USE ONLY EAST OF THE ROCKY MOUNTAINS.

Apply this tank mixture in water or sprayable fluid fertilizer solutions for control of triazine resistant lambsquarter and pigweed in addition to yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

[[DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.]]

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHOD

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

NOTE: PLANT-ONLY CORN, PEANUTS, SORGHUM, OR SOYBEANS* can be planted the year following use of this mixture.

- * There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	Less than 3% organic matter		
	Partner* (pounds)	+ atrazine 4L*** (quarts)	+ linuron 4L*** (quarts)
Coarse-sandy loam only	2.3 to 3	0.75 to 2	0.33
Medium	2.3 to 3	1 to 2	0.5
Fine	3 to 3.4	1.25 to 2	0.5

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE		
	3% or more organic matter		
	Partner* (pounds)	+ atrazine 4L*** (quarts)	+ linuron 4L*** (quarts)
Coarse-sandy loam only	2.3 to 3	1 to 2	0.5
Medium	3 to 3.4	1.25 to 2	0.5
Fine	3 to 4.1	1.25 to 2	0.5

36 9 97

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

***Use rates listed in this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50% dry flowable formulations. One quart of linuron 4L equals 2 pounds of linuron 50WP or 2 pounds of linuron 50% dry flowable.

NOTE: Use the higher rates in the recommended ranges in areas of heavy weed infestations.

**Conservation or Minimum
Tillage Systems
TANK MIXTURES**

AT-PLANTING APPLICATIONS

When applied as directed under the conditions described, these tank mixtures control emerged annual weeds, suppress many emerged perennial weeds and give preemergence control of many annual grasses and weeds when corn will be planted directly into a cover crop, established sod or in previous crop residues. These tank mixtures will not control regrowth from perennial weeds.

DO NOT APPLY BY AIR.

Refer to specific product labels for crop rotation restriction and cautionary statements of all products used in the tank mixture.

For mixing instructions, see the "MIXING, SPRAYING AND HANDLING INSTRUCTIONS" section of this label.

Partner and the following tank mixtures can be tank mixed with Roundup, Gramoxone Extra and/or 2,4-D.

- **PARTNER® plus ~~GRAMOXONE-SUPER~~** ■
or

- **PARTNER plus ATRAZINE plus ~~GRAMOXONE-EXTRA~~** ■
or

- **PARTNER plus BLADIX plus ~~GRAMOXONE-SUPER~~** ■
or

- **PARTNER plus EXTRAZINE II** ■

37 5 77

■ **PARTNER plus PRINCEP**
plus GRAMOXONE-SUPER
OF

■ **PARTNER plus ATRASINE plus**
BLADEX plus GRAMOXONE-SUPER
OF

■ **PARTNER plus ATRASINE plus**
PRINCEP plus GRAMOXONE-SUPER

~~Apply these tank mixtures with Gramoxone Extra in 20 to 60 gallons of water spray solution per acre~~ Apply these tank mixtures with Roundup or 2,4-D (amine or low-volatile ester) in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre, or the tank mixtures with Gramoxone Extra per in 20 to 60 gallons of water or clear liquid fertilizer per acre immediately before, during or after planting, but BEFORE CROP EMERGENCE. As density of stubble, crop residue or weeds increases, spray gallonage and rate should be increased within the recommended ranges to insure complete coverage. In the absence of emerged vegetation, delete the Roundup, Gramoxone Super Extra or 2,4-D portion of these tank mixtures.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of these products using the system selected.

CONTROL OR SUPPRESSION OF EMERGED WEEDS

ATTENTION

AVOID DRIFT - EXTREME CARE MUST BE USED WHEN APPLYING THESE TANK MIXTURES TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS. Do not allow spray mist to drift since even minute quantities of spray can cause severe damage or destruction to nearby crops, plants or other areas on which treatment is not intended. Do not apply when wind are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow drift to occur. When spraying avoid combinations of pressure and nozzle type that will result in fine particles (mist) which are more likely to drift.

■ **ROUNDUP®** ■

Annual Weeds-Apply 1 to 1.5 pints of Roundup per acre in these tank mixtures when weeds are less than 6 inches tall. Use 2 pints of Roundup per acre for control of fall panicum, barnyardgrass, crabgrass and shattercane up to 2 inches tall; Pennsylvania smartweed up to 6 inches tall; other annual weeds 6 to 12 inches tall. For emerged annual weeds controlled, see the "WEEDS CONTROLLED" section of the label for Roundup.

Perennial Weeds—At normal application dates in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the "GENERAL INFORMATION" section of the label for Roundup for the proper stage of growth for perennial weeds. Use of 2 to 4 quarts of Roundup per acre in the above mixtures under these conditions provides top kill and reduces competition from many emerged perennial grasses and broadleaf weeds.

For emerged perennial weeds controlled see the "WEEDS CONTROLLED" section of the label for Roundup. To obtain control of perennial weeds, follow recommendations on the label for Roundup for stage of growth and rate of application.

USE OF THIS MIXTURE FOR BERMUDAGRASS OR JOHNSONGRASS CONTROL IS NOT RECOMMENDED.

NOTE: When using these tank mixtures, do not exceed 4 quarts of Roundup herbicide per acre.

Ammonium Sulfate—The addition of ammonium sulfate in the spray solution may increase the performance of Roundup tank mixtures on emerged annual weeds under adverse growing conditions. When using ammonium sulfate, add 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water. Ammonium sulfate should be added to the water in the spray tank and completely dissolved prior to adding the herbicide or surfactant. Do not mix ammonium sulfate in fluid fertilizer solutions.

If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet lines. Nozzle tip plugging may result from the use of low-quality ammonium sulfate. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for one minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to adding to the spray tank.

Following use of ammonium sulfate, thoroughly rinse the spray system with clean water to reduce corrosion.

Surfactants—Nonionic surfactants which are labeled for use with herbicides may be used with Roundup. Do not reduce rates of Roundup when adding surfactant. Use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 50 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

■ **GRAMOXONE SUPER EXTRA** ■

When used as directed. Gramoxone Super Extra—~~in tank mixture~~ or labeled tank mixtures controls many emerged annual weeds and suppresses many emerged perennial weeds.

31571

Broadcast Treatment - Apply 1.5 to 2.5 pints of Gramoxone Super in these tank mixtures immediately before, during or after planting but BEFORE CROP EMERGENCE. Use the 2.5 pints rate if any weeds are 4 to 6 inches. As density of stubble, crop residue or weeds increases, spray gallonage should be increased within the recommended range for complete coverage. Add a nonionic spreader surfactant (approved for use on crops) containing at least 75 percent surfactant active agent at 8 ounces per 100 gallons of diluted spray. REFER TO THE GRAMOXONE SUPER LABEL FOR PRECAUTIONARY STATEMENTS.

■ **2,4-D** ■

When used as directed, 2,4-D in labeled tank mixtures controls many emerged annual and perennial broadleaf weeds. For emerged weeds controlled, see the "WEEDS CONTROLLED" section of the label for 2,4-D.

Broadcast Treatment-Apply 1 to 2 pints of 2,4-D (amine or low-volatile ester) in these tank mixtures. Applications should be made 7 to 14 days before planting or 3 to 5 days after planting but BEFORE CORN EMERGES. As density of stubble, crop residue or weeds increases, spray gallonage should be increased within the recommended range for complete coverage.

DO NOT use 2,4-D on light, sandy soils, or where soil moisture is inadequate for normal weed growth. Observe all precautions and limitations on the 2,4-D label booklet.

PREEMERGENCE WEED CONTROL

■ **PARTNER®** ■

For weeds controlled preemergence, see the sections of this label on Partner. See the following table for recommended rates of Partner in this tank mixture on various soil types:

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)
	Partner®
Coarse	3.8
Medium	4.5
Fine	4.5 to 5.3

* Refer to the "SOIL TEXTURE" section of the label to determine the corresponding soil textural group for the soil to be treated.

Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestation, or when organic matter content is 3 percent or more.

■ **PARTNER® plus ATRAZINE** ■

40 11
For weeds controlled preemergence, see the sections of this label for Partner and Partner plus atrazine.

[[DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.]]

See the following table for recommended rates of Partner plus atrazine in this tank mixture on various soil types.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (QUARTS)	
	Partner® (pounds)	+ atrazine 4L** (quarts)
Coarse-sandy loam only	2.6 to 3	1 to 2
Medium	3 to 3.8	1.2 to 2
Fine	3.8 to 4.5	1.6 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present.

Use the higher rate of atrazine in the recommended ranges on soils with greater than 3% organic matter.

NOTE: ~~PLANT ONLY~~ CORN, PEANUTS, SORGHUM OR SOYBEANS can be planted the year following use of this mixture.

There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

■ PARTNER® plus BLADEX ■

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of the label for Partner and this tank mixture.

See the following table for recommended rates of Partner plus Bladex in this tank mixture with Roundup on various soil types.

BROADCAST RATE PER ACRE

SOIL TEXTURAL GROUP*	Partner® (pounds)	+	Bladex 4L** (quarts)
Coarse-sandy loam only	2.6 to 3		1 to 1.6
Medium	3 to 3.8		1.2 to 1.6
Fine	3.8 to 4.5		1.6 to 2.2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**When using Bladex 80W use equivalent rates. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.

Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present.

Use the higher rate of Bladex in the recommended ranges on soils with greater than 3% organic matter.

NOTE: Do not use this mixture on sand or loamy sand soils with less than 2% organic matter.

■ **PARTNER® plus EXTRAZINE II** ■

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of the label for Partner plus Bladex.

See the following table for recommended rates of Partner plus Extrazine II in this tank mixture on various soil types.

BROADCAST RATE PER ACRE (QUARTS)

SOIL TEXTURAL GROUP*	Partner®	+	Extrazine II 4L***
Coarse**	2.7 to 3.5		1.25 to 2
Medium	3 to 3.8		1.5 to 2.75
Fine	3.8 to 4.6		2 to 3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**On coarse textured soils containing less than 2% organic matter, use Partner plus Extrazine II only on sandy loam. Do not use on sand and loamy sand with less than 2% organic matter.

42 71

*****Use equivalent rates when using Extrazine II DF. One quart of Extrazine II 4L equal 1.1 pounds of Extrazine II DF.**

Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present.

Use the higher rate of Extrazine II in the recommended ranges on soils with greater than 3% organic matter.

■ **PARTNER® plus PRINCEP 4L** ■

For weeds controlled preemergence see the 'WEEDS CONTROLLED' section of the labels for Partner and Princep herbicides.

See the following table for recommended rates of Partner plus Princep in this tank mixture on various soil types.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE	
	Partner® (pounds)	+ Princep 4L** (quarts)
Coarse-sandy loam only	2.6 to 3	1 to 1.2
Medium	3 to 3.8	1.2 to 1.6
Fine	3.8 to 4.5	1.6 to 2.2

*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

**When using Princep 80W use equivalent rates. One quart of Princep 4L equals 1.25 pounds of Princep 80W.

Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present.

Use the higher rate of Princep in the recommended ranges on soils with greater than 3% organic matter.

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP, PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDED ROTATIONAL CROPS.

■ **PARTNER® plus ATRASINE plus BLADEX** ■

This tank mixture may be applied for additional preemergence control of heavy weed infestation and/or hard-to-control weeds listed in the 'WEEDS CONTROLLED' section of this label.

43 7 77

[[DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.]]

[[Refer to the following rate table to determine the recommended rates for this tank mixtures on various soil types.]]

BROADCAST RATE PER ACRE

SOIL TEXTURAL GROUP*	Partner® (pounds)	+ atrazine 4L** (quarts)	+ Bladex 4L*** (quarts)
Coarse****	2.6 to 3	0.75 to 1	1 to 1.25
Medium	3 to 3.8	1 to 1.25	1 to 2
Fine	3.4 to 4.5	1 to 1.25	1 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use rates listed in this label when using atrazine 4L. Use equivalent rates when using atrazine 80W or 90% dry flowable formulations. One quart of atrazine 4L equals 1.25 pounds of atrazine 80W or 1.1 pounds of atrazine 90% dry flowable.

***Use rates listed in this label when using Bladex 4L. Use equivalent rates when using Bladex 80W. One quart of Bladex 4L equals 1.25 pounds of Bladex 80W.

****DO NOT use this mixture on sand or loamy sand soils with less than 2% organic matter.

Use the higher rates in the recommended ranges in areas of heavy weed infestations.

NOTE: ~~PLANT-ONLY~~ Corn, peanuts, sorghum or soybeans* the year following use of this mixture.

There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year. DO NOT plant soybeans the year following use of this tank mixture on furrow irrigated corn.

■ PARTNER® plus ATRAZINE plus PRINCEP 4L ■

For additional preemergence control of heavy weed infestation and/or hard-to-control weeds, a three-way tank mixture of Partner herbicide plus atrazine plus Princep may be applied.

[[DO NOT graze treated area or feed treated forage to livestock for 21 days following application of this tank mixture.]]

[[Refer to the following rate table to determine the recommended rates for this tank mixture on various soil types.]]

4-1

BROADCAST RATE PER ACRE

SOIL TEXTURAL GROUP*	Partner® (pounds)	+ atrazine 4L*** (quarts)	+ Princep 4L*** (quarts)
Coarse****	2.6 to 3	0.75 to 1	0.75 to 1
Medium	3 to 3.8	1 to 1.25	1 to 1.5
Fine	3.4 to 4.5	1 to 1.25	1 to 1.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rates of atrazine and Princep in the recommended ranges on soils with greater than 3% organic matter.

***When using 4L formulations use the rates listed. Use equivalent rates when using 80W or 90% dry flowable formulations. One quart of the 4L formulation of these products equals 1.25 pounds of the 80W formulation. One quart of atrazine 4L or Princep 4L equals 1.1 pounds of the 90% dry flowable formulations of these products.

In the areas of heavy grass infestation, use the higher rate of Partner in the recommended ranges when fall panicum, crabgrass, or soils with greater than 3% organic matter will be present.

NOTE: LAND TREATED WITH PRINCEP SHOULD NOT BE PLANTED TO ANY CROP EXCEPT CORN FOR ONE YEAR FOLLOWING TREATMENT AS CROP INJURY MAY OCCUR. AFTER HARVEST OF TREATED CROP, PLOW AND THOROUGHLY TILL THE SOIL IN THE FALL OR SPRING TO MINIMIZE POSSIBLE INJURY TO SPRING SEEDING ROTATIONAL CROPS. PLANT ONLY CORN, PEANUTS, SORGHUM OR SOYBEANS CAN BE PLANTED THE YEAR FOLLOWING USE OF THIS MIXTURE.

There is possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year.

DO NOT plant soybeans the year following use of this tank mixture on furrowed irrigated corn.

EARLY PREPLANT APPLICATION

For use in no-till and other conservation tillage systems in CONNECTICUT, DELAWARE, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA, MISSOURI, NEBRASKA, NEW JERSEY, NEW YORK, NORTH CAROLINA, OHIO, PENNSYLVANIA, SOUTH DAKOTA, VERMONT, VIRGINIA, AND WISCONSIN.

If emerged weeds are present at the time of treatment, Roundup, Gramoxone Extra or 2,4-D should be added to the product according to the directions for use on their respective product labels. If unsatisfactory weed control occurs (due to excessively dry or excessively wet conditions) following the earlier application, a postemergence application of an appropriate labelled grass and/or broadleaf weed herbicide may be used. If a postemergence treatment

includes the herbicide used early preplant, do not exceed the labelled rate for corn on a given soil texture. Observe all precautions and limitations on the labels for Partner, Roundup, and Gramoxone Extra, 2,4-D and other postemergence herbicides before use of these products.

DO NOT apply tank mixture containing Roundup, Gramoxone Extra or other contact herbicides by air.

PARTNER®

This product, when applied in a single application or split application will provide preemergence control or reduced competition of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Aerial - Fixed wing; helicopter

Center Pivot Irrigation

Dry Bulk Fertilizer Impregnation

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application—Application of this product should be made less than 30 days before planting but prior to crop emergence.

NOTE: Application on coarse soils should not be made more than two weeks prior to planting.

Split application—Apply 60 percent of the recommended rate as a split application prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting but before crop emergence.

See the following table for recommended broadcast rates per acre for single and split applications.

BROADCAST RATE PER ACRE (POUNDS)	
SOIL TEXTURAL GROUP*	Partner®
Coarse	3.8
Medium	4.6
Fine	5.4

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

In order to provide broad spectrum weed control, both single and split applications of this product must be followed with a planned postemergence application of a labeled broadcast and/or grass herbicide. Observe the directions for use, precautions and restrictions on the label of the postemergence herbicide before use of these products.

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

**PARTNER® plus ATRASINE
(TANK MIXTURE)**

~~For use in no-till and other conservation tillage systems in DELAWARE, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MARYLAND, MINNESOTA, MISSOURI, NEBRASKA, NORTH CAROLINA, OHIO, SOUTH DAKOTA, VIRGINIA, AND WISCONSIN.~~

This tank mixture, when applied in a single application (alone or in a 3-way combination with Bladex or Princep), split application or as a sequential application to Bladex or Princep in early preplant programs, will provide preemergence control of reduced competition of listed annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

~~If emerged weeds are present at the time of treatment Gramoxone Super or 2,4-D should be added to the product according to the directions for use on their respective product labels. If unsatisfactory weed control occurs (due to excessively dry or excessively wet conditions) following the earlier application, a postemergence application of an appropriate labeled grass and/or broadleaf weed herbicide may be used. If a postemergence treatment includes the herbicide used early preplant, do not exceed the labeled rate for corn on a given soil texture. Observe all precautions and limitations on the labels for Partner, atrazine, Bladex, Princep, Gramoxone Super, 2,4-D and other postemergence herbicides before use of these products.~~

RECOMMENDED RATE AND TIMING OF APPLICATION:

DO NOT graze treated area or feed treated forage to livestock for 21 days following application of these tank mixtures.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Aerial - fixed wing

Dry Bulk Fertilizer Impregnation

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

Single application - Application of this tank mixture should be made less than 30 days before planting but prior to weed emergence.

This application should be restricted to areas north of Interstate 70 when west of the Mississippi River and north of Interstate 64 when east of the Mississippi River where cool spring weather minimizes or slows biodegradation of herbicides.

See the following table for recommended rates.

BROADCAST RATE PER ACRE**		
SOIL TEXTURAL GROUP*	Partner® (pounds)**	+ atrazine 4L (quarts)**
Coarse***	3.4 to 3.8	1.25 to 1.5
Medium	3.8 to 4.5	1.5 to 2
Fine	4.5 to 5-6 5.4	1.75 to 2.25

This tank mixture may also be mixed in a 3-way combination with 1 to 1.25 quarts per acre of Princep or 1.5 to 2 quarts per acre of Bladex to provide improved control of fall panicum and crabgrass.

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the lower rate where light weed infestation is expected and the higher rate where heavy weed infestation is expected.

[[***Application on coarse soils should not be made more than 2 weeks prior to planting.]]

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

Split application - Apply 60 percent of the recommended rate as a split treatment prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting but before crop emergence.

FOR CONTROL OF FALL PANICUM, CRABGRASS OR BROADLEAF SIGNALGRASS USE THE SEQUENTIAL APPLICATION OF THIS TANK MIXTURE FOLLOWING BLADEX OR PRINCEP.

See the following table for recommended rates.

BROADCAST RATE PER ACRE**		
SOIL TEXTURAL GROUP*	Partner** (pounds)*	+ atrazine 4L (quarts)*
Coarse	3.4 to 4.1	1.25 to 1.75
Medium	3.8 to 4.5	1.5 to 2
Fine	4.5 to 5-6 5.4	1.75 to 2.25

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

18791

[[**Use the lower rate where light weed infestation is expected and the higher rate where heavy weed infestation is expected.]]

Sequential application - Application of this tank mixture following Bladex or Princep should be utilized for the control of fall panicum, crabgrass or broadleaf signalgrass. Apply 1.5 to 2 quarts per acre of Bladex or 1 to 1.25 quarts per acre of Princep prior to weed emergence and no more than 45 days prior to planting. At or immediately following planting, but before crop emergence, apply the recommended rate of this tank mixture.

Following applications of Bladex or Princep see the following table for recommended rates.

BROADCAST RATE PER ACRE

SOIL TEXTURAL GROUP	Partner® (pounds)*	+	atrazine 4L (quarts)*
Coarse*	2.6 to 3.8		1 to 1.5
Medium	3.4 to 4.1		1.25 to 1.75
Fine	3.8 to 4.5		1.5 to 2

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

[[**Use the lower rate where light weed infestation is expected and the higher rate where heavy weed infestation is expected.]]

Use rates listed in this label when using the 4L formulation of atrazine, Bladex or Princep. Use equivalent rates of the 80W or 90% dry flowable formulation of atrazine, Bladex or Princep. One quart of the 4L formulation of atrazine, Bladex or Princep equals 1.25 pounds of the 80W and 1.1 pounds of the 90% dry flowable formulations of atrazine, Bladex or Princep.

DO NOT use Bladex on coarse textured soils or soils of less than 1 percent organic matter.

~~Plant only~~ **CORN, PEANUTS, SORGHUM (MILO) OR SOYBEANS** can be planted the year following the use of this program. There is a possibility of crop injury due to carryover of atrazine if soybeans or other nonlabeled crops are planted the following year.

NOTE: Soil treated with Princep should not be planted to any crop except corn for one year following treatment as crop injury may occur. After harvest of the treated crop, plow and thoroughly till the soil in the fall or spring to minimize possible injury to spring seeded rotational crops.

SOYBEANS

When making applications on soybeans, do not make more than one two application per year or exceed a total of ~~4 pounds of alachlor~~ 4.5 pounds of Partner per acre per year.

PARTNER®

Apply this product in water or sprayable fluid fertilizer solution for control of yellow nutsedge, and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

DO NOT feed forage, hay or straw from soybeans treated with this product or tank mixtures of this product. Do not ensile soybeans treated with this product.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

Aerial-Fixed wing; helicopter

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface - Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

37 17

[[Preplant Incorporated Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse	3.8	3.8
Medium	3.8 to 4.5	3.8 to 4.5
Fine	3.8 to 4.5	4.5 to 5.3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate in the recommended range in areas of heavy weed infestations. Use a ~~minimum of 4.5~~ pounds per acre on coarse textured soils for control of yellow nutsedge or red rice. Rates may be increased to a maximum of 6 4.5 pounds on any soil type when heavy infestations of yellow nutsedge or red rice are present.

On coarse soils with 10% or more organic matter use 6 4.5 pounds of Partner per acre.]]

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse***	3 to 3.4	3 to 3.4
Medium	3 to 4.1	3 to 4.1
Fine	3 to 4.1	3.8 to 4.9 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate in the recommended range in areas of heavy weed infestations. Use a minimum of 3.75 pounds of this product per acre on coarse soils and 4.5 ~~to 6~~ pounds on medium or fine textured soils to control eastern black or hairy nightshade or to reduce competition from hard-to-control weeds listed in this label.

***When applied under sprinkler irrigation systems on coarse soils, use a ~~minimum of 4.5~~ pounds per acre of this product.

On coarse soils with 10% or more organic matter use 6 4.5 pounds of Partner per acre.

**~~PARTNER plus AMIBEN~~
(TANK MIXTURE)**

~~Apply this tank mixture in water or sprayable fluid fertilizer solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.~~

51797

~~DO NOT use this tank mixture on sand or loamy sand soils as crop injury may result.~~

~~APPROVED APPLICATION SYSTEMS~~

~~Ground Broadcast boom, banded~~

~~Aerial Fixed wing, helicopter~~

~~Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.~~

~~APPROVED APPLICATION METHODS~~

~~Preplant Incorporated Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.~~

~~Preemergence Surface Apply after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.~~

~~Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for the application timing and method selected.~~

~~RECOMMENDED RATES:~~

~~Preplant Incorporated Apply 3.8 to 4.5 pounds of Partner plus 4 quarts of Amiben per acre.~~

~~NOTE: Use the 4.5 pound per acre rate of Partner in areas of heavy weed infestation. Use 4.5 to 6 pounds of Partner when eastern black or hairy nightshade, yellownutsedge or red rice is present.~~

~~Preemergence Surface Apply 3 pounds of Partner plus 4 quarts of Amiben per acre.~~

~~NOTE: Use a minimum of 3.8 pounds of Partner on coarse soils and 4.5 to 6 pounds in this tank mixture on medium or fine textured soils to control black or hairy nightshade.~~

**PARTNER® plus CANOPY
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Canopy labels including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

DO NOT plant rice or peanuts within 18 months after application of this tank mixture.

52-77

APPROVED APPLICATION SYSTEM

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of the label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight soybean injury early, but soybeans should recover with no subsequent effect on growth or yield.

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Preplant Incorporated Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**				
	1/2 to 3% organic matter		3 to 5% organic matter		
	Partner® (pounds)	+ Canopy (ounces)	Partner + (pounds)	Canopy (ounces)	
Coarse-loamy sand, sandy loam only	3	to 3.8	6	3 to 3.8	8
Medium***	3	to 3.8	8	3.4 to 4.1	10
Fine	3.4	to 4.1	10	3.8 to 4.5	12

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. See "WEEDS CONTROLLED" section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.

***In southeast KS, southwest MO and northeast OK on soils with 1/2 to 3% organic matter, use 5 to 6 ounces Canopy per acre. On silt loam soils in TN and KY, use 6 to 8 ounces of Canopy per acre.

NOTE: In the states of IA, IL, IN, KS, MI, MO, MN, NE, OK, OH, and WI on soils with 1/2 to 4% organic matter, reduced rates of Canopy are recommended for broadleaf weed control. Use 5 ounces of Canopy per acre on coarse soils (loamy sand and sandy loam only); 6 ounces on medium soils; and 6 to 7 ounces of fine soils. Do not apply to soils with greater than 6.8 pH or to soils with less than 1/2% organic matter or greater than 4% organic matter.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command herbicide per acre to this tank mixture and preplant incorporate. Observe all precautions and limitations on the Canopy and Command labels before using this tank mixture.

Surface Application Rates

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	1/2 to 3% organic matter		3 to 5% organic matter	
	Partner® (pounds)	+ Canopy (ounces)	Partner (pounds)	+ Canopy (ounces)
Coarse-loamy sand, sandy loam only	2.6 to 3.4	6	2.6 to 3.4	8
Medium***	2.6 to 3.4	8	3 to 3.8	10
Fine	3 to 3.8	10	3.4 to 4.1	12

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present.

***In southeast KS, southwest MO and northeast OK on soils with 1/2 to 3% organic matter, use 5 to 6 ounces Canopy per acre. On silt loam soils in TN and KY, use 6 to 8 ounces of Canopy per acre.

NOTE: In the states of IA, IL, IN, KS, MI, MO, MN, NE, OK, OH, and WI on soils with 1/2 to 4% organic matter, reduced rates of Canopy are recommended for broadleaf weed control. Use 5 ounces of Canopy per acre on coarse soils (loamy sand and sandy loam only); 6 ounces on medium soils, and 6 to 7 ounces on fine soils. Do not apply to soils with greater than 6.8 pH or to soils with less than 1/2% organic matter or greater than 4% organic matter.

Surface Application Rates

54 7 91

~~BROADCAST RATES PER ACRE**~~

SOIL TEXTURAL GROUP*	1/4 to 3%	3% or more	Canopy (ounces)
	organic matter	organic matter	
Partner® (pounds)	Partner (pounds)		
Coarse-sandy-loam only	2.6 to 3.4	2.6 to 3.4	6
Medium	2.6 to 3.4	3 to 3.8	8
Fine	3 to 3.8	3.4 to 4.1	10

~~*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/4 organic matter.~~

~~**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present.~~

~~Preplant Incorporated Application Rate~~

~~BROADCAST RATES PER ACRE**~~

SOIL TEXTURAL GROUP*	1/4 to 3%	3% or more	Canopy (ounces)
	organic matter	organic matter	
Partner® (pounds)	Partner (pounds)		
Coarse-sandy-loam only	3 to 3.8	3 to 3.8	6
Medium	3 to 3.8	3.4 to 4.1	8
Fine	3.4 to 4.1	3.8 to 4.5	10

~~*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group of the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/4 organic matter.~~

~~**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. See the 'WEEDS CONTROLLED' section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.~~

PARTNER® plus COMMAND (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of johnsongrass and the annual grasses on broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

55 9 97

NOTE: Observe all precautions and limitations on the Partner and Command labels before use of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 3 to 3.8 pounds of Partner plus 1 pint of Command 4E per acre. In areas of heavy pigweed and lambsquarter infestations, use 3.8 to 4.5 pounds of Partner per acre in the tank mixture.

PARTNER® plus GEMINI (TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Gemini labels including precautions on soil pH, minimum recropping interval and rotational guidelines.

DO NOT plant rice or peanuts within 18 months after application of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATES PER ACRE**		
	1/2 to 3% organic matter	3% or more organic matter	
	Partner® (pounds)	Partner® (pounds)	Gemini (ounces)
Coarse-sandy loam only	2.6 to 3.4	2.6 to 3.4	12 to 16
Medium	2.6 to 3.4	3 to 3.8	16 to 20
Fine	3 to 3.8	3.4 to 4.1	20 to 24

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural for the soil to be treated.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. Use the higher rate of Gemini in the recommended ranges on soils with higher organic matter or heavy pressure from large deep germinating weed seeds.

**PARTNER® plus LEXONE
OR
PARTNER plus SENCOR
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: DO NOT use on muck soils or on alkaline soils with a pH greater than 7.4. Crop injury may result if atrazine was applied the year preceding use of this tank mixture.

DO NOT REPLANT CROPS OTHER THAN SOYBEANS FOR 120 DAYS AFTER APPLICATION OF THESE TANK MIXTURES.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Aerial - Fixed wing; helicopter

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight soybean injury early, but soybeans should recover with no subsequent effect on growth or yield.

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

[[Preplant Incorporated Application Rates

BROADCAST RATES PER ACRE					
Less than 3% organic matter			3% or more organic matter		
SOIL TEXTURAL GROUP*	Partner® (pounds)	+ Lexone 4L** or Sencor 4F** (pint)	Partner (pounds)	+ Lexone 4L** or Sencor 4F** (pint)	
Coarse-sandy loam only	3 to 3.8	0.5	3 to 3.8	0.75	
Medium	3 to 3.8	0.5 to 0.75	3.4 to 4.1	0.75	
Fine	3.4 to 4.1	0.75 to 1	3.8 to 4.5	1	

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the rates listed in this label when using Lexone 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lexone 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

***Use the 4.5 pounds per acre rate of Partner in areas of heavy weed infestation. Use 4.5 to 6 pounds of Partner when eastern black or hairy nightshade, yellow nutsedge or red rice are present.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command per acre to this tank mixture and preplant incorporate. Observe all precautions and limitations on the Lexone, Sencor and Command labels before using this tank mixture.

NOTE: Do not use on coarse textured soils. Do not use on medium or fine textured soils with less than 1% organic matter. These mixtures, preplant incorporated, may cause slight soybean injury early, but the soybeans should recover and subsequent growth or yield should not be affected.]]

Surface Application Rates

BROADCAST RATES PER ACRE					
Less than 3% organic matter			3% or more organic matter		
SOIL TEXTURAL GROUP*	Partner® (pounds)	+ Lexone 4L** or Sencor 4F** (pint)	Partner (pounds)	+ Lexone 4L** or Sencor 4F** (pint)	
Coarse - sandy					
loam only	2.6 to 3.4	0.5	2.6 to 3.4	0.75	
Medium	2.6 to 3.4	0.5 to 0.75	3 to 3.8	0.75	
Fine	3 to 3.8	0.75 to 1	3.4 to 4.1	1	

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the rates listed in this label when using Lexone 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lexone 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

NOTE: DO NOT use on sand or loamy sand soil. Use 1.5 pints of Lexone 4L or Sencor 4F per acre in this tank mixture when applying on heavy clay soils of the Mississippi Delta. Use a minimum of 3.8 pounds of Partner on coarse soils and 4.5 to 6 pounds in this tank mixture on medium or fine textured soils to control eastern black or hairy nightshade.

7377

~~PARTNER plus AMIBEN plus LEXONE or
GENCOA (TANK MIXTURES)~~

~~Apply this tank mixture in water solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.~~

~~NOTE: DO NOT use on sand, loamy sand, or on soils with less than 1% organic matter. DO NOT use the foliage from soybeans treated with Partner plus Amiben plus Lexone or Sencor for feed or forage.~~

~~NOTE: DO NOT use on muck soils or on alkaline soils with greater than 7.4 pH. Crop injury may result if atrazine was applied the year preceding use of this tank mixture.~~

~~APPROVED APPLICATION SYSTEMS~~

~~Ground — Broadcast boom~~

~~Aerial — Fixed wing, helicopter~~

~~Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.~~

~~APPROVED APPLICATION METHODS~~

~~Preplant Incorporation — Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight soybean injury early but soybeans should recover with no subsequent effect on growth or yield.~~

~~Preemergence Surface — Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.~~

~~Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing methods selected.~~

~~RECOMMENDED RATES: Refer to the following tables for recommended broadcast treatment rates. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.~~

60 7 97

BROADCAST RATE PER ACRE
Less than 1% organic matter

SOIL	Partner®	Amiben	Lexone 4L**
TEXTURAL	(pounds)	(quarts)	Sencor 4F
GROUPS	(pounds)	(quarts)	(pint)

Coarse			
sandy loam			
only	2.6 to 3.4	3	0.5
Medium	2.6 to 3.4	3 to 4	0.5
Fine	3 to 3.8	4 to 5	0.75

BROADCAST RATE PER ACRE
1% or more organic matter

SOIL	Partner®	Amiben	Lexone 4L**
TEXTURAL	(pounds)	(quarts)	Sencor 4F
GROUPS	(pounds)	(quarts)	(pint)

Coarse			
sandy loam			
only	2.6 to 3.4	3 to 4	0.5 to 0.75
Medium	3 to 3.8	4 to 5	0.5 to 0.75
Fine	3.4 to 4.1	4 to 6	0.75 to 1

~~Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated. On alkaline (calcareous) soils with a pH of 7.4 or greater, apply Lexone 4L or Sencor 4F at 0.5 pint per acre on medium soils and 0.5 to 0.75 pint or corresponding equivalent of WP or DF formulations on fine soils. Use the higher rates of Partner in the recommended ranges in areas of heavy weed infestations and for preplant incorporation applications. Use a minimum of 3.8 pounds of Partner on coarse soils and 4.5 to 6 pounds in this tank mixture on medium or fine soils, when eastern black or hairy nightshade are present. Use 4.5 to 6 pounds of Partner per acre preplant incorporated for control of yellow nutsedge.~~

~~**Use rates listed in this label when using Lexone 4L or Sencor 4F. Use equivalent rates when using the 50WP or DF formulations of these products. One pint of Lexone 4L or Sencor 4F equals 1 pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.~~

PARTNER® plus LINURON
(TANK MIXTURE)

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Plant seed at least 1-3/4 inches deep and DO NOT use on sand or loamy sand or on soils with less than 1% organic matter as crop injury may occur. DO NOT spray over the top of emerged soybeans.

Ground - Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface - Apply after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATES PER ACRE			
	1½ to 3½ organic matter		3½ or more organic matter	
	Partner® + (pounds)	linuron** 4L (quarts)	Partner + (pounds)	linuron** 4L (quarts)
Coarse-sandy loam only	2.6 to 3.4	0.5	2.6 to 3.4	0.5 to 0.75
Medium	2.6 to 3.4	0.5 to 0.75	3 to 3.8	0.75 to 1
Fine	3 to 3.8	1	3.4 to 4.1	1.25 to 1.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the rates listed on this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50% dry flowable formulations. One quart of linuron 4L equals 2 pounds of linuron 50WP or 2 pounds of linuron 50% dry flowable.

**PARTNER® plus LOROX plus
(TANK MIXTURE)**

62797

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Lorox Plus labels including precautions on soil pH, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**		
	<u>1/2 to 3%</u> organic matter	<u>3% or more</u> organic matter	
	Partner® (pounds)	Partner® (pounds)	Lorox Plus (ounces)
Coarse- sandy loam only	2.6 to 3.4	2.6 to 3.4	12 to 14
Medium	2.6 to 3.4	3 to 3.8	14 to 16
Fine	3 to 3.8	3.4 to 4.2	16 to 18

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. Use the higher rate of Lorox Plus in the recommended ranges on soils with higher organic matter or heavy pressure from large deep germinating weed seeds.

60 . 97

**PARTNER® plus PREVIEW
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Preview label including precautions of soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of the label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight soybean injury early, but soybeans should recover with no subsequent effect on growth or yield.

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

64 97

[[Preplant Incorporated Application Rates

BROADCAST RATES PER ACRE**

SOIL TEXTURAL GROUP*	<u>½ to 3½</u> <u>organic matter</u>		<u>3½ or more to 5½</u> <u>organic matter</u>	
	Partner® (pounds)	Preview (ounces)	Partner (pounds)	Preview (ounces)
Coarse-sandy loam only	3 to 3.8	6	3 to 3.8	7
Medium	3 to 3.8	7	3.4 to 4.1	8
Fine	3.4 to 4.1	8	3.8 to 4.5	9

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group of the soil to be tested. DO NOT apply to soil with less than 1/2% organic matter.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. See "WEEDS CONTROLLED" section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.]]

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command herbicide per acre to this tank mixture and preplant incorporate. Observe all precautions and limitations on the Preview and Command labels before using this tank mixture.

Surface Application Rates

BROADCAST RATES PER ACRE**

SOIL TEXTURAL GROUP*	<u>½ to 3½</u> <u>organic matter</u>		<u>3½ or more to 5½</u> <u>organic matter</u>	
	Partner® (pounds)	Preview (ounces)	Partner (pounds)	Preview (ounces)
Coarse-sandy loam only	2.6 to 3.4	6	2.6 to 3.4	7
Medium	2.6 to 3.4	7	3 to 3.8	8
Fine	3 to 3.8	8	3.4 to 4.1	9

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT apply to soil with less than 1/2% organic matter.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present.

**PARTNER® plus PREVIEW plus LOROX PLUS
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of annual grasses and broadleaf weeds listed for Partner plus Preview in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the labels for Partner, Preview and Lorox Plus including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated—Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Preplant incorporation may cause slight soybean injury early, but soybeans should recover with no subsequent affect on growth or yield.

Preemergence Surface—Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following tables for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Preplant Incorporated Application Rates

BROADCAST RATE PER ACRE**

1/2 to 3 1/2% organic matter

SOIL TEXTURAL GROUP*	Partner® (pounds)	+ Preview (ounces)	+ Lorox Plus (ounces)
Coarse-sandy loam only	3 to 3.8	3	6
Medium	3 to 3.8	3.5	7
Fine	3.4 to 4.1	4	8

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present. See the "WEEDS CONTROLLED" section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command herbicide per acre to this tank mixture and preplant incorporate. Observe all precautions and limitations on the Preview and Command labels before using this tank mixture.

Surface Application Rates

BROADCAST RATE PER ACRE**

1/2 to 3 1/2% organic matter

SOIL TEXTURAL GROUP*	Partner® (pounds)	+ Preview (ounces)	+ Lorox Plus (ounces)
Coarse-sandy loam only	2.6 to 3.4	3	6
Medium	2.6 to 3.4	3.5	7
Fine	3 to 3.8	4	8

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when broadleaf signalgrass is present.

**PARTNER® plus PURSUIT
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Pursuit labels before use of this tank mixture, including precautions on minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated—Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface—Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury. See the "WEEDS CONTROLLED" section of this label for recommended rates and application methods for control of certain weeds.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	Less than 3%	3% or more	+ Pursuit (ounces)
	organic matter	organic matter	
	Partner® (pounds)	Partner (pounds)	
Coarse	2.3 to 3	2.3 to 3	4
Medium	2.3 to 3	2.6 to 3.4	4
Fine	2.6 to 3.4	3 to 3.8	4

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when the following weeds are present: fall panicum, barnyardgrass, broadleaf signalgrass, seedling johnsongrass, red rice or yellow nutsedge.

NOTE: Preplant incorporate this tank mixture for more consistent control of common cocklebur, common lambsquarters, jimsonweed, prickly sida, velvetleaf and sunflower and for reduced competition of morningglory and giant ragweed.

**PARTNER® plus SCEPTER
(TANK MIXTURE)**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

Note: Observe all precautions and limitations on the Partner and Scepter labels before use of this tank mixture including precautions on minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Aerial - Fixed wing; helicopter

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface - Apply this tank mixture after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury. See the "WEEDS CONTROLLED" section of this label for recommended rates and application methods for control of certain weeds.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	Less than 3%	3% or more	Scepter*** (pints)
	<u>organic matter</u>	<u>organic matter</u>	
	Partner® (pounds)	Partner (pounds)	
Coarse	2.3 to 3	2.3 to 3	1/2 to 2/3
Medium	2.3 to 3	2.6 to 3.4	1/2 to 2/3
Fine	2.6 to 3.4	3 to 3.8	1/2 to 2/3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when the following weeds are present: fall panicum, barnyardgrass, broadleaf signalgrass, seedling johnsongrass, red rice or yellow nutsedge.

***Use the higher rate of Scepter in the recommended range when red rice, sicklepod, beggarweed or velvetleaf are present.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command 4E herbicide per acre to this tank mixture and preplant incorporate. Observe all precautions and limitations on the Scepter and Command labels before use of this tank mixture.

**PARTNER® plus TREFLAN
(TANK MIXTURE)**

Apply this tank mixture in water solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: See The 'MIXING, SPRAYING AND HANDLING INSTRUCTION' section of this label for special mixing directions for Partner plus Treflan.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Aerial - Fixed wing; helicopter

Reference: The 'APPLICATION SYSTEMS' section of the label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil within 8 hours after application.

Reference: The 'APPLICATION TIMING AND METHODS' section of this label provides detailed information and procedures for the application timing and method selected.

RECOMMENDED RATES: Preplant Incorporated - Apply 3.8 to 4.5 pounds of Partner plus 0.5 quart of Treflan per acre. Applications which are not consistent with the recommendation in this label may result in unsatisfactory weed control or crop injury.

Conservation or Minimum Tillage Systems TANK MIXTURES

SOYBEANS ONLY

When applied as directed under the conditions described, these tank mixtures control many emerged annual weeds, suppress many emerged perennial weeds and give preemergence control of many annual grasses and weeds when soybeans will be planted directly into a cover crop, stale seedbed, or in previous crop residues such as wheat stubble.

These tank mixtures will not control regrowth from perennial weeds.

DO NOT APPLY BY AIR.

For mixing instructions, see the 'MIXING, SPRAYING AND HANDLING INSTRUCTIONS' section of this label.

~~PARTNER plus GRAMOXONE SUPER~~
OF

~~PARTNER plus CANOPY plus~~
~~GRAMOXONE SUPER~~
OF

- ~~PARTNER plus GEMINI plus~~
~~GRAMOXONE SUPER~~
~~OF~~ ■
- ~~PARTNER plus LEXONE plus~~
~~GRAMOXONE SUPER~~
~~OF~~ ■
- ~~PARTNER plus LINURON plus~~
~~GRAMOXONE SUPER~~
~~OF~~ ■
- ~~PARTNER plus LOROX plus~~
~~GRAMOXONE SUPER~~
~~OF~~ ■
- ~~PARTNER plus PREVION plus~~
~~GRAMOXONE SUPER~~
~~OF~~ ■
- ~~PARTNER plus SCIFTER plus~~
~~GRAMOXONE SUPER~~
~~OF~~ ■
- ~~PARTNER plus SENCON plus~~
~~GRAMOXONE SUPER~~ ■

~~Apply these tank mixtures in 20 to 60 gallons of water per acre immediately before, during or after planting, but BEFORE CROP EMERGENCE. As density of stubble, crop residue or weeds increases, spray gallonage and rate should be increased within the recommended ranges to insure complete coverage. In the absence of emerged vegetation delete the Gramoxone Extra portion of these mixes.~~

Partner and the following tank mixtures can be tank-mixed with Roundup and Gramoxone Extra.

- PARTNER ■
- PARTNER plus CANOPY ■
- PARTNER plus GEMINI ■
- PARTNER plus LEXONE ■
- PARTNER plus LINURON ■
- PARTNER plus LOROX PLUS ■
- PARTNER plus PREVION ■
- PARTNER plus PREVION plus LOROX PLUS ■
- PARTNER plus PURSUIT ■
- PARTNER plus SCIFTER ■
- PARTNER plus SENCON ■

Apply the tank mixture with Roundup in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre, or the tank mixture with Gramoxone Extra in 20 to 60 gallons of water or clear liquid fertilizer per acre immediately before, during or after planting, but BEFORE CROP EMERGENCE. As density of stubble, crop residue or weeds increases, spray gallonage and rate should be increased within the recommended ranges to insure complete coverage. In the absence of emerged vegetation, delete the Roundup or Gramoxone Extra portion of these tank mixtures.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of these products using the system selected.

CONTROL OR SUPPRESSION OF EMERGED WEEDS

ROUNDUP®

Annual Weeds-Apply 1 to 1.5 pints of Roundup per acre in these tank mixtures when weeds are less than 6 inches tall. Use 2 pints of Roundup per acre for control of fall panicum, barnyardgrass, crabgrass and shattercane up to 2 inches tall; Pennsylvania smartweed up to 6 inches tall; other annual weeds 6 to 12 inches tall. For emerged annual weeds controlled, see the "WEEDS CONTROLLED" section of the label for Roundup.

Perennial Weeds-At normal application dates in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the "GENERAL INFORMATION" section of the label for Roundup for the proper stage of growth for perennial weeds. Use of 2 to 4 quarts of Roundup per acre in the above mixtures under these conditions provides top kill and reduces competition from many emerged perennial grasses and broadleaf weeds.

For emerged perennial weeds controlled see the "WEEDS CONTROLLED" section of the label for Roundup. To obtain control of perennial weeds, follow recommendations on the label for Roundup for stage of growth and rate of application.

USE OF THIS MIXTURE FOR BERMUDAGRASS OR JOHNSONGRASS CONTROL IS NOT RECOMMENDED.

NOTE: When using these tank mixtures, do not exceed 4 quarts of Roundup herbicide per acre.

Ammonium Sulfate-The addition of ammonium sulfate in the spray solution may increase the performance of Roundup tank mixtures on emerged annual weeds under adverse growing conditions. When using ammonium sulfate, add 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water. Ammonium sulfate should be added to the water in the spray tank and completely dissolved prior

to adding the herbicide or surfactant. Do not mix ammonium sulfate in fluid fertilizer solutions.

If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet lines. Nozzle tip plugging may result from the use of low-quality ammonium sulfate. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for one minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to adding to the spray tank.

Following use of ammonium sulfate, thoroughly rinse the spray system with clean water to reduce corrosion.

Surfactants-Nonionic surfactants which are labeled for use with herbicides may be used with Roundup. Do not reduce rates of Roundup when adding surfactant. Use 0.5 percent surfactant concentration (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 50 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

■ **GRAMOXONE SUPER EXTRA** ■

When used as directed, Gramoxone Super Extra tank-mixed with labeled Partner tank mixtures controls many emerged annual weeds and suppresses many emerged perennial weeds.

Broadcast Treatment - Apply 1.5 to 2.5 pints of Gramoxone Super in this tank mixture immediately before, during or after planting but **BEFORE CROP EMERGENCE**. Use the 2.5 pint rate if any weeds are 4 to 6 inches tall. This mixture may not control weeds taller than 6 inches. As density of stubble, crop residue or weeds increases, spray volume should be increased within the recommended range for complete coverage. Add a nonionic spreader surfactant (approved for use in growing crops) containing at least 75 percent surface active agent at 8 ounces per 100 gallons of diluted spray. **REFER TO THE GRAMOXONE SUPER EXTRA LABEL FOR PRECAUTIONARY STATEMENTS.**

PREEMERGENCE WEED CONTROL

PARTNER®

74 997

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of this label.

See the following table for recommended rates of Partner on various soil types.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (POUNDS)	
	Partner®	
Coarse	3.8	
Medium	4.5	
Fine	4.5 to 5.3	

*Refer to the "SOIL TEXTURE" section of the label to determine the corresponding soil textural group for the soil to be treated.

Use the higher rate in the recommended ranges in areas of heavy grass infestation, or when organic matter content is 3 percent or more.

PARTNER® plus CANOPY

For weeds controlled preemergence see the "WEEDS CONTROLLED" section of this label for Partner and Partner plus Canopy.

NOTE: Observe all precautions and limitations on the Partner and Canopy labels including precautions on soil pH, sensitive soybean varieties and rotational crop restrictions.

See the following table for recommended rates of Partner and Canopy in this tank mixture.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**		
	Partner® (pounds)	+	Canopy (ounces)
Coarse - sandy loam only	2.6 to 3.4		6
Medium	3 to 3.8		8
Fine	3.8 to 4.5		10

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate of Partner and Canopy in areas of heavy weed infestation.

PARTNER® plus GEMINI

For weed controlled preemergence see the "WEEDS CONTROLLED" section of this label for Partner plus Gemini.

Note: Observe all precautions and limitations on the Partner and Gemini labels including precautions on soil pH and rotational crop restrictions.

See the following table for recommended rates of Partner and Gemini in this tank mixture.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**	
	Partner® (pounds)	+ Gemini (ounces)
Coarse - sandy loam only	2.6 to 3.4	12 to 16
Medium	3 to 3.8	16 to 20
Fine	3.8 to 4.5	20 to 24

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate of Partner and Gemini in areas of heavy weed infestation.

- PARTNER® plus LEXONE
- or
- PARTNER® plus SENCOR

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of this label for Partner and Partner plus Lexone 4L or Sencor 4F.

See the following table for recommended rates of Partner plus Lexone 4L or Partner plus Sencor 4F in this tank mixture on various soil types.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE	
	Partner® (pounds)	+ Lexone 4L** or Sencor 4F** (pints)
Coarse - sandy loam only	2.6 to 3.4	0.5 to 0.75
Medium	3 to 3.8	0.75 to 1
Fine	3.8 to 4.5	1 to 1.5***

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the rates listed in this label when using Lexone 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lexone 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

76897
***On the heavy clay soils of the Mississippi Delta, use 1.5 pints of Lexone 4L or Sencor 4F per acre.

Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestations or when fall panicum or crabgrass will be present.

Use the higher rate of Lexone or Sencor in the recommended ranges on soils with greater than 3% organic matter.

DO NOT use this mixture on sand or loamy sand soils as crop injury from Lexone or Sencor may occur.

DO NOT use on muck soils.

DO NOT apply on alkaline soils with a pH greater than 7.4.

Crop injury may occur if atrazine was applied the year before use of this tank mixture.

DO NOT REPLANT CROPS OTHER THAN SOYBEANS FOR 120 DAYS AFTER APPLICATION OF THESE TANK MIXTURES.

■ **PARTNER® plus LINURON** ■

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of this label for Partner and Partner plus linuron.

See the following table for recommended rates of Partner and linuron in this tank mixture.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (QUARTS)	
	Partner® (pounds)	+ linuron 4L** (quarts)
Coarse - sandy loam only	2.6 to 3.4	0.5 to 0.75
Medium	3 to 3.8	0.75 to 1
Fine	3.8 to 4.5	1 to 1.5

*Refer to the "SOIL TEXTURE" section of the label to determine the corresponding soil textural group for the soil to be treated.

**Use the rates listed on this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50 percent dry flowable formulations. One quart of linuron 4L equals 2 pounds of linuron 50WP or 50 percent dry flowable formulation.

Use the higher rate of Partner in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present.

Use the higher rate of linuron in the recommended ranges on soils with greater than 3% organic matter.

11 74 11

DO NOT use this tank mixture on sand or loamy sand or on soil with less than 1% organic matter as crop injury from linuron may occur.

PARTNER® plus LOROX PLUS

For weeds controlled preemergence see the "WEEDS CONTROLLED" section of this label for Partner and Partner plus Lorox Plus.

NOTE: Observe all precautions and limitations on the Partner and Lorox Plus labels including precautions on soil pH and rotational crop restrictions.

See the following table for recommended rates of Partner and Lorox Plus in this tank mixture.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**	
	Partner® (pounds)	+ Lorox Plus (ounces)
Coarse - sandy loam only	2.6 to 3.4	12 to 14
Medium	3 to 3.8	14 to 16
Fine	3.8 to 4.5	16 to 18

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate of Partner and Lorox Plus in areas of heavy weed infestation.

PARTNER® plus PREVIEW

For weeds controlled preemergence see the "WEEDS CONTROLLED" section of this label for Partner and Partner plus Preview.

NOTE: Observe all precautions and limitations on the Partner and Preview labels including precautions on soil pH, sensitive soybean varieties and rotation crop restrictions. See the following table for recommended rates of Partner and Preview in this tank mixture.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**	
	Partner® (pounds)	+ Preview (ounces)
Coarse - sandy loam only	2.6 to 3.4	6
Medium	3 to 3.8	7
Fine	3.8 to 4.5	8

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

****Use the higher rate of Partner and Preview in areas of heavy weed infestation.**

■ PARTNER® plus PREVIEW plus LOROX PLUS ■

For weeds controlled preemergence, see the 'WEEDS CONTROLLED' section of this label for Partner plus Preview.

NOTE: Observe all precautions and limitations on the labels for Partner, Preview and Lorox Plus, including precautions on soil pH, sensitive soybean varieties and rotational crop restrictions.

See the following table for recommended rates of Partner, Preview and Lorox Plus in this tank mixture.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**		
	Partner® (pounds)	+ Preview + (ounces)	Lorox Plus (ounces)
Coarse-sandy loam only	2.6 to 3.4	3	6
Medium	3 to 3.8	3.5	7
Fine	3.8 to 4.5	4	8

*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

****Use the higher rate of Partner in areas of heavy weed infestation.**

■ PARTNER® plus PURSUIT ■

For weeds controlled preemergence, see the 'WEEDS CONTROLLED' section of this label for Partner plus Pursuit.

NOTE: Observe all precautions and limitations on the Partner and Pursuit labels before use of this tank mixture, including minimum recropping interval and rotational guidelines.

See the following table for recommended rates of Partner and Pursuit in this tank mixture.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**	
	Partner® (pounds)	+ Pursuit (ounces)
Coarse	2.6 to 3.4	4
Medium	3 to 3.8	4
Fine	3.8 to 4.5	4

*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding textural group for the soil to be treated.

77 9 11
**Use the higher rate of Partner in areas of heavy weed infestations.

PARTNER® plus SCEPTER

For weeds controlled preemergence, see the "WEEDS CONTROLLED" section of this label for Partner and Partner plus Scepter.

Note: Observe all precautions and limitations on the Partner and Scepter labels before use of this tank mixture including minimum recropping interval and rotational guidelines.

See the following table for recommended rates of Partner and Scepter in this tank mixture.

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**	
	Partner® (pounds)	+ Scepter (pints)
Coarse - sandy loam only	2.6 to 3.4	2/3
Medium	3 to 3.8	2/3
Fine	3.8 to 4.5	2/3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate of Partner in areas of heavy weed infestations.

EARLY PREPLANT APPLICATION

For use in no-till and other conservation tillage systems in CONNECTICUT, DELAWARE, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA, MISSOURI, NEBRASKA, NEW JERSEY, NEW YORK, NORTH CAROLINA, OHIO, PENNSYLVANIA, SOUTH DAKOTA, VERMONT, VIRGINIA, AND WISCONSIN.

If emerged weeds are present at the time of treatment, Roundup, or Gramoxone Extra should be added to the product according to the directions for use on their respective product labels. If unsatisfactory weed control occurs (due to excessively dry or excessively wet conditions) following the earlier application, a postemergence application of an appropriate labelled grass and/or broadleaf weed herbicide may be used. If a postemergence treatment includes the herbicide used early preplant, do not exceed the labelled rate for corn on a given soil texture. Observe all precautions and limitations on the labels for Micro-Tech, Roundup, Gramoxone Extra and other postemergence herbicides before use of these products.

DO NOT apply tank mixture containing Roundup, Gramoxone Extra or other contact herbicides by air.

PARTNER®

This product, when applied in a single application or split application, will provide postemergence control or reduced competition of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Aerial - Fixed wing; helicopter

Center Pivot Irrigation

Dry Bulk Fertilizer Impregnation

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application—Application of this product should be made less than 30 days before planting but prior to weed emergence.

NOTE: Application on coarse soils should not be made more than two weeks prior to planting.

Split application—Apply 60 percent of the recommended rate as a split application prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting but before crop emergence.

See the following table for recommended broadcast rates per acre for single and split applications.

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE (POUNDS) Partner®
Coarse	3.8
Medium	4.5
Fine	4.5

In order to provide broad spectrum weed control, both single and split applications of this product must be followed with a planned postemergence application of a labeled broadcast and/or grass herbicide. Observe the directions for use, precautions and restrictions on the label of the postemergence herbicide before use of these products.

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

**PARTNER® plus CANOPY
(TANK MIXTURE)**

This tank mixture, when applied in a single application or split application in a early preplant program, will provide preemergence control or reduced competition of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Canopy labels, including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

DO NOT plant rice or peanuts within 18 months after application of this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application-Application of this product should be made less than 30 days before planting but prior to weed emergence.

NOTE: Application on coarse soils should not be made more than two weeks prior to planting.

Split application-Apply 60 percent of the recommended rate as a split application prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting but before crop emergence.

See the following table for recommended broadcast rates per acre for single and split applications.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE**	
	Partner® (pounds)	+ Canopy (ounces)
Coarse	3.8	6 to 8
Medium	3.8 to 4.5	8 to 10
Fine	4.5	10 to 12

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

** Use the higher rate of Partner and Canopy in areas of heavy weed infestation.

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

**PARTNER® plus LOROX PLUS
(TANK MIXTURE)**

This tank mixture, when applied in a single application or split application in a early preplant program, will provide preemergence control or reduced competition of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Lorox Plus labels, including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application—Application of this product should be made less than 30 days before planting but prior to weed emergence.

NOTE: Application on coarse soils should not be made more than two weeks prior to planting.

Split application—Apply 60 percent of the recommended rate as a split application prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting but before crop emergence.

See the following table for recommended broadcast rates per acre for single and split applications.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	Partner® (pounds)	Lorox Plus (ounces)
Coarse	3.8	12 to 14
Medium	3.8 to 4.5	14 to 16
Fine	4.5	16 to 18

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

** Use the higher rate of Partner and Lorox Plus in areas of heavy weed infestation.

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

**PARTNER® plus PREVIEW
(TANK MIXTURE)**

This tank mixture, when applied in a single application or split application in a early preplant program, will provide preemergence control or reduced competition of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Preview labels, including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application—Application of this product should be made less than 30 days before planting but prior to weed emergence.

NOTE: Application on coarse soils should not be made more than two weeks prior to planting.

Split application—Apply 60 percent of the recommended rate as a split application prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting but before crop emergence.

See the following table for recommended broadcast rates per acre for single and split applications.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	Partner® (pounds)	+ Preview (ounces)
Coarse	3.8	6 to 7
Medium	3.8 to 4.5	7 to 8
Fine	4.5	8 to 9

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

** Use the higher rate of Partner and Preview in areas of heavy weed infestation.

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

**PARTNER® plus PREVIEW plus LOROX PLUS
(TANK MIXTURE)**

This tank mixture, when applied in a single application in a early preplant program, will provide preemergence control or reduced competition of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner, Preview and Lorox Plus labels, including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application—Application of this product should be made less than 30 days before planting but prior to weed emergence.

NOTE: Application on coarse soils should not be made more than two weeks prior to planting.

See the following table for recommended broadcast rates per acre for single and split applications.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	Partner® (pounds)	+ Preview + (ounces)	Lorox Plus (ounces)
Coarse	3.8	3	6
Medium	3.8 to 4.5	3.5	7
Fine	4.5	4	8

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

** Use the higher rate of Partner in areas of heavy weed infestation.

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

**PARTNER® plus PURSUIT
(TANK MIXTURE)**

This tank mixture, when applied in a single application or split application in a early preplant program, will provide preemergence control or reduced competition of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Pursuit labels, including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application—Application of this product should be made less than 30 days before planting but prior to weed emergence.

NOTE: Application on coarse soils should not be made more than two weeks prior to planting.

Split application—Apply 60 percent of the recommended rate as a split application prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting but before crop emergence.

See the following table for recommended broadcast rates per acre for single and split applications.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	Partner®	+	Pursuit
	(pounds)		(ounces)
Coarse	3.8		4
Medium	3.8 to 4.5		4
Fine	4.5		4

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

** Use the higher rate of Micro-Tech in areas of heavy weed infestation.

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

**PARTNER plus SCEPTER
(TANK MIXTURE)**

This tank mixture, when applied in a single application or split application in a early preplant program, will provide preemergence control or reduced competition of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the Partner and Scepter labels, including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

RECOMMENDED RATE AND TIMING OF APPLICATION

Single application—Application of this product should be made less than 30 days before planting but prior to weed emergence.

NOTE: Application on coarse soils should not be made more than two weeks prior to planting.

Split application—Apply 60 percent of the recommended rate as a split application prior to weed emergence and no more than 45 days prior to planting and the remaining 40 percent at or immediately following planting but before crop emergence.

See the following table for recommended broadcast rates per acre for single and split applications.

BROADCAST RATE PER ACRE**

SOIL TEXTURAL GROUP*	Partner™ (pounds)	+	Scepter (pints)
Coarse	3.8		2/3
Medium	3.8 to 4.5		2/3
Fine	4.5		2/3

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

** Use the higher rate of Partner in areas of heavy weed infestation.

If emerged weeds exist at planting, the application of a contact herbicide or tillage is recommended when possible to eliminate existing weeds.

DRY BEANS

When making applications on dry beans, do not make more than one application per year or exceed a total of ~~3 pounds of alachlor~~ 4.5 pounds of ~~Partner~~ per acre per year.

Dry Beans (All Types)

87997

FOR USE ONLY WEST OF THE MISSISSIPPI RIVER, EXCEPT IN KERN COUNTY, CALIFORNIA.

PARTNER®

Apply this product in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: DO NOT APPLY ON DRY BEANS AFTER PLANTING AS CROP INJURY MAY OCCUR. This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

Do not feed forage or hay from treated crop.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 3.8 to 4.5 pounds of this product per acre and shallowly incorporate prior to planting. Use the higher rate in areas of heavy weed infestations or when eastern black or hairy nightshade or yellow nutsedge are present. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

~~PARTNER® FOLLOWED BY
ANIBEN~~

~~Dry Beans (All Types)~~

~~FOR USE IN MICHIGAN ONLY~~

~~Apply Aniben as a sequential pre-emergence application to Partner~~

~~Ground Broadcast boom~~

~~Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of these products using the system selected.~~

~~APPROVED APPLICATION METHODS~~

~~PARTNER~~

~~Preplant Incorporated — Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.~~

~~Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.~~

~~AMIBEN~~

~~Preemergence Surface — Apply Amiben after planting and before crop and weed emergence. Read and carefully follow all use instructions and precautionary statements listed on the Amiben label.~~

~~Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.~~

~~RECOMMENDED RATES: PARTNER — Apply 3 pounds of this product per acre and shallowly incorporate or Surface Blend prior to planting.~~

~~NOTE: DO NOT USE ON SAND OR LOAMY SANDS. DO NOT APPLY ON DRY BEANS AFTER PLANTING AS CROP INJURY MAY OCCUR. This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.~~

~~AMIBEN — Apply 4 quarts of this product per acre.~~

~~NOTE: When Partner followed by Amiben is used on soils with 6 percent or more organic matter, weed control may be reduced.~~

**PARTNER® plus EPTAM
(TANK MIXTURE)
Dry Beans (All Types)**

**FOR USE ONLY IN MICHIGAN AND
WEST OF THE MISSISSIPPI RIVER,
EXCEPT IN CALIFORNIA**

Apply this tank mixture in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Read the label for Eptam thoroughly for rotational crop restrictions. This tank mixture may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

Do not feed forage or hay from treated crop.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate into the upper 2 inches of soil immediately.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 3 to 4.5 pounds of Partner plus 2 to 3 pints of Eptam 7-E per acre. Use the higher recommended rate of Partner for this tank mixture when heavy weed infestations, eastern black or hairy nightshade or yellow nutsedge are present. **USE ONLY THE 3-POUND RATE OF PARTNER IN MICHIGAN.** Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

**PARTNER® plus TREFLAN
(TANK MIXTURE)
Dry Beans (All Types)**

**FOR USE ONLY WEST OF THE MISSISSIPPI
RIVER, EXCEPT IN KERN COUNTY, CALIFORNIA.**

Apply this tank mixture in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Read the label for Treflan thoroughly for rotational crop restrictions. This tank mixture may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

Do not feed forage or hay from treated crop.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

90 8 97

Reference: The 'APPLICATION SYSTEMS' section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this tank mixture within 7 days prior to planting and shallowly incorporate into the upper 2 inches of soil within 8 hours after application.

Reference: The 'APPLICATION TIMING AND METHODS' section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 3.8 to 4.5 pounds of this product plus 1/2 quart of Treflan per acre. Use the higher rate of this product for this tank mixture in areas of heavy weed infestation or when eastern black or hairy nightshade or yellow nutsedge are present. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

PARTNER®

Red Kidney Bean Types Only

FOR USE IN ILLINOIS AND WISCONSIN ONLY

Apply this product in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

NOTE: This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

Do not feed forage or hay from treated crop.

~~**NOTE: Only one application of Partner may be made per cropping season.**~~

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Reference: The 'APPLICATION SYSTEMS' section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface - Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

91 9 97

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 3.8 to 4.5 pounds of this product per acre. Use the higher recommended rate when heavy weed infestations, eastern black or hairy nightshade or yellow nutsedge are present. Preplant incorporate only for control of yellow nutsedge.

LIMA BEANS (GREEN)

When making applications on lima beans, do not make more than one application per year or exceed a total of ~~3 pounds of alachlor~~ 4.5 pounds of Partner per acre per year.

FOR USE IN ALL STATES,
EXCEPT IN KERN COUNTY, CALIFORNIA

PARTNER®

Apply this product in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: This product may delay crop maturity and/or reduce yield if cold, wet soil conditions occur after planting.

Do not feed forage or hay from treated crop.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated - Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil.

Preemergence Surface - Apply this product after planting and before crop and weed emergence and within 5 days after the last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following table for recommended broadcast treatment rates.

BROADCAST RATE PER ACRE (POUNDS)

SOIL TEXTURAL GROUP*	Partner®
-----------------------------	-----------------

Coarse	3
Medium	3.8 to 4.5
Fine**	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate in the recommended range in areas of heavy weed infestation or when eastern black or hairy nightshade or yellow nutsedge are present.

Surface Application Rates

BROADCAST RATE PER ACRE (POUNDS)

SOIL TEXTURAL GROUP*	Partner®
-----------------------------	-----------------

Coarse	2.3 to 3
Medium	3.8 to 4.5
Fine**	3.8 to 4.5

*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

**Use the higher rate in the recommended range in areas of heavy weed infestation or when eastern black or hairy nightshade are present.

NOTE: Preemergence applications are not recommended for use in Illinois or Wisconsin as crop injury may occur.

MUNG BEANS

When making applications on mung beans, do not make more than one application per year or exceed a total of ~~2 pounds of alachlor~~ 3 pounds of Partner per acre per year.

FOR USE IN OKLAHOMA ONLY

PARTNER®

Apply this product in water solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

APPROVED APPLICATION SYSTEMS

93 9 97

Ground - Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of product using the system selected.

Do not feed forage or hay from treated crop.

APPROVED APPLICATION METHODS

Preemergence Surface - Apply this product after planting, before crop and weed emergence and within 5 days after last preplant tillage operation.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Refer to the following table for recommended broadcast rates.

Surface Application Rates

SOIL TEXTURAL GROUP	BROADCAST RATE PER ACRE (POUNDS)
	Partner®
Coarse*	2.3
Medium	3
Fine	3

*DO NOT USE ON SAND AS CROP INJURY MAY OCCUR.

WOODY ORNAMENTALS

PARTNER®

FOR USE ON JUNIPER
(JUNIPERUS SPP.)
AND YEW (TAXUS SPP.)

Apply this product in water solution for control of the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Topical application or contact of this product with green foliage of the woody ornamental may cause injury and is not recommended. In case green foliage is accidentally sprayed, overhead irrigation within 1 hour after application will minimize injury symptoms (i.e., leaf tip burn, discoloration, distortion, etc.). This product is recommended for this use for application only by certified, professional applicators only.

DO NOT USE THIS PRODUCT ON SEEDBEDS, UNROOTED CUTTINGS OR PRIOR TO TRANSPLANTING. DO NOT USE THIS PRODUCT IN GREENHOUSES, OR OTHER ENCLOSED STRUCTURES, APPLIED THROUGH IRRIGATION SYSTEMS OR MIXED WITH SPRAYABLE FLUID FERTILIZERS. APPLICATIONS DURING PERIODS WHEN TEMPERATURE EXCEEDS 90°F MAY CAUSE INJURY TO ORNAMENTALS.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Reference: The "APPLICATION SYSTEMS" section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Post-Transplant Directed - Apply this product as a directed spray after transplanting or to established plantings prior to weed emergence.

Reference: The "APPLICATION TIMING AND METHODS" section of this label provides detailed information and procedures for application timing and method selected.

RECOMMENDED RATES: Apply 6 pounds of this product per acre as a broadcast treatment. Applications which are not consistent with the recommendations in this label may result in unsatisfactory weed control or injury to ornamental plantings.

NOTE: Repeat applications of this product may be required for sustained weed control. DO NOT exceed ~~3 applications per year~~ 2 applications or 9 pounds of Partner per acre per year or retreat within 21 days.

APPENDIX

LIST OF COMMON AND SCIENTIFIC NAME OF ANNUAL GRASSES AND BROADLEAF WEEDS.

ANNUAL GRASSES

Barnyardgrass

Echinochloa crus-galli

Crabgrass

Digitaria ischaemum

Digitaria sanguinalis

Cupgrass, woolly

Eriochloa villosa

Foxtail, giant

Setaria faberi

**Foxtail, green, robust
purple, robust white
Setaria viridis**

**Foxtail, yellow
Setaria lutescens**

**Goosegrass
Eleusine indica**

**Johnson grass, seedling
Sorghum halepense**

**Millet, proso
Panicum miliaceum**

**Panicum, browntop
Panicum fasciculatum**

**Panicum, fall
Panicum dichotomiflorum**

**Panicum, Texas
Panicum texanum**

**Rice, red
Oryza sativa**

**Sandbur, Grassbur
Cenchrus incertus**

**Shattercane, wild cane
Sorghum bicolor**

**Signalgrass, broadleaf
Brachiaria platyphylla**

**Sprangletop, red
Leptochloa filiformis**

**Witchgrass
Panicum capillare**

ANNUAL BROADLEAVES

**Beggarsweed, Florida
Desmodium tortuosum**

**Carpetweed
Mollugo verticillata**

**Cocklebur
Xanthium strumarium**

**Coffaweed,
Henp sesbania
Sesbania sesban**

Acalypha spp.

Galinsoga

Galinsoga spp.

Groundcherry, annual

Physalis spp.

Groundcherry, cutleaf

Physalis angulata

Jimsonweed

Datura stramonium

Kochia

Kochia scoparia

Lambsquarters

Chenopodium album

Morningglory:

Tall

Ipomoea purpurea

Pitted

Ipomoea lacunosa

Ivyleaf

Ipomoea hederacea

Entireleaf

Ipomoea hederacea

var. *intergriuscula*

Smallflower

Jacquemontia tannifolia

Mustard

Brassica spp.

Nightshade, black

Solanum nigrum

Nightshade, hairy

Solanum sarrachoides

Pigweed, Carelessweed

Amaranthus spp.

Purslane

Portulaca oleracea

Pusley, Florida

Richardia scabra

Ragweed, common

Ambrosia artemisiifolia

Ragweed, giant

Ambrosia trifida

Sicklepod
Cassia obtusifolia

Sida, prickly; Teaweed
Sida spinosa

Smartweed
Polygonum pensylvanicum
Polygonum persicaria

Starbur, bristly
Acanthospermum
hispidum

Sunflower, common
Helianthus annuus

Velvetleaf, Buttonweed
Abutilon theophrasti

Waterhemp
Amaranthus tuberculatos

SEDGE

Nutsedge, yellow
Cyperus esculentus

EPA Reg. No. 524-403

~~C 843.12-000.32/53843.12-000.13/53~~

In case of an emergency involving this product,
Call Collect, day or night, (314) 694-4000.

©MONSANTO COMPANY ~~1990~~ 1992

MONSANTO COMPANY
AGRICULTURAL PRODUCTS
ST. LOUIS, MISSOURI, 63167 U.S.A.