

RESTRICTED USE PESTICIDE

due to oncogenicity. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

SUPPLEMENTAL LABELING DIRECTIONS FOR USE

Read the entire label for Partner® Herbicide before proceeding with the directions for use contained in this supplemental labeling. Follow all applicable restrictions, precautions, and Worker Protection Standard Requirements as well as all applicable application directions.

"Label" as used in this supplemental labeling includes ALL product labeling (label booklet, container labeling, and these directions for use) for Partner Herbicide.

RECD EPA/019/DPD2

'96 AUG 27 09:16

ACCEPTED
JUL -2 1996
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-403



EPA Reg. No. 524-403

Partner® is a registered trademark of Monsanto Company.

The use of this product may be hazardous to your health. This product contains alachlor, which has been determined to cause tumors in laboratory animals.

PARTNER HERBICIDE FOR USE IN PEANUTS

Keep out of reach of children

WARNING! AVISO!

In case of emergency involving this product, Call Collect, day or night, (314)694-4000

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in possession of the user at the time of application.

PARTNER HERBICIDE FOR USE IN PEANUTS

When used according to label directions, this product and certain tank-mixtures may be applied to control or reduce competition from weeds listed in the "WEEDS CONTROLLED" section of the label.

When making applications on peanuts, do not make more than two applications per year or exceed a total of 4.5 pounds of Partner per acre per year.

Do not feed forage or hay from treated crop:

Partner

Apply this product in water solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of the label.

APPROVED APPLICATION SYSTEMS

Ground - Broadcast boom; banded

Aerial - Fixed wing; helicopter

Reference: The "APPLICATION SYSTEMS" section of the label provides detailed information on the application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated* - Apply within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. For control of Texas panicum, this application should be made as a tank mixture with a labeled rate of Balan EC, Prowl, or Sonolan DF Herbicide.

Preemergence Surface - Apply after planting, before crop and weed emergence and within 5 days after the last preplant tillage operation.

Ground-crack Surface - Apply this product at the ground-crack stage of peanuts.

Reference: The "APPLICATION TIMING AND METHODS" section of the label provides detailed information and procedures for application timing and method selected.

*Incorporation on coarse soils may result in reduced length of weed control due to leaching with rainfall or irrigation. For improved control of yellow nutsedge and labeled broadleaf weeds, a labeled sequential ground-crack herbicide application should be made.

Balan and Sonolan are trademarks of DowElanco Product Company.

PARTNER HERBICIDE FOR USE IN PEANUTS - page 2

RECOMMENDED RATES: Refer to the following table for recommended broadcast treatment rates for this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

<u>APPLICATION METHOD</u>	<u>PARTNER® (POUNDS)</u>
Preplant Incorporated	4.5
Preemergence or Ground-crack Surface	4.5

Monsanto Company
800 North Lindbergh Boulevard
St. Louis, Missouri 63167

©Monsanto Company 1996