

15 1/8" LAYFLAT

1/4" NO PRINT ZONE

### LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein and additional conditions and limitations fully stated in the label booklet.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the

soil, crop or treated vegetation. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

© MONSANTO COMPANY 1996

Registered trademark of Monsanto Company.



MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167  
U.S.A.

EPA Est. 524-1A-1



HERBICIDE BY **Monsanto**

Emulsifiable herbicide for weed control in soybeans only.

Read the entire label before using this product. Use only according to label instructions.

Carefully read entire "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION OR REPACKAGING IS PROHIBITED.

Keep out of reach of children.

**WARNING! AVISO!**

Read precautions on back panel.

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

### RESTRICTED USE PESTICIDE

due to oncogenicity. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

The use of this product may be hazardous to your health. This product contains alachlor which has been determined to cause tumors in laboratory animals.

Carefully follow detailed instructions in label booklet.

#### ACTIVE INGREDIENTS:

Alachlor .....	31.7%
Trifluralin .....	3.9%
INERT INGREDIENTS: .....	64.4%
	100.0%

Contains petroleum distillates.

\*Contains 320 grams/litre or 2.67 pounds/gallon of 2-chloro-2',6'-diethyl-N-(methoxy-methyl) acetanilide and 40 grams/litre or 0.33 pound/gallon of (a,a,a-trifluoro-2',6'-dinitro-N,N-dipropyl-p-toluidine).

**NET 2.5 GAL.**

EPA Reg. No. 524-422

12013U2-5M

**ACCEPTED**

MAY 2 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 524-422

PM 2.5  
524-422  
1740

1740

2940

15 1/8" LAYFLAT

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

WARNING! AVISO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY. CAUSES SKIN IRRITATION. HARMFUL IF SWALLOWED. MAY CAUSE RESPIRATORY TRACT IRRITATION. MAY CAUSE ALLERGIC SKIN REACTION.

Do not get in eyes, on skin or on clothing. Avoid breathing vapors or spray mist.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

IF ON SKIN, immediately flush with plenty of water while removing contaminated clothing. Sensitized persons should avoid further contact and reuse of contaminated clothing. Get medical attention.

IF SWALLOWED, do NOT induce vomiting. Give large quantities of water. Get medical attention. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

IF INHALED, remove individual to fresh air. Get medical attention if breathing difficulty develops.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instruction for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: coveralls over short-sleeved shirt and short pants; chemical-resistant gloves, such as barrier laminate or viton; chemical-resistant footwear plus socks; protective eye wear; chemical-resistant apron when cleaning equipment, mixing, or loading; chemical-resistant headgear for overhead exposures.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Users should:
• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

In case of an emergency involving this product, Call Collect, day or night, (314) 594-4000.

Environmental Hazards

This product contains alachlor which has been identified in limited ground water sampling and there is the possibility that it can leach through the soil to ground water, especially where soils are coarse and ground water is near the surface. Following application, and during rainfall events that cause runoff, this chemical (Continued on back panel)

(Continued from side panel)

may reach surface water bodies including streams, rivers and reservoirs.

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Care must be taken when using this product to prevent back-siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Check valves or antisiphoning devices must be used on all mixing equipment. Disposal of excess pesticide, spray mixtures, or rinsate should be according to label use instructions or according to the state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

Physical or Chemical Hazards

In case of SPILL or LEAK, soak up, remove and keep away from humans and domestic animals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: STORE ABOVE 32°F TO KEEP PRODUCT IN SOLUTION.

Below 32°F crystals may form and settle to the bottom. If crystals form, place in a warm room and roll and/or shake the container frequently to redissolve before using.

DISPOSAL: Wastes of this pesticide may cause eye irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed.

Do not reuse container. Triple rinse container. Then puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(See label booklet for Complete Directions for Use.)

For more product information, call toll-free 1-800-332-3111.

3/4" SEAL AREA

4 3/4" CUT OFF 4 27/32" STEP

A. New text aquatic warning.
B. New text spray drift.

2

3740

RESTRICTED USE PESTICIDE

due to oncogenicity. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FREEDOM®  
HERBICIDE by MONSANTO

Emulsifiable herbicide for weed control in soybeans only.

The use of this product may be hazardous to your health. This product contains alachlor which has been determined to cause tumors in laboratory animals.

In case of an emergency involving this product or for user safety information on this product, Call Collect, day or night (314) 694-4000.

EPA Reg. No. 524-422

Complete Directions for Use

®Freedom is a registered trademark of Monsanto Company.

~~1996-1~~ ~~12013U2-2/53~~

Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

REFORMULATION IS PROHIBITED. See individual container label for repackaging limitations.

LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based on contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products

other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Keep out of reach of children.

WARNING! AVISO!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

- CAUSES SUBSTANTIAL BUT TEMPORARY EYE INJURY.
- CAUSES SKIN IRRITATION.
- HARMFUL IF SWALLOWED.
- MAY CAUSE RESPIRATORY TRACT IRRITATION.
- MAY CAUSE ALLERGIC SKIN REACTION.

Do not get in eyes, on skin or on clothing.

Avoid breathing vapors or spray mist.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

IF ON SKIN, immediately flush with plenty of water while removing contaminated clothing. Sensitized persons should avoid further contact and reuse of contaminated clothing. Get medical attention.

IF SWALLOWED, do NOT induce vomiting. Give large quantities of water. Get medical attention. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

IF INHALED, remove individual to fresh air. Get medical attention if breathing difficulty develops.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instruction for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: coveralls over short-sleeved shirt and short pants; chemical-resistant gloves, such as barrier laminate or viton; chemical-resistant footwear plus socks; protective eyewear; chemical-resistant

5940

apron when cleaning equipment, mixing or loading; chemical-resistant headgear for overhead exposure.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4 - 6), the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

Environmental Hazards

This product contains alachlor which has been identified in limited ground water sampling and there is the possibility that it can leach through the soil to ground water, especially where soils are coarse and ground water is near the surface. Following application, and during rainfall events that cause runoff, this chemical may reach surface water bodies including streams, rivers and reservoirs.

This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner which will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Care must be taken when using this product to prevent back-siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Check valves or antisiphoning devices must be used on all mixing equipment. Disposal of excess pesticide, spray mixtures, or rinsate should be according to label use instructions or according to the state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Physical or Chemical Hazards

In case of: SPILL or LEAK, soak up, remove and keep away from humans and domestic animals.

ACTIVE INGREDIENTS:\*

Alachlor.....	31.7%
Trifluralin.....	3.9%
INERT INGREDIENTS:.....	64.4%
	100.0%

Contains petroleum distillates.

\*Contains 320 grams/litre or 2.67 pounds/gallon of 2-chloro-2',6'-diethyl-N-(methoxymethyl)acetanilide and 40 grams/litre or 0.33 pound/gallon of (α,α,α-trifluoro-2',6'-dinitro-N, N-dipropyl-p-toluidine).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short-sleeved shirt and short pants; chemical-resistant gloves, such as barrier laminate or viton; chemical-resistant footwear plus socks; protective eye wear; chemical-resistant headgear for overhead exposure.

For more product information, call toll free 1-800-332-3111.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE: STORE ABOVE 32°F TO KEEP PRODUCT IN SOLUTION.

Below 32°F crystals may form and settle to the bottom. If crystals form, place in a warm room and roll and/or shake the container frequently to redissolve before using. For bulk containers, see the container label for alternate storage information.

DISPOSAL: Wastes of this pesticide may cause eye irritation and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label use instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA regional office for guidance.

Emptied container retains vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed. DO NOT CUT OR WELD ON OR NEAR THIS CONTAINER.

(See the individual container label for disposal information.)

GENERAL INFORMATION

Freedom® is recommended for control of yellow nutsedge and annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label. This product may be applied either as a surface application before or after planting or shallowly incorporated prior to planting to blend the herbicide into the upper 1 to 2 inches of soil.

Performance is often improved when sufficient rainfall or irrigation occurs to moisten the soil within the weed germination zone. If adequate moisture is not received prior to weed emergence, then shallow cultivation is recommended. Refer to the 'APPLICATION AND TIMING METHODS' section of this label for details.

Read and carefully observe the precautionary statements and all other information appearing on the labels of all products used in tank-mixtures and sequential treatments.

Do not flood irrigate following application of this product or tank mixtures.

~~Do not apply when conditions favor drift.~~ Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

Drain and thoroughly flush sprayer with clean water after use and also prior to utilizing spray equipment for treating crops other than soybeans.

Do not graze treated area or feed treated soybean forage, hay or straw to livestock following application.

SOIL TEXTURE

Applicators should evaluate soil and weed conditions carefully to assure that they choose the lowest effective label rate.

The recommended use rates of this product and the other herbicides labeled for use in tank mixtures with this product generally vary with soil texture. Unless soil texture is specifically named, rate tables throughout this label refer to only three soil textural groups: coarse, medium and fine. The following is a complete listing of soil textures included in each of these three soil textural groups:

SOIL TEXTURAL GROUP	SOIL TEXTURE
COARSE	sand, loamy sand, sandy loam
MEDIUM	loam, silt loam, silt, sandy clay loam
FINE	silty clay loam, clay loam, sandy clay, silty clay, clay

Refer to the above table to determine the corresponding soil textural group for the soil to be treated.

**MIXING, SPRAYING AND HANDLING INSTRUCTIONS**

**NOTE**

The use of a mechanical transfer system is required by all mixer/loaders and/or applicators who treat 300 acres or more annually with pesticides containing alachlor as an active ingredient.

Direct contact or exposure to this product or spray mixtures of this product should be minimized. The following instructions for transfer, mixing, cleaning or repairing equipment should be followed in order to minimize this exposure. Review the protective clothing requirements as listed in the "PRECAUTIONARY STATEMENTS" section of this label (front panel) and do not use this product until you are wearing the necessary protective clothing.

**SPRAY DRIFT REQUIREMENTS**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the following information:

**AERIAL DRIFT REDUCTION ADVISORY INFORMATION**



INFORMATION ON DROPLET SIZE The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

WIND Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind, direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the

light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

2.5 Gallon Containers

Open pouring from these containers can result in exposure from splashing or spilling. Special care in lifting and pouring is strongly recommended.

Bulk Containers

This product should be transferred from these containers to the mix or spray tank using pumps or transfer probes. The probe or pump should not be removed from the container or disconnected until the container is emptied and rinsed. Use the pump or probe system to rinse the empty container and transfer the rinsate directly to the mix or spray tank.

Equipment Cleaning and Repair

Cleaning and repair of transfer systems or application equipment is a source of exposure to this product. Care should be taken to minimize exposure during cleaning and repair of transfer systems and application equipment. Whenever possible, these systems or this equipment should be rinsed before being cleaned or repaired. When repairs must be made during transfer or application, the equipment must be shut down and special care should be taken to avoid contact with the pesticide.

Always predetermine the compatibility of this product or labeled tank mixtures of this product with water carrier or sprayable fluid fertilizer carrier by mixing small proportional quantities in advance. See the "STANDARD SPRAYABLE FLUID FERTILIZER COMPATIBILITY TEST" section in this label to determine the compatibility of this product and the labeled tank mixtures recommended for use with sprayable fluid fertilizer carrier.

Mix this product or labeled tank mixture of this product with the appropriate carrier as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with the appropriate carrier.
3. If a compatibility agent is necessary to improve mixing, or to prevent the formation of undesirable and unsprayable gels or precipitates, while agitating, add it to the carrier already in the tank. Use only compatibility agents cleared by FDA for this use. Read and follow all directions for use, precautionary statements and all other information

appearing on the selected compatibility agent label. Check for adequate agitation.

4. If a wettable powder or dry flowable formulation is used, make a slurry with water and add it slowly through the screen into the tank. Continue agitation.
5. If a flowable formulation is used, add slowly through the screen into the tank. Mixing and compatibility may be improved when flowable is premixed one part flowable with one part water and added to the tank in diluted form.
6. Add Freedom herbicide slowly through the screen into the tank. Mixing and compatibility may be improved when this product is prediluted with two parts of water and added to the tank in diluted form.
7. Complete filling the sprayer tank with carrier. If Roundup® herbicide or Gramoxone™ Extra is used, add the required amount near the end of the filling process. Remove hose from tank immediately after filling to avoid siphoning back into the carrier source.

Maintain good agitation at all times until the contents of the tank are sprayed. NOTE: If spray mixture is allowed to settle at any time, thorough agitation is required to resuspend the mixture before spraying is resumed.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select the correct nozzle to avoid spraying a fine mist. Check for even distribution of spray droplets. To reduce loss of chemical due to drift of a fine mist, apply at nozzle pressures below 40 psi.

Certain tank mixtures of Freedom and broadleaf herbicides may be combined with applications of Roundup herbicide or Gramoxone Extra for control of many emerged weeds prior to soybean emergence. Refer to the Roundup or Gramoxone Extra label for a list of emerged weeds controlled by tank mixtures of these products.

®Roundup is a registered trademark of Monsanto Company.

™Gramoxone is a trademark of a ZENECA Group Company.

STANDARD SPRAYABLE FLUID  
FERTILIZER COMPATIBILITY TEST

Herbicides may not always mix evenly throughout a sprayable fluid fertilizer or the components may separate too quickly to make their combined use of practical value. This may be due to certain characteristics of the different fluid fertilizers. A simple test using small quantities of the components is suggested to determine compatibility potential.

The test follows:

A. Materials Required For A Compatibility Test

1. Two one-quart jars with lid or stopper (marked 'with' and 'without').
2. TEAspoon (for a more exacting test, a five to ten milliliter (ml) pipette or graduated cylinder is desirable).
3. Sprayable fluid fertilizer to be tested.
4. The herbicides to be mixed.
5. A compatibility agent (the purpose of the adjuvant is to help keep the fertilizer and crop protection chemical in suspension, if this assistance is needed).

B. Procedure

1. Add one pint of the sprayable fluid fertilizer that will be used or other herbicide carrier to each jar marked 'with' and 'without'.

Add One Pint Liquid Fertilizer  
To Two Quart Jars

WITH

WITHOUT

2. To the jar marked 'with', add 1/4 TEAspoon or 1.2 milliliters of a suitable compatibility agent, shake gently for five to ten seconds to mix (1/4 TEAspoon in one pint is the equivalent of two pints per 100 gallons of liquid fertilizer).

To Jar Marked 'With'  
Add Compatibility Agent  
And Shake To Mix

WITH

WITHOUT

3. To each jar add the appropriate amount of herbicide(s) listed on the following tables. If more than one is used, add them separately with the wettable powders or dry flowables added first, flowables second and liquid last. Shake gently five to ten seconds after each addition.

Add Herbicide(s) To Both Jars  
And Shake To Mix

WITH

WITHOUT

13740

Amount To Be  
Added Per Pint Of  
Sprayable Fluid Fertilizer  
(Assuming Volume is 25 gal/a)

Herbicide	Rate/A	Level	
		TEAspoons	
Wettable Powders or	1 lb =	1.5	
	2 lb =	3	
	3 lb =	4.5	
Dry Flowables	4 lb =	6	
	5 lb =	7.5	

Herbicide	Rate/A	Level		Milliliters
		TEAspoons		
Emulsifiable Concentrates or	1 pint =	0.5	or	2.4
	1 quart	1	or	4.7
Flowables or	2 quarts =	2	or	9.5
	3 quarts =	3	or	14.2
Liquids or Solutions	1 gallon =	4	or	19.0
	5 quarts =	5	or	23.8

The compatibility test is designed for 25 gallons of spray per acre. For changes in spray volume or herbicide rate, make appropriate changes in the ingredients of the test. Regardless of spray volume, the amount of compatibility agent should be equal to two or three pints (two pints = 1/4 TEAspoon or 1.2 milliliters, three pints = 3/8 TEAspoon or 1.8 milliliters per pint of sprayable fluid fertilizer) per 100 gallons of liquid fertilizer.

C. Observations and Decisions

1. If the herbicide(s) and the sprayable fluid fertilizer are compatible.
2. If a compatibility agent is necessary.

Five minutes after the final addition and mixing, observe both jars for the formation of large flakes, sludge, gels or other precipitates. Observe if the herbicide(s) cannot be physically mixed with the liquid fertilizer but remains as small oily particles in the solution.

If incompatibility in any form described above occurs in the jar "with" the compatibility agent added, the liquid fertilizer and the herbicide(s) should not be used together in the same spray tank.

If incompatibility as described above occurs in the jar "without" the adjuvant but not in the jar "with" adjuvant, the use of a compatibility adjuvant is recommended.

Both jars should be allowed to stand and be observed periodically for one-half hour. If the separate layers of liquid fertilizer and additives can be resuspended by shaking, commercial application is possible. An emulsifiable concentrate normally will go to the top after standing, wettable powders will either settle to the bottom of the tank or jar, or float to the top, depending upon the density of the fertilizers.

If the herbicide(s) is compatible with fluid fertilizer in the foregoing test without using a compatibility agent, fluid fertilizer may be used for

premixing. If it is not compatible without the compatibility agent, the herbicide(s) should be premixed with water before adding to the spray tank.

APPLICATION INFORMATION

Do not apply this product through any type of irrigation system.

APPLICATION SYSTEMS

Ground Broadcast Treatment-Apply this product and the labeled tank mixtures in 10 or more gallons of solution per acre using broadcast boom equipment. The carrier may be either water or sprayable fluid fertilizer. Do not apply during periods of gusty winds, when winds are in an excess of 15 mph or when other conditions favoring drift exist.

Aerial-Fixed wing; helicopter-Apply the recommended rates of this product or approved tank mixtures in 3 to 10 gallons of water per acre. ~~AVOID DRIFT DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT.~~ Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray or fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Ensure uniform application. To avoid streaked, uneven or overlapped application, use mechanical marking systems only. **HUMAN FLAGGERS PROHIBITED!** See Spray Drift Requirements and Aerial Drift Advisory Information in the MIXING, SPRAYING AND HANDLING INSTRUCTIONS portion of this label for further information on reducing spray drift during aerial application.

Application with Dry Bulk Fertilizer-The herbicide-dry fertilizer impregnation process (In-Plant and On-Board systems) must be completed only by commercial fertilizer or chemical dealerships properly equipped for this procedure. Contact Monsanto Company for additional information regarding recommended equipment and methods for herbicide-fertilizer impregnation applications.

Dry bulk fertilizer may be impregnated with this product or the tank mixtures of this product plus Lexone™ or Sencor™. Freedom herbicide and these tank mixtures must be applied with 200 to 450 pounds of dry bulk fertilizer per acre and shallowly incorporated within 7 days prior to planting. On medium and fine textured soils in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, applications can be made up to 7 days before planting to allow moisture to move the herbicide-fertilizer mixture into the soil. The correct amount of herbicide must be applied per acre in the blend recommended in this label for the crop selected and weed species to be controlled.

The following table provides a reference to determine the amount of LIQUID herbicide to be mixed per ton of dry bulk fertilizer for a range of herbicide recommendations and fertilizer rates per acre.

15 of 40

RECOMMENDED QUARTS  
LIQUID HERBICIDE/ACRE

Fertilizer Rate (lb/a)	Acres Covered (per ton)	3	3.5	4
		(quarts of herbicide/ton dry bulk fertilizer)		
200	10	30	35	40
250	8	24	28	32
300	6.7	20	23.3	26.6
350	5.7	17.1	20	22.8
400	5	15	17.5	20
450	4.5	13.3	15.6	17.8

To determine the amount of herbicide needed for rates not included in the preceding table, use the following formula:

$$\frac{\text{Recommended herbicide rate (lb or qt/acre} \times 2000)}{\text{pounds fertilizer/acre}} = \frac{\text{pounds or quarts of herbicide}}{\text{per ton of dry bulk fertilizer}}$$

With In-Plant systems, mix and blend the dry fertilizer and herbicide mixture in a closed rotary drum-type mixer, allowing sufficient time to ensure uniform coverage. Use at least one ton of dry fertilizer per mixing operation. Inject the herbicide into the drum over a minimum of a 2-minute period and allow at least 2 additional minutes mixing time to ensure uniformity. The nozzle used to spray the herbicide treatment must be placed inside the mixer to provide uniform spray coverage of the tumbling fertilizer.

If the dry fertilizer used has inadequate absorptive capacity, a highly absorptive material such as Agsorb™, MP-79™, or Microcel™ E can be added to provide a free-flowing mixture. Contact Monsanto Company for specific guidelines with regard to the sequence of addition for the various components and the amount of drying agent to add to provide a free-flowing mixture. The absorptive material and dry fertilizer should be thoroughly mixed prior to adding liquid herbicide. Generally the drying material weight is no more than 2 to 4 percent of the total weight of the dry fertilizer. The following dry fertilizers may be impregnated with this product or tank mixtures of this product with Sencor or Lexone: ammonium sulfate, ammonium phosphate-sulfate, diammonium phosphate, potassium chloride, single super-phosphate, treble super-phosphate and urea.

™Agsorb is a trademark of Oil Dri Corporation.

™Lexone is a trademark of E. I. du Pont de Nemours and Company.

™Microcel is a trademark of Manville Products Corporation.

™MP-79 is a trademark of Eagle-Picher Industries, Inc.

™Sencor is a trademark of the parent company of Farbenfabriken Bayer GmbH.

NOTE: DO NOT impregnate this product or tank mixtures of this product with other herbicides on fertilizers containing ammonium nitrate, potassium nitrate or sodium nitrate.

Spread the dry fertilizer-herbicide mixture uniformly with an adequately calibrated dribble, pneumatic (air flow) or spin applicator. When using spin

applicators, the herbicide-dry fertilizer mixture must be spread uniformly. Nonuniform spreading of the dry fertilizer-herbicide mixture may result in unsatisfactory weed control or crop injury.

APPLICATION TIMING AND METHODS

Preplant Incorporation-This product and the labeled tank mixtures may be mixed into the soil using shallow incorporation equipment any time within 14 days prior to planting. Apply the recommended treatment rate to the soil surface as a broadcast application and shallowly incorporate into the upper 2 inches of soil within 24 hours after application. If weeds emerge after treatment, rotary hoe or shallowly cultivate immediately to improve performance.

1-Pass Incorporation (Surface Blend™): Shallowly incorporate (Surface Blend) the treatment into the upper 1 to 2 inches of the soil. Equipment should be operated at manufacturer's designated speed for incorporation to ensure adequate mixing and distribution of the herbicide treatment in the soil. Equipment design, including any drag attachments, must be adequate to avoid soil ridging which may result in streaked or reduced weed control. Soil conditions, including moisture content and crop residue levels, must be suitable to allow thorough and uniform mixing with the equipment used for 1-pass incorporation.

Contact Monsanto Company for additional information regarding recommended equipment and methods for Surface Blend application.

2-Pass Incorporation: When 2-pass incorporation is used, shallowly incorporate the herbicide treatment into the upper 1 to 2 inches of the soil with equipment set to work the soil NO DEEPER THAN 4 INCHES. The second pass must be made at an angle and no deeper than the first pass to ensure uniform distribution of the herbicide treatment in the soil.

Preemergence Surface Application-This product and all labeled tank mixtures except Freedom plus Command™, may be applied to the soil surface after planting and prior to either crop or weed emergence. Apply within 5 days of last preplant tillage. If weeds emerge after treatment, or if treatment is applied more than 5 days after last preplant tillage, rotary hoe or shallowly cultivate immediately to improve performance. Precipitation or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone. The amount of precipitation or overhead sprinkler irrigation required depends on existing soil moisture, soil type and percent organic matter content, but 1/3 to 3/4 inch is normally adequate. Performance is improved when moisture is received within 7 days after application and prior to weed emergence. High intensity or excessive rainfall or excessive irrigation after application may reduce control.

Cultivation Information-Delay cultivation after application for as long as possible unless weeds or grasses emerge. Shallowly cultivate or rotary hoe immediately if weeds or grasses emerge. If cultivation is necessary because of soil crusting or compaction, set equipment shallow and minimize lateral soil movement to avoid dilution or displacement of the herbicide treatment.

™Surface Blend is a trademark of Monsanto Company.

™Command is a trademark of FMC Corporation.



WEEDS CONTROLLED

Applicators should evaluate soil and weed conditions carefully to ensure that they choose the lowest effective label rate. When applied as directed under conditions described, this product and tank mixtures of this product will control or reduce competition from the weeds listed.

C = Control

R = Reduced Competition

1	Freedom®	6	Freedom + Lorox™Plus
2	Freedom + Canopy™	7	Freedom + Preview™
3	Freedom + Command	8	Freedom + Pursuit™
4	Freedom + Gemini™	9	Freedom + Scepter™
5	Freedom + linuron	10	Freedom + Sencor™/Lexone™

	1	2	3	4	5	6	7	8	9	10
--	---	---	---	---	---	---	---	---	---	----

ANNUAL GRASSES

Barnyardgrass	C	C	C	C	C	C	C	C	C	C
Crabgrass: hairy, smooth	C	C	C	C	C	C	C	C	C	C
Cupgrass, woolly <sup>1</sup>	R	R	R	.	.	.	R	R	R	R
Foxtail: giant, green, robust purple, robust white, yellow	C	C	C	C	C	C	C	C	C	C
Goosegrass	C	C	C	C	C	C	C	C	C	C
Johnsongrass, seedling <sup>2</sup>	R	R	R	R	R	R	R	C	C	R
Panicum: browntop, fall	C	C	C	C	C	C	C	C	C	C
Panicum, Texas <sup>3</sup>	R	R	R	R	R	R	R	R	R	R
Rice, red <sup>4</sup>	R	R	R	.	.	.	R	C	C	R
Sandbur, Grassbur <sup>2</sup>	R	R	R	.	.	.	R	R	R	R
Shattercane, wild cane <sup>1</sup>	R	R	R	.	.	.	R	R	R	R
Signalgrass, broadleaf	C	C	C	C	C	C	C	C	C	C
Sprangletop, red	C	C	C	C	C	C	C	C	C	C
Witchgrass	C	C	C	C	C	C	C	C	C	C

ANNUAL BROADLEAVES

Beggarweed, Florida <sup>2</sup>	R	C	.	C	.	.	R	.	.	C
Carpetweed	C	C	C	C	C	C	C	C	C	C
Cocklebur	.	C	.	C	.	C	C	C	C	R
Coffeeweed:										
hemp sesbania	.	C	.	C	.	.	C	.	.	.
Copperleaf, hophornbeam	.	C	.	.	.	.	C	.	.	.
Galinsoga	C	C	C	C	C	C	C	C	C	C
Groundcherry, annual	.	C	.	C	C	C	C	C	C	C
Groundcherry, cutleaf <sup>5</sup>	C	C	C	C	C	C	C	C	C	C
Jimsonweed	.	C	C	C	.	C	C	C	C	R
Kochia	.	C	C	.	.	.	C	C	C	C
Lambsquarters <sup>5</sup>	R	C	C	C	C	C	C	C	C	C
Morningglory:										
tall <sup>6</sup> , ivyleaf <sup>6</sup> ,	.	C	.	C	.	R	R	R	R	.
entireleaf <sup>6</sup> ,	.	C	.	C	.	R	R	R	R	.
pitted,	.	C	.	C	.	R	R	R	C	.
smallflower	R	C	.	C	.	R	.	C	C	.
Mustard	.	C	.	C	C	C	C	C	C	.
Nightshade:										

18g 40

eastern black <sup>4</sup> , hairy <sup>4</sup>	C	C	C	C	C	C	C	C	C	C
Pigweed, Carelessweed	C	C	C	C	C	C	C	C	C	C
Purshlane	C	C	C	C	C	C	C	C	C	C
Pusley, Florida	C	C	C	C	C	C	C	C	C	C
Ragweed, common <sup>5</sup>	R	C	C	C	C	C	C	R	C	C
Ragweed, giant	•	R	•	R	•	R	•	R	R	•
Sicklepod	•	C	•	R	•	•	•	•	R	R
Sida, prickly; Teaweed <sup>5</sup>	R	C	C	C	C	C	C	C	C	•
Smartweed <sup>5</sup>	R	C	R	C	C	C	C	C	C	C
Starbur, bristly	R	C	•	•	•	•	•	•	C	C
Sunflower, common	•	R	•	•	•	•	R	C	C	•
Velvetleaf; Buttonweed	•	C	C	C	R	C	C	C	C	R
Waterhemp	C	C	C	C	C	C	C	C	C	C

SEDGE

Nutsedge, yellow <sup>4</sup>	C	C	C	•	•	•	C	C	C	C
-------------------------------	---	---	---	---	---	---	---	---	---	---

TM Canopy, Gemini, Lorox and Preview are trademarks of E. I. du Pont de Nemours and Company.  
 TM Command is a trademark of FMC Corporation.  
 TM Pursuit and Scepter are trademarks of American Cyanamid Company.

<sup>1</sup>Apply 4.5 quarts of this product per acre alone or in tank mixture on this weed.

<sup>2</sup>Apply 4 to 4.5 quarts of this product per acre alone or in tank mixture on this weed.

<sup>3</sup>Apply 4.5 quarts of this product per acre, followed by sprinkler irrigation, if rainfall is not expected.

<sup>4</sup>Apply 4.5 quarts of this product per acre alone or in tank mixture on this weed. When tank mixed with Pursuit or Scepter, use 3 to 4 quarts of this product per acre.

<sup>5</sup>Apply 4 to 4.5 quarts of this product per acre when used alone on this weed.

<sup>6</sup>For best results of these species when using Freedom plus Scepter tank mixture, use preplant incorporated treatments.

For scientific names for the weeds listed above, refer to the appendix at the end of this booklet.

SOYBEANS

Do not make more than one application per year or exceed a total of 4.5 quarts of Freedom per acre per year.

Do not feed forage, hay or straw from soybeans treated with this product or tank mixtures of this product. Do not ensile soybeans treated with this product.

FREEDOM®

Apply this product in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

APPROVED APPLICATION SYSTEMS

- Ground-Broadcast boom
- Aerial-Fixed wing; helicopter
- Dry Bulk Fertilizer Impregnation

Reference: The 'APPLICATION SYSTEMS' section of this label provides detailed information on the application of this product using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated-Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Irrigation within 10 days following planting may improve weed control.

Preemergence Surface-Apply this product after planting, before crop and weed emergence and within 5 days of last preplant tillage operation.

RECOMMENDED RATES: Refer to the following tables for the recommended rates for this product. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

Preplant Incorporated Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (QUARTS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse	3.5 to 4	3.5 to 4
Medium	3.5 to 4.5	3.5 to 4.5
Fine	4 to 4.5	4.5

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate in the recommended range in areas of heavy weed infestation. Use a minimum of 4.5 quarts per acre for control of yellow nutsedge.

Surface Application Rates

SOIL TEXTURAL GROUP*	BROADCAST RATE PER ACRE (QUARTS)**	
	Less than 3% organic matter	3% or more organic matter
Coarse	3 to 3.5	3 to 3.5
Medium	3 to 4.25	3 to 4.25
Fine	3 to 4.25	3.75 to 4.5

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate in the recommended range in areas of heavy weed infestations. Use a minimum of 4.5 quarts per acre for control of yellow nutsedge.

209 40

FREEDOM® plus CANOPY  
TANK MIXTURE

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the labels for Freedom and Canopy herbicides, including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

DO NOT plant rice or peanuts within 18 months after application of this tank mixture.

APPROVED APPLICATION SYSTEM

Ground-Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of the label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated-Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Irrigation within 10 days following planting may improve weed control.

Preemergence Surface-Apply this product after planting, before crop and weed emergence and within 5 days of last preplant tillage operation.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

21740

Preplant Incorporated Application Rates

BROADCAST RATES PER ACRE\*\*

SOIL TEXTURAL GROUP*	1/2% to 3% organic matter		3% to 5% organic matter	
	Freedom® + (quarts)	Canopy (ounces)	Freedom + (quarts)	Canopy (ounces)
Coarse-loamy sand, sandy				
Loam only	2.75 to 3.5	6	3 to 3.5	8
Medium***	2.75 to 3.5	8	3.5 to 4	10
Fine	3.5 to 4	10	3.5 to 4.5	12

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

\*\*Use the higher rate of Freedom herbicide in the recommended ranges in areas of heavy grass infestation, heavy crop residue or when broadleaf signalgrass is present. See 'WEEDS CONTROLLED' section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.

\*\*\*On medium soils in southeast KS, southwest MO and northeast OK with 1/2 to 3% organic matter, use 5 to 6 ounces Canopy of per acre. On silt loam soils in TN and KY, use 6 to 8 ounces of Canopy per acre.

NOTE: In the states of IA, IL, KS, MO, MN, NE, OK and WI on soils with 1/2 to 4% organic matter, reduced rates of Canopy are recommended for broadleaf weed control. Use 5 ounces of Canopy per acre on coarse soils (loamy sand and sandy loam only); 6 ounces on medium soils; and 6 to 7 ounces on fine soils. Do not apply to soils with greater than 6.8 pH or to soils with less than 1/2% organic matter or greater than 4% organic matter.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command herbicide per acre to this tank mixture.

Observe all precautions and limitations on the labels for Canopy and Command before using this tank mixture.

22740

Surface Application Rates

BROADCAST RATES PER ACRE\*\*

SOIL TEXTURAL GROUP*	1/2% to 3% organic matter		3% to 5% organic matter	
	Freedom® + (quarts)	Canopy (ounces)	Freedom + (quarts)	Canopy (ounces)
Coarse-loamy sand, sandy loam only	2.5 to 3.5	6	2.5 to 3.5	8
Medium***	2.5 to 3.5	8	3.0 to 3.75	10
Fine	3.0 to 3.75	10	3.5 to 4.0	12

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

\*\*Use the higher rate of Freedom herbicide in the recommended ranges in areas of heavy grass infestation, heavy crop residue or when broadleaf signalgrass is present. See 'WEEDS CONTROLLED' section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.

\*\*\*On medium soils in southeast KS, southwest MO and northeast OK with 1/2 to 3% organic matter, use 5 to 6 ounces Canopy per acre. On silt loam soils in TN and KY, use 6 to 8 ounces of Canopy per acre.

NOTE: In the states of IA, IL, KS, MO, MN, NE, OK and WI on soils with 1/2 to 4% organic matter, reduced rates of Canopy are recommended for broadleaf weed control. Use 5 ounces of Canopy per acre on coarse soils (loamy sand and sandy loam only); 6 ounces on medium soils; and 6 to 7 ounces on fine soils. Do not apply to soils with greater than 6.8 pH or to soils with less than 1/2% organic matter or greater than 4% organic matter.

FREEDOM® plus COMMAND  
TANK MIXTURE

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

NOTE: Observe all precautions and limitations on the labels for Freedom and Command herbicides before using this tank mixture.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom

Reference: The 'APPLICATION SYSTEMS' section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHOD

Preplant Incorporated-Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Irrigation within 10 days following planting may improve weed control.

RECOMMENDED RATES: Apply 3 to 3.5 quarts of Freedom herbicide plus 1 pint of Command 4E per acre. In areas of heavy pigweed and lambsquarters infestation, use 3.5 to 4 quarts of Freedom per acre in the tank mixture.

■ FREEDOM® plus LEXONE ■  
or  
■ FREEDOM plus SENCOR ■  
TANK MIXTURES

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

NOTE: DO NOT use on muck soils or on alkaline soils with greater than 7.4 pH. Crop injury may result if atrazine was applied the year preceding use of this tank mixture.

Observe all precautions and limitations on the labels for Freedom, Lexone and Sencor herbicides before using this tank mixture.

DO NOT REPLANT CROPS OTHER THAN SOYBEANS FOR 120 DAYS AFTER APPLICATION OF THESE TANK MIXTURES.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom

Aerial-Fixed wing helicopter

Dry Bulk Fertilizer Impregnation

Reference: The 'APPLICATION SYSTEMS' section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated-Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Irrigation within 10 days following planting may improve weed control.

Preemergence Surface-Apply this product after planting, before crop and weed emergence and within 5 days of last preplant tillage operation.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

24740

Preplant Incorporated Application Rate

BROADCAST RATES PER ACRE\*\*

SOIL TEXTURAL GROUP*	Less than 3% organic matter		3% or more organic matter	
	Freedom® + (quarts)	Sencor 4L** (pint)	Freedom + (quarts)	Sencor 4L** (pint)
Coarse-Sandy				
Loam Only	2.75 to 3.5	0.5	3 to 3.5	0.75
Medium	2.75 to 3.5	0.5 to 0.75	3.5 to 4	0.75
Fine-	3.5 to 4	0.75 to 1	3.5 to 4.5	1

\*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the rates listed in this label when using Lexone 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lexone 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

NOTE: DO NOT use on sand or loamy sand soil. Use 1.5 pints of Lexone 4L or Sencor 4F per acre in this tank mixture when applying on heavy clay soils of the Mississippi Delta. Use 4.5 quarts of Freedom herbicide in this tank mixture on medium or fine textured soils to control black or hairy nightshade.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command herbicide per acre to this tank mixture.

Observe all precautions and limitations on the labels for Lexone, Sencor and Command before using these tank mixtures.

Surface Application Rate

BROADCAST RATES PER ACRE\*\*

SOIL TEXTURAL GROUP*	Less than 3% organic matter		3% or more organic matter	
	Freedom® + (quarts)	Sencor 4L** (pint)	Freedom + (quarts)	Sencor 4L** (pint)
Coarse-Sandy				
Loam Only	2.5 to 3.5	0.5	2.5 to 3.5	0.75
Medium	2.5 to 3.5	0.5 to 0.75	3.0 to 3.75	0.75
Fine	3.0 to 3.75	0.75 to 1	3.5 to 4.0	1

\*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.



25740

\*\*Use the rates listed in this label when using Lexone 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lexone 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

NOTE: DO NOT use on sand or loamy sand soil. Use 1.5 pints of Lexone 4L or Sencor 4F per acre in this tank mixture when applying on heavy clay soils of the Mississippi Delta. Use 4.5 quarts of Freedom herbicide in this tank mixture on medium or fine textured soils to control black or hairy nightshade.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command herbicide per acre to this tank mixture. Observe all precautions and limitations on the labels for Lexone, Sencor and Command before using these tank mixtures.

**FREEDOM® plus PREVIEW  
TANK MIXTURE**

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the "WEEDS CONTROLLED" section of this label.

NOTE: Observe all precautions and limitations on the labels for Freedom and Preview, including precautions on soil pH, sensitive soybean varieties, minimum recropping interval and rotational guidelines.

**APPROVED APPLICATION SYSTEMS**

Ground-Broadcast boom

Reference: The "APPLICATION SYSTEMS" section of the label provides detailed information on the application of this tank mixture using the system selected.

**APPROVED APPLICATION METHODS**

Preplant Incorporated-Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Irrigation within 10 days following planting may improve weed control.

Preemergence Surface-Apply this product after planting, before crop and weed emergence and within 5 days of last preplant tillage operation.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury.

26740

Preplant Incorporated Application Rate

BROADCAST RATE PER ACRE\*\*

SOIL TEXTURAL GROUP*	1/2% to 3% organic matter		3% to 5% organic matter	
	Freedom® (quarts)	Preview (ounces)	Freedom® (quarts)	Preview (ounces)
Coarse-Sandy Loam Only	2.75 to 3.5	6	3 to 3.5	7
Medium	2.75 to 3.5	7	3.5 to 4	8
Fine	3.5 to 4	8	3.5 to 4.5	9

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

\*\*Use the higher rate of Freedom herbicide in the recommended ranges in areas of heavy grass infestation, heavy crop residue or when broadleaf signalgrass is present. See 'WEEDS CONTROLLED' section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.

For improved control of heavy velvetleaf infestations, add 1/2 pint of Command herbicide per acre to this tank mixture.

Observe all precautions and limitations on the labels for Preview and Command before using this tank mixture.

Surface Application Rate

BROADCAST RATE PER ACRE\*\*

SOIL TEXTURAL GROUP*	1/2% to 3% organic matter		3% to 5% organic matter	
	Freedom® (quarts)	Preview (ounces)	Freedom® (quarts)	Preview (ounces)
Coarse-Sandy Loam Only	2.5 to 3.5	6	2.5 to 3.5	7
Medium	2.5 to 3.5	7	3 to 3.75	8
Fine	3 to 3.75	8	3.5 to 4	9

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated. DO NOT APPLY TO SOIL WITH LESS THAN 1/2% organic matter.

\*\*Use the higher rate of Freedom herbicide in the recommended ranges in areas of heavy grass infestation, heavy crop residue or when broadleaf signalgrass is present. See 'WEEDS CONTROLLED' section of this label for specific rate recommendations on seedling johnsongrass, red rice and yellow nutsedge.

FREEDOM® plus PURSUIT  
TANK MIXTURE

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

NOTE: Observe all precautions and limitations on the labels for Freedom and Pursuit herbicides before using this tank mixture including precautions on minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom

Aerial-Fixed wing; helicopter

Reference: The 'APPLICATION SYSTEMS' section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated-Apply this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Irrigation within 10 days following planting may improve weed control.

Preemergence Surface-Apply this product after planting, before crop and weed emergence and within 5 days of last preplant tillage operation.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury. See the 'WEEDS CONTROLLED' section of this label for recommended rates and application methods for control of certain weeds.

BROADCAST RATES PER ACRE\*\*

SOIL TEXTURAL GROUP*	Less than 3% organic matter	3% or more organic matter	
	Freedom® (quarts)	Freedom (quarts)	Pursuit (ounces)
Coarse	2.5 to 3	2.5 to 3	4
Medium	2.5 to 3	2.75 to 3.25	4
Fine	2.75 to 3.25	3 to 3.5	4

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate of Freedom herbicide in the recommended ranges in areas of heavy grass infestations, heavy crop residue or when the following weeds are present: fall panicum, barnyardgrass, broadleaf signalgrass, seedling johnsongrass, red rice or yellow nutsedge.

FREEDOM® plus SCEPTER  
TANK MIXTURE

Apply this tank mixture in water or sprayable fluid fertilizer solution for control of yellow nutsedge and the annual grasses and broadleaf weeds listed in the 'WEEDS CONTROLLED' section of this label.

NOTE: Observe all precautions and limitations on the labels for Freedom and Scepter herbicides before using this tank mixture including precautions on minimum recropping interval and rotational guidelines.

APPROVED APPLICATION SYSTEMS

Ground-Broadcast boom

Aerial-Fixed wing; helicopter

Reference: The 'APPLICATION SYSTEMS' section of this label provides detailed information on the application of this tank mixture using the system selected.

APPROVED APPLICATION METHODS

Preplant Incorporated-Applly this product within 7 days prior to planting and shallowly incorporate or Surface Blend into the upper 1 to 2 inches of soil. Irrigation within 10 days following planting may improve weed control.

Preemergence Surface-Applly this product after planting, before crop and weed emergence and within 5 days of last preplant tillage operation.

RECOMMENDED RATES: Refer to the following table for the recommended broadcast treatment rates for this tank mixture. Applications which are not consistent with recommendations in this label may result in unsatisfactory weed control or crop injury. See the 'WEEDS CONTROLLED' section of this label for recommended rates and application methods for control of certain weeds.

BROADCAST RATES PER ACRE\*\*

SOIL TEXTURAL GROUP*	Less than 3% organic matter	3% or more organic matter	SCEPTER*** (pints)
	Freedom® (quarts)	Freedom (quarts)	
Coarse	2.5 to 3	2.5 to 3	1/2 to 2/3
Medium	2.5 to 3	2.75 to 3.25	1/2 to 2/3
Fine	2.75 to 3.25	3 to 3.5	1/2 to 2/3

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate of Freedom herbicide in the recommended ranges in areas of heavy grass infestation, heavy crop residue or when the following weeds are present: fall panicum, barnyardgrass, broadleaf signalgrass, seedling johnsongrass, red rice or yellow nutsedge.

\*\*\*Use the higher rates of Scepter in the recommended ranges when red rice, sicklepod, Florida beggarweed or velvetleaf are present.

For improved control of heavy velvetleaf infestation, add 1/2 pint of Command 4E herbicide per acre to this tank mixture.

Observe all precautions and limitations on the labels for Scepter and Command before using these tank mixtures.

Conservation or Minimum

Tillage Systems

TANK MIXTURES

SOYBEANS ONLY

When applied as directed under the conditions described, these tank mixtures control many emerged annual weeds, suppress many emerged perennial weeds and give preemergence control of many annual grasses and weeds when soybeans will be planted directly into a cover crop, stale seedbed, or in previous crop residues such as wheat stubble.

These tank mixtures will not control regrowth from perennial weeds.

DO NOT APPLY BY AIR.

For mixing instructions, see the 'MIXING, SPRAYING AND HANDLING INSTRUCTIONS' section of this label.

Freedom and the following tank-mixtures can be tank-mixed with Roundup or Gramoxone Extra.

- FREEDOM® ■
- FREEDOM plus CANOPY ■
- FREEDOM plus GEMINI ■
- FREEDOM plus LEXONE ■
- FREEDOM plus SENCOR ■
- FREEDOM plus LINURON ■
- FREEDOM plus LOROX PLUS ■
- FREEDOM plus PREVIEW ■
- ~~FREEDOM plus SENCOR~~ ■
- FREEDOM plus PURSUIT ■
- FREEDOM plus SCEPTER ■
- ~~FREEDOM plus SENCOR~~ ■

Apply the tank mixtures with Roundup in 10 to 20 gallons of water or 10 to 60 gallons of nitrogen solution per acre, or the tank mixtures with Gramoxone Extra in 20 to 60 gallons of water or clear liquid fertilizer per acre immediately before, during or after planting, but BEFORE CROP EMERGENCE. As density of stubble, crop residue or weeds increases, spray gallonage and rate should be increased within the recommended ranges to ensure complete coverage. In the absence of emerged vegetation delete the Roundup or Gramoxone Extra portion of these mixes.

CONTROL OR SUPPRESSION OF EMERGED WEEDS

- ROUNDUP® ■

30740

Annual Weeds—Apply 1 to 2 pints of Roundup per acre in these tank mixtures when weeds are less than 6 inches tall. Use 2 pints of Roundup per acre for control of fall panicum, barnyardgrass, crabgrass and shattercane up to 2 inches tall; Pennsylvania smartweed up to 6 inches tall. 3 pints on other annual weeds 6 to 12 inches tall. For emerged annual weeds controlled, see the "WEEDS CONTROLLED" section of the label for Roundup.

Perennial Weeds—At normal application dates in minimum tillage systems, perennial weeds may not be at the proper stage of growth for control. See the "GENERAL INFORMATION" section of the label for Roundup for the proper stage of growth for perennial weeds. Use of 2 to 4 quarts of Roundup per acre in this tank mixture under these conditions provides top kill and reduces competition from many emerged perennial grasses and broadleaf weeds. For perennial weeds controlled, see the "WEEDS CONTROLLED" section of the label for Roundup.

To obtain control of perennial weeds, follow recommendations on the label for Roundup for stage of growth and rate of application.

USE OF THIS TANK MIXTURE FOR BERMUDAGRASS OR JOHNSONGRASS CONTROL IS NOT RECOMMENDED.

NOTE: When using these tank mixtures, do not exceed 4 quarts of Roundup herbicide per acre.

Ammonium Sulfate—The addition of ammonium sulfate in the spray solution may increase the performance of Roundup tank mixtures on emerged annual weeds under adverse growing conditions. When using ammonium sulfate, add 2 percent dry ammonium sulfate by weight or 17 pounds per 100 gallons of water. Ammonium sulfate should be added to the water in the spray tank and completely dissolved prior to adding the herbicide or surfactant. Do not mix ammonium sulfate in fluid fertilizer solutions.

If ammonium sulfate is added directly to the spray tank, add slowly with agitation. Adding too quickly may clog outlet lines. Nozzle tip plugging may result from the use of low-quality ammonium sulfate. To determine quality, perform a jar test by adding 1/3 cup of ammonium sulfate to 1 gallon of water and agitate for one minute. If undissolved sediment is observed, predissolve the ammonium sulfate in water and filter prior to adding to the spray tank.

Following use of ammonium sulfate, thoroughly rinse the spray system with clean water to reduce corrosion.

Surfactants—Nonionic surfactants which are labeled for use with herbicides may be used with Roundup. Do not reduce rates of Roundup when adding surfactant. Use 0.5 percent surfactant concentrate (2 quarts per 100 gallons of spray solution) when using surfactants which contain at least 50 percent active ingredient or a 1 percent surfactant concentration (4 quarts per 100 gallons of spray solution) for those surfactants containing less than 50 percent active ingredient. Read and carefully observe surfactant cautionary statements and other information appearing on the surfactant label.

GRAMOXONE EXTRA

When used as directed, Gramoxone Extra in tank-mixture with labeled Freedom® tank-mixtures controls many emerged annual weeds and suppresses many emerged perennial weeds.

Broadcast Treatment—Apply 1.5 to 3 pints of Gramoxone Extra in this tank mixture immediately before, during or after planting but BEFORE CROP EMERGENCE. Use 1.5 to 2 pints when weeds are 1 to 3 inches tall, 2 to 2.5

pints when weeds are 3 to 6 inches tall and 2.5 to 3 pints when weeds are 6 inches tall. This mixture may not control weeds taller than 6 inches. As density of stubble, crop residue or weeds increases, spray gallonage should be increased within the recommended range for complete coverage. Add a nonionic spreader surfactant (approved for use in crops) containing at least 75 percent surface active agent at 8 fluid ounces per 100 gallons of diluted spray. REFER TO THE LABEL FOR GRAMOXONE EXTRA FOR PRECAUTIONARY STATEMENTS.

PREEMERGENCE WEED CONTROL

FREEDOM®

For weeds controlled preemergence, see the 'WEEDS CONTROLLED' section of this label.

See the following table for recommended rates of Freedom on various soil types.

BROADCAST RATE PER ACRE (QUARTS)	
SOIL TEXTURAL GROUP*	Freedom®
Coarse	3.75
Medium	4.5
Fine	4.5

\*Refer to the 'SOIL TEXTURE' section of the label to determine the corresponding soil textural group for the soil to be treated.

FREEDOM® plus CANOPY

For weeds controlled preemergence see the 'WEEDS CONTROLLED' section of this label for Freedom® plus Canopy.

NOTE: Observe all precautions and limitations on the Freedom and Canopy labels including precautions on soil pH, sensitive soybean varieties and rotational crop restrictions.

See the following table for recommended rates of Freedom and Canopy in this tank mixture.

BROADCAST RATE PER ACRE\*\*

SOIL TEXTURAL GROUP*	Freedom® (quarts)	+	Canopy (ounces)
Coarse-sandy loam only	2.5 to 3.5		6
Medium	3 to 3.75		8
Fine	3.75 to 4.5		10

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate of Freedom and Canopy in areas of heavy weed infestation.

■ FREEDOM® plus GEMINI ■

For weed controlled preemergence, see the 'WEEDS CONTROLLED' section of this label for Freedom plus Gemini.

NOTE: Observe all precautions and limitations on the Freedom and Gemini labels including precautions on soil pH and rotational crop restrictions.

See the following table for recommended rates of Freedom and Gemini in this tank mixture.

BROADCAST RATE PER ACRE\*\*

SOIL TEXTURAL GROUP*	Freedom® (quarts)	+	Gemini (ounces)
Coarse-sandy loam only	2.5 to 3.5		12 to 16
Medium	3 to 3.75		16 to 20
Fine	3.75 to 4.5		20 to 24

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate of Freedom and Gemini in areas of heavy weed infestation.

■ FREEDOM® plus LEXONE ■

or

■ FREEDOM plus SENCOR ■

For weeds controlled preemergence, see the 'WEEDS CONTROLLED' section of this label for Freedom plus Lexone 4L or Sencor 4F.

See the following table for recommended rates of Freedom plus Lexone 4L or Freedom plus Sencor 4F in this tank mixture on various soil types.

BROADCAST RATE PER ACRE\*\*



SOIL TEXTURAL GROUP*	Freedom® (quarts)	+	Lexone 4L** or Sencor 4F** (pints)
Coarse-sandy loam only	2.5 to 3.5		0.5 to 0.75
Medium	3 to 3.75		0.75 to 1
Fine	3.75 to 4.5		1 to 1.5***

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the rates listed in this label when using Lexone 4L or Sencor 4F. Use the equivalent rates when using the 50WP or DF formulations of these products. One pint of Lexone 4L or Sencor 4F equals one pound of the corresponding 50WP formulation or 2/3 pound of the corresponding DF formulation.

\*\*\*On the heavy clay soil of the Mississippi Delta, use 1.5 pints of Lexone 4L or Sencor 4F per acre.

Use the higher rate of Freedom in the recommended ranges in areas of heavy grass infestations or when fall panicum or crabgrass will be present.

Use the higher rate of Lexone or Sencor in the recommended ranges on soils with greater than 3% organic matter.

DO NOT use this mixture on sand or loamy sand soils as crop injury from Lexone or Sencor may occur.

DO NOT use on muck soils.

DO NOT apply on alkaline soils with a pH greater than 7.4.

Crop injury may occur if atrazine was applied the year before use of this tank mixture.

DO NOT REPLANT CROPS OTHER THAN SOYBEANS FOR 120 DAYS AFTER APPLICATION OF THESE TANK MIXTURES.

■ FREEDOM® plus LINURON ■

For weeds controlled preemergence, see the 'WEEDS CONTROLLED' section of this label for Freedom plus linuron.

See the following table for recommended rates of Freedom and linuron in this tank mixture.

347 40

BROADCAST RATE PER ACRE\*\*

SOIL TEXTURAL GROUP*	Freedom® (quarts)	+	linuron 4L** (quarts)
Coarse-sandy loam only	2.5 to 3.5		0.5 to 0.75
Medium	3 to 3.75		0.75 to 1
Fine	3.75 to 4.5		1 to 1.5

\*Refer to the "SOIL TEXTURE" section of the label to determine the corresponding textural group for the soil to be treated.

\*\*Use the rates listed on this label when using linuron 4L. Use equivalent rates when using linuron 50WP or 50% dry flowable formulation. One quart of linuron 4L equals 2 pounds of linuron 50WP or 2 pounds of linuron 50% dry flowable.

Use the higher rate of Freedom in the recommended ranges in areas of heavy grass infestation or when fall panicum or crabgrass will be present.

Use the higher rate of linuron in the recommended ranges on soils with greater than 3% organic matter.

DO NOT use this tank mixture on sand or loamy sand or on soil with less than 1% organic matter as crop injury from linuron may occur.

FREEDOM® plus LOROX PLUS

For weeds controlled preemergence see the "WEEDS CONTROLLED" section of this label for Freedom plus Lorox Plus.

NOTE: Observe all precautions and limitations on the labels for Freedom and Lorox Plus, including precautions on soil pH and rotational crop restrictions.

See the following table for recommended rates of Freedom and Lorox Plus in this tank mixture.

BROADCAST RATE PER ACRE\*\*

SOIL TEXTURAL GROUP*	Freedom® (quarts)	+	Lorox Plus (ounces)
Coarse-sandy loam only	2.5 to 3.5		12 to 14
Medium	3 to 3.75		14 to 16
Fine	3.75 to 4.5		16 to 18

\*Refer to the "SOIL TEXTURE" section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate of Freedom and Lorox Plus in areas of heavy weed infestation.

FREEDOM® plus PREVIEW

For weeds controlled preemergence see the 'WEEDS CONTROLLED' section of this label for Freedom plus Preview.

NOTE: Observe all precautions and limitations on the labels for Freedom and Preview, including precautions on soil pH, sensitive soybean varieties and rotational crop restrictions.

See the following table for recommended rates of Freedom and Preview in this tank mixture.

BROADCAST RATE PER ACRE**		
SOIL TEXTURAL GROUP*	Freedom® (quarts)	Preview (ounces)
Coarse-sandy loam only	2.5 to 3.5	6
Medium	3 to 3.75	7
Fine	3.75 to 4.5	8

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding soil textural group for the soil to be treated.

\*\*Use the higher rate of Freedom and Preview in areas of heavy weed infestation.

FREEDOM® plus PURSUIT

For weeds controlled preemergence, see the 'WEEDS CONTROLLED' section of this label for Freedom plus Pursuit.

NOTE: Observe all precautions and limitations on the Freedom and Pursuit labels before use of this tank mixture including minimum recropping interval and rotational guidelines.

See the following table for recommended rates of Freedom and Pursuit in this tank mixture.

BROADCAST RATE PER ACRE**		
SOIL TEXTURAL GROUP*	Freedom® (quarts)	Pursuit (ounces)
Coarse	2.5 to 3.5	4
Medium	3 to 3.75	4
Fine	3.75 to 4.5	4

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding textural group for the soil to be treated.

\*\*Use the higher rate of Freedom in areas of heavy weed infestations.

36740

FREEDOM® plus SCEPTER

For weeds controlled preemergence, see the 'WEEDS CONTROLLED' section of this label for Freedom plus Scepter.

NOTE: Observe all precautions and limitations on the Freedom and Scepter labels before use of this tank mixture including minimum recropping interval and rotational guidelines.

See the following table for recommended rates of Freedom and Scepter in this tank mixture.

BROADCAST RATE PER ACRE**		
SOIL TEXTURAL GROUP*	Freedom® (quarts)	Scepter (pints)
Coarse	2.5 to 3.5	2/3
Medium	3 to 3.75	2/3
Fine	3.75 to 4.5	2/3

\*Refer to the 'SOIL TEXTURE' section of this label to determine the corresponding textural group for the soil to be treated.

\*\*Use the higher rate of Freedom in areas of heavy weed infestations.

APPENDIX

LIST OF COMMON AND SCIENTIFIC NAME OF ANNUAL GRASSES AND BROADLEAF WEEDS

ANNUAL GRASSES

Barnyardgrass  
*Echinochloa crus-galli*

Crabgrass  
*Digitaria ischaemum*  
*Digitaria sanguinalis*

Cupgrass, woolly  
*Eriochloa villosa*

Foxtail, giant  
*Setaria faberi*

Foxtail, green, robust  
purple, robust white  
*Setaria viridis*

Foxtail, yellow  
*Setaria lutescens*

Goosegrass  
*Eleusine indica*

Johnsongrass  
*Sorghum halepense*

Panicum, browntop  
*Panicum fasciculatum*

Panicum, fall  
*Panicum*  
*dichotomiflorum*

Panicum, Texas  
*Panicum texanum*

Rice, red  
*Orvza sat\_vva*

Sandbur, Grassbur  
*Cenchrus rcertus*

Shattercane wild cane  
*Sorghum b color*

Signalgrass broadleaf  
*Brachiaria platyphylla*

Sprangletop red  
*Leptochloa filiformis*

Witchgrass

38740

Panicum capillare

ANNUAL BROADLEAVES

Beggarweed, Florida  
*Desmodium tortuosum*

Carpetweed  
*Mollugo verticillata*

Cocklebur  
*Xanthium strumarium*

Coffeeweed  
*Hesperis sesbania*  
*Sesbania exaltata*

Copperleaf, hophornbeam  
*Acalypha spp*

Galinsoga  
*Galinsoga spp*

Groundcherry, annual  
*Physalis spp*

Groundcherry, cutleaf  
*Physalis angulata*

Jimsonweed  
*Datura stramonium*

Kochia  
*Kochia scoparia*

Lambsquarters  
*Cenopodium album*

Morningglory  
Tall  
*Ipomoea purpurea*

Pitted  
*Ipomoea lacunosa*

Ivyleaf  
*Ipomoea nederacea*

Entireleaf  
*Ipomoea hederacea*  
var *intergriuscula*

Smallflower  
*Jacquemontia*  
*tamnifolia*

Mustard

Brassica spp

Nightshade black  
*Solanum nigrum*

Nightshade, hairy  
*Solanum sarrachoides*

Pigweed, Carelessweed  
*Amaranthus spp*

Purslane  
*Portulaca oleracea*

Pusley, Florida  
*Richardia scabra*

Ragweed, common  
*Ambrosia artēmisifolia*

Ragweed, giant  
*Ambrosia trifida*

Sicklepod  
*Cassia obtusifolia*

Sida, prickly Teaweed  
*Sida spinosa*

Smartweed  
*Polygonum pensylvanicum*  
*Polygonum persicaria*

Starbur, bristly  
*Acanthospermum hispidum*

Sunflower, common  
*Helianthus annuus*

Velvetleaf, Buttonweed  
*Abutilon theophrasti*

Waterhemp  
*Amaranthus tuberculatos*

SEDGE

Nutsedge yellow  
*Cyperus esculentus*

STATE REGISTRATIONS

Freedom® herbicide has been approved by the U.S. Environmental Protection Agency for the crop, uses and sites listed in this label. This product is registered for use in the following states:

Alabama	Kentucky	Oklahoma
Arkansas	Louisiana	South Carolina
Colorado	Maryland	South Dakota
Delaware	Minnesota	Tennessee
Georgia	Mississippi	Texas
Illinois	Missouri	Virginia
Indiana	Nebraska	West Virginia
Iowa	North Carolina	Wisconsin
Kansas	North Dakota	

EPA Reg. No. 524-422

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

©MONSANTO COMPANY, 1997/1996

1201302-2/53

MONSANTO COMPANY  
ST. LOUIS, MISSOURI 63167 U.S.A.