

RAMROD® + ATRAZINE FG  
FLOWABLE GRANULAR HERBICIDE  
BY MONSANTO

Selective herbicide for preemergence weed control in grain sorghum (milo), field corn, hybrid seed corn, silage corn and sweet corn.

Complete Directions for Use

Ramrod® is a registered trademark of Monsanto Company.


EPA Reg. No. 524-

1989-1

848.48-000.01/53

Read each of these sections of this label for essential product performance information.

USE THESE CONVENIENT SECTION MARKERS.



PRECAUTIONARY STATEMENTS

GENERAL INFORMATION

MIXING AND SPRAYING INSTRUCTIONS


FLUID FERTILIZER COMPATIBILITY TEST

APPLICATION INFORMATION

CULTIVATION INFORMATION

WEEDS CONTROLLED

CROPS



Read the entire label before using this product.

Use only according to label instructions.

Read "LIMIT OF WARRANTY AND LIABILITY" before buying or using. If terms are not acceptable, return at once unopened.

NOT FOR REFORMULATION OR REPACKAGING.

LIMIT OF WARRANTY AND LIABILITY


This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OF MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application ranges set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range specified in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE ELECTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.



PRECAUTIONARY STATEMENTS

Hazards to  
Humans and Domestic Animals

Keep out of reach of children.

DANGER!

- CAUSES EYE BURNS.
- HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN.
- CAUSES NOSE AND THROAT IRRITATION.
- MAY CAUSE SKIN IRRITATION.
- MAY CAUSE ALLERGIC SKIN REACTION.

Do not get in eyes, on skin or on clothing.  
 Wear goggles, face shield or safety glasses and rubber gloves in transferring and mixing and when adjusting, repairing or cleaning equipment.  
 Do not breathe dust.  
 Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

IF ON SKIN, immediately flush with plenty of water while removing contaminated clothing. Get medical attention. As soon as soap is available, wash skin thoroughly with soap and water. Wash clothing before reuse. Sensitized persons should avoid further contact and reuse of contaminated clothing.

IF INHALED, remove individual to fresh air. Get medical attention if breathing difficulty develops.

IF SWALLOWED, induce vomiting immediately by giving two glasses of water and sticking finger down throat. Get medical attention. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

---

In case of an emergency involving this product,  
 Call Collect, day or night, (314) 694-4000.

---

Environmental Hazards

This chemical has been identified in limited ground water sampling and there is the possibility that it can leach through the soil to ground water, especially where soils are coarse and ground water is near the surface. Following application, and during rainfall events that cause runoff, this chemical may reach surface water bodies including streams, rivers and reservoirs.

Care must be taken when using this product to prevent back-siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Check valves or antisiphoning devices must be used on all mixing and/or irrigation equipment. Disposal of excess pesticide, spray mixtures or rinsate should be according to label use instructions or according to the state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office.

This product is toxic to fish. Do not apply directly to water or wetlands. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, collect for use or properly dispose of granules.

ACTIVE INGREDIENTS:

Propachlor, (2-chloro-N-isopropylacetanilide) . . . . .	48.0%
Atrazine, [2-Chloro-4-(ethylamino)-6-(isopropylamino)-s-triazine]. . . . .	15.5%
Related atrazine compounds. . . . .	0.5%
INERT INGREDIENTS: . . . . .	36.0%
	100.0%

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

STORAGE:

Keep bag closed to prevent spills and contamination.

DISPOSAL:


Wastes of this pesticide may cause eye burns and may be dangerous. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Emptied container retains product residue. Observe all labeled safeguards until container is destroyed.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

This herbicide mixture is recommended for control of many annual grasses and certain broadleaf weeds. For best results, it should be applied to the soil surface before crops or weeds emerge. The seedbed should be fine, firm and free of clods and trash; application should be made within 5 days after the last tillage for weed control. Do not apply when conditions favor drift.



Precipitation or overhead sprinkler irrigation is required after application to move the herbicide treatment into the weed germination zone. The amount of precipitation or overhead sprinkler irrigation required depends on existing soil mixture, soil type and percent organic matter content, but 1/3 to 3/4 inch is normally adequate. Performance is improved when moisture is received within 7 days after application and prior to weed emergence. High intensity or excessive rainfall or excessive irrigation after application may reduce control.

NOTE: Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

Flush sprayer with clean water after use and dispose of rinsate according to labeled use or disposal instructions.

Do not use this product in fields where flood irrigation is to be practiced.

DO NOT apply in sprayable fluid fertilizer after crop emergence since crop injury may result.

Do not graze or feed sorghum (milo) forage, silage or fodder (stubble) from treated fields to dairy animals.

PLANT ONLY CORN, SORGHUM OR SOYBEANS THE YEAR FOLLOWING THE USE OF THIS TANK MIXTURE. IF SOYBEANS ARE TO BE PLANTED, THERE IS A POSSIBILITY OF CROP INJURY DUE TO CARRYOVER OF ATRAZINE. Do not plant soybeans in areas where furrow irrigation is practiced.

In Texas, on grain sorghum, use only on the fine-textured soils of the Gulf Coast.


Applications made to sorghum growing on alkali soils or where cuts, fills or erosions have exposed calcareous or alkali subsoils may result in crop injury.

### MIXING AND SPRAYING INSTRUCTIONS

Always predetermine the compatibility of this product with water carrier or sprayable fluid fertilizer carrier by mixing small proportional quantities in advance. See the "Standard Sprayable Fluid Fertilizer Compatibility Test" section on this label.

Mix this product with the appropriate carrier as follows:

1. Place a 20 to 35 mesh screen or wetting basket over filling port.
2. Through the screen, fill the sprayer tank one-half full with the appropriate carrier.
3. If a compatibility agent is necessary to improve mixing or to prevent the formation of undesirable and unsprayable gels or precipitates, while agitating add it to the carrier already in the tank.



Use only compatibility agents cleared by the Federal Food and Drug Administration for this use. Follow the directions for use on the compatibility agent label. Check for adequate agitation.

- 4. Add the required amount of this product. Continue agitation.
- 5. Complete filling the sprayer tank with carrier. Remove hose from tank immediately after filling to avoid siphoning back into carrier source. Maintain good agitation at all times until the contents of the tank are sprayed, otherwise settling, which can lead to clogging, may occur. Close tank valves and drain all lines when spraying is delayed.

Keep by-pass line on or near the bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select proper nozzle to avoid spraying a fine mist. For best results, use flat fan or flood jet nozzles. Check for even distribution of spray droplets. To reduce loss of the chemical due to drift of a fine mist, apply at pressures less than 40 psi.

STANDARD SPRAYABLE  
FLUID FERTILIZER  
COMPATIBILITY TEST

Herbicides may not always mix evenly throughout a sprayable fluid fertilizer or the components may separate too quickly to make their combined use of practical value. This may be due to certain characteristics of the different fluid fertilizers. A simple test using small quantities of the components is suggested to provide compatibility potential.


The test follows:

A. Materials Required For A Compatibility Test

- 1. Two one-quart jars with lid or stopper (marked "with" and "without").
- 2. TEAspoons (for a more exacting test, a five to ten milliliter (ml.) pipette or graduated cylinder is desirable).
- 3. A minimum of one quart of the sprayable fluid fertilizer to be tested.
- 4. The herbicide(s) to be mixed.
- 5. A compatibility agent (the purpose of the adjuvant is to help keep the fertilizer and herbicide(s) in suspension, if this assistance is needed).

B. Procedure

- 1. Add one pint of the sprayable fluid fertilizer that will be used or other herbicide carrier to each jar marked "with" and "without".



Add One Pint Liquid Fertilizer  
To Two Quart Jars.

WITH

WITHOUT

- 2. To the jar marked "with", add 1/4 TEAspoon or 1.2 milliliters of a suitable compatibility agent; shake gently for five to ten seconds to mix. (1/4 TEAspoon in one pint is the equivalent of two pints per 100 gallons of liquid fertilizer.)

To Jar Marked "With"  
Add Compatibility Agent  
And Shake to Mix.

WITH

WITHOUT

- 3. To each jar add the appropriate amount of herbicide(s). If more than one is used, add them separately with the wetttable powders added first, flowables second and liquid last. Shake gently five to ten seconds after each addition.

Add Herbicide(s) To Both Jars  
And Shake to Mix.

WITH

WITHOUT

Amount to Be  
Added Per Pint Of  
Sprayable fluid Fertilizer  
(Assuming Volume is 25 gal/A)

HERBICIDE	RATE/A	Level TEAspoons
This Product	1 lb.	= 1.5
	2 lb.	= 3
	3 lb.	= 4.5
	4 lb.	= 6
	5 lb.	= 7.5

This compatibility test is designed for 25 gallons of spray per acre with the maximum labeled rate of herbicide. For changes in spray volume or herbicide concentration, make appropriate changes in the ingredients of the test. Regardless of spray volume, the amount of compatibility agent should be equal to two or three pints (two pints = 1/4 TEAspoon or 1.2 milliliters, three pints = 3/8 TEAspoon or 1.8 milliliters per pint of sprayable fluid fertilizer) per 100 gallons of liquid fertilizer.

C. Observations and Decisions

1. If the herbicide(s) and the sprayable fluid fertilizer are compatible.
2. If a compatibility agent is necessary.

Five minutes after the final addition and mixing, observe both jars for the formation of large flakes, sludge, gels or other precipitates. Observe if the herbicide(s) cannot be physically mixed with the liquid fertilizer but remains as small oily particles in the solution.

If incompatibility in any form described above occurs in the jar "with" the compatibility agent added, the liquid fertilizer and the herbicide(s) should not be used together in the same spray tank.

If incompatibility as described above occurs in the jar "without" the adjuvant but not in the jar "with" adjuvant, the use of a compatibility adjuvant is recommended.

Both jars should be allowed to stand and be observed periodically for one-half hour. If the separate layers of liquid fertilizer and additives can be resuspended by shaking, commercial application is possible. Wettable powders will either settle to the bottom of the tank or jar, or float to the top, depending upon the density of the fertilizers.


If this product is compatible with fluid fertilizer in the foregoing test without having to use a compatibility agent, fluid fertilizer may be used for the premixing. If it is not compatible, the herbicide(s) should be premixed with water before adding to the spray tank.

APPLICATION INFORMATION  
APPLICATION SYSTEMS

Ground Broadcast Treatment--Apply this product in 15 or more gallons of water or sprayable fluid fertilizer per acre. Do not apply during periods of gusty winds or winds in excess of 15 MPH.

Band Treatment--Apply a broadcast equivalent rate and volume per acre. To determine these:

Band width in inches	X	Broadcast RATE per acre	=	Band RATE per acre
----- Row width in inches				
Band width in inches	X	Broadcast VOLUME per acre	=	Band VOLUME per acre
----- Row width in inches				





Mount spray equipment on the planter with nozzles behind press wheel and apply this product at planting time.

NOTE: For band application nozzle placement, refer to the nozzle manufacturer's specification as a guide for determining proper nozzle height.

### AERIAL APPLICATION

Fixed wing; helicopter - Apply the recommended rates of this product in 3 to 10 gallons of water per acre. AVOID DRIFT - DO NOT APPLY DURING INVERSION CONDITIONS, WHEN WINDS ARE GUSTY, OR UNDER ANY OTHER CONDITION WHICH WILL ALLOW DRIFT. Coarse sprays are less likely to drift, therefore, do not use nozzles or nozzle configurations which dispense spray or fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure. Ensure uniform application. To avoid streaked, uneven or overlapped application use mechanical marking systems only. HUMAN FLAGGERS PROHIBITED!

### CULTIVATION INFORMATION

As long as weeds are controlled, delay cultivation as long as possible.

If weed seedlings emerge, shallowly cultivate or rotary hoe to improve weed control.

If cultivation or rotary hoeing is necessary because of soil crusting or compaction, tillage should be shallow to minimize dilution and displacement of this product.

### WEEDS CONTROLLED

When applied as directed, under the conditions described, this product CONTROLS the following weeds:

Barnyardgrass  
*Echinochloa crus-galli*


Buckwheat, wild  
*Polygonum convolvulus*

Carpetweed  
*Mollugo verticillata*

Crabgrass  
*Digitaria spp.*

Foxtail, giant  
*Setaria faberi*

Foxtail, green  
*Setaria viridis*



Foxtail, yellow  
*Setaria lutescens*

Goosegrass  
*Eleusine indica*

Groundsel  
*Senecio vulgaris*

Kochia  
*Kochia scoparia*

Lambsquarters  
*Chenopodium album*

Mustard  
*Brassica spp.*

Nightshade, black  
*Eleusine indica*

Nightshade, hairy  
*Solanum sarrachoides*

Panicum, fall  
*Panicum dichotomiflorum*

Pigweed  
*Amaranthus spp.*

Purslane  
*Portulaca oleracea*

Pusley, Florida  
*Richardia scabra*


Ragweed  
*Ambrosia artemisiifolia*

Ryegrass, annual  
*Lolium spp.*

Smartweed  
*Polygonum pennsylvanicum*  
*Polygonum persicaria*

When applied immediately after planting, under the conditions described, this product at a rate of 6.25 to 8.6 pounds per acre REDUCES COMPETITION. From the following hard to control weeds:

Buttonweed, Velvetleaf  
*Abutilon theophrasti*



Cocklebur  
Xanthium  
strumarium


Jimsonweed  
Datura stramonium

Morningglory, annual  
Ipomoea purpurea

Signalgrass, broadleaf  
Brachiaria platyphylla

### PREEMERGENCE SURFACE TREATMENT

For control of listed annual grasses and broadleaf weeds, apply this product in 15 or more gallons of water or sprayable fluid fertilizer per acre on the soil surface any time from immediately after planting and before the crop or weeds emerge. See the following table for recommended rates. In areas of heavy grass or broadleaf weed infestations or when listed hard to control weeds are present, use the higher rate in the recommended ranges.



05900084

BROADCAST RATE PER ACRE (POUNDS)

<u>SOIL TEXTURAL GROUP</u>	<u>Less than 3% organic matter</u>	<u>More than 3% organic matter</u>
COARSE*		
Sand thru sandy loam	5.5	6.25
MEDIUM		
Loam thru sandy clay loam	6.25	7
FINE		
Silty clay loam thru clay	6.25 to 7	7 to 8.6

\*Do not use on grain sorghum on coarse texture soils with less than 3 percent organic matter.

For Corn Only--This product can be applied until weeds reach the 2-leaf stage but prior to corn emergence. Applications beyond the 2-leaf stage will not provide satisfactory results. Better results will be obtained if rainfall occurs within 5 to 7 days after application. If rainfall does not occur, use shallow cultivation with a rolling cultivator or similar implement to remove emerged weeds. Avoid cultivation methods which will displace the treated soil layer. See the rate table in the preemergence treatment section of this label for the recommended rate for the soil texture and organic matter content to be treated.


For Lister or Deep Furrow Planted Corn--Apply this product at the higher broadcast rate listed in the rate table in the preemergence section of this label for the soil texture and organic matter content to be treated.

If heavy rains follow application, the treatment on the top and sides of the furrow may not provide weed control as good as that obtained in the bottom of the furrow.

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

MONSANTO COMPANY 1989

MONSANTO COMPANY  
AGRICULTURAL PRODUCTS  
ST. LOUIS, MISSOURI 63167 U.S.A.



0600084

# RAMBROD

## + ATRAZINE FG


FLOWABLE GRANULAR HERBICIDE BY

**Monsanto**

Selective herbicide for preemergence weed control in grain sorghum (milo), field corn, hybrid seed corn, silage corn and sweet corn.

**DANGER!**  
 Read the entire label before using this product.  
 See the directions on the label for use.  
 Read the label on the container and label for use.  
 Mixing or using in other than the recommended amount is  
 not approved.  
 See the label for use on other crops.

<b>ACTIVE INGREDIENTS</b>	
Propachlor, (2-chloro-N-isopropylacetanilide) . . .	48.0%
Atrazine, [2-Chloro-4-(ethylamino)-6-(isopropylamino)-s-triazine] . . . . .	15.5%
Related atrazine compounds . . . . .	0.5%
<b>INERT INGREDIENTS</b> . . . . .	36.0%
	<b>100.0%</b>



**NET 18.75 LBS**

# + ATRAZINE FG

FLOWABLE GRANULAR HERBICIDE BY

**Monsanto**

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

Keep out of reach of children.

### DANGER!

CAUSES EYE BURNS.  
HARMFUL IF SWALLOWED OR ABSORBED THROUGH SKIN.  
CAUSES NOSE AND THROAT IRRITATION.  
MAY CAUSE SKIN IRRITATION.  
MAY CAUSE ALLERGIC SKIN REACTION.

Do not get in eyes, on skin or on clothing.  
Wear goggles, face shield or safety glasses and rubber gloves in transferring and mixing and when adjusting, repairing or cleaning equipment.  
Do not breathe dust.  
Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

IF ON SKIN, immediately flush with plenty of water while removing contaminated clothing. Get medical attention. As soon as soap is available, wash skin thoroughly with soap and water. Wash clothing before reuse. Sensitized persons should avoid further contact and reuse of contaminated clothing.

IF INHALED, remove individual to fresh air. Get medical attention if breathing difficulty develops.

IF SWALLOWED, induce vomiting immediately by giving two glasses of water and sticking finger down throat. Get medical attention. NEVER GIVE ANYTHING BY MOUTH BY AN UNCONSCIOUS PERSON.

In case of an emergency involving this product,  
Call Collect, day or night, (314) 694-4000.

### Environmental Hazards

This chemical has been identified in limited ground water sampling and there is the possibility that it can leach through the soil to ground water, especially where soils are coarse and ground water is near the surface. Following application, and during rainfall events that cause runoff, this chemical may reach surface water bodies including streams, rivers and reservoirs.

Care must be taken when using this product to prevent back-siphoning into wells, spills or improper disposal of excess pesticide, spray mixtures or rinsates.

Check valves or antisiphoning devices must be used on all mixing and/or irrigation equipment. Disposal of excess pesticide, spray mixtures or rinsate should be according to label use instructions or according to the state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA regional office.

This product is toxic to fish. Do not apply directly to water or wetlands. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, collect for use or properly dispose of granules.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

### Storage and Disposal

Do not contaminate water, foodstuffs, feed or seed by storage or disposal.

**STORAGE:**  
Keep bag closed to prevent spills and contamination.

**DISPOSAL:**  
Wastes of this pesticide may cause eye burns and may be dangerous. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of according to label use instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Emptied container retains product residue. Observe all labeled safeguards until container is destroyed.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.

(See pamphlet for Complete Directions for Use.)

### LIMIT OF WARRANTY AND LIABILITY

This Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes set forth in the Complete Directions for Use label booklet ("Directions") when used in accordance with those Directions under the conditions described therein. NO OTHER EXPRESS WARRANTY OR IMPLIED WARRANTY OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY IS MADE. This warranty is also subject to the conditions and limitations stated herein.

Buyer and all users shall promptly notify this Company of any claims whether based in contract, negligence, strict liability, other tort or otherwise.

Buyer and all users are responsible for all loss or damage from use or handling which results from conditions beyond the control of this Company, including, but not limited to, incompatibility with products other than those set forth in the Directions, unusual weather, weather conditions which are outside the range considered normal at the application site and for the time period when the product is applied, as well as weather conditions which are outside the application range set forth in the Directions, application in any manner not explicitly set forth in the Directions, moisture conditions outside the moisture range set forth in the Directions, or the presence of products other than those set forth in the Directions in or on the soil, crop or treated vegetation.

THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE LIMIT OF THE LIABILITY OF THIS COMPANY OR ANY OTHER SELLER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT (INCLUDING CLAIMS BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, OTHER TORT OR OTHERWISE) SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER FOR THE QUANTITY OF THIS PRODUCT INVOLVED, OR, AT THE OPTION OF THIS COMPANY OR ANY OTHER SELLER, THE REPLACEMENT OF SUCH QUANTITY, OR, IF NOT REQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. IN NO EVENT SHALL THIS COMPANY OR ANY OTHER SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.

Buyer and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

MONSANTO COMPANY 1989

EPA Reg. No. 524-